



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 02:09 PM GMT

PDB ID : 2QWW
Title : Crystal structure of multiple antibiotic-resistance repressor (MarR)
(YP_013417.1) from *Listeria monocytogenes* 4b F2365 at 2.07 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2007-08-10
Resolution : 2.07 Å (reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1224 (2.08-2.04)
Clashscore	79885	1390 (2.08-2.04)
Ramachandran outliers	78287	1381 (2.08-2.04)
Sidechain outliers	78261	1381 (2.08-2.04)
RSRZ outliers	66119	1225 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	154	
1	B	154	
1	C	154	
1	D	154	
1	E	154	
1	F	154	
1	G	154	
1	H	154	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8738 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator, MarR family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	Se	0	2	0
			1094	696	181	211	6			
1	B	144	Total	C	N	O	Se	0	2	0
			1075	691	174	205	5			
1	C	138	Total	C	N	O	Se	0	4	0
			1041	665	169	203	4			
1	D	139	Total	C	N	O	Se	0	0	0
			1032	658	169	201	4			
1	E	143	Total	C	N	O	Se	0	1	0
			1047	667	173	203	4			
1	F	139	Total	C	N	O	Se	0	0	0
			1032	657	169	201	5			
1	G	140	Total	C	N	O	Se	0	0	0
			1036	662	174	195	5			
1	H	143	Total	C	N	O	Se	0	3	0
			1048	669	173	202	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP Q722C0
B	0	GLY	-	LEADER SEQUENCE	UNP Q722C0
C	0	GLY	-	LEADER SEQUENCE	UNP Q722C0
D	0	GLY	-	LEADER SEQUENCE	UNP Q722C0
E	0	GLY	-	LEADER SEQUENCE	UNP Q722C0
F	0	GLY	-	LEADER SEQUENCE	UNP Q722C0
G	0	GLY	-	LEADER SEQUENCE	UNP Q722C0
H	0	GLY	-	LEADER SEQUENCE	UNP Q722C0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	49	Total 49	O 49	0	0
2	B	45	Total 45	O 45	0	0
2	C	50	Total 50	O 50	0	0
2	D	37	Total 37	O 37	0	0
2	E	41	Total 41	O 41	0	0
2	F	40	Total 40	O 40	0	0
2	G	36	Total 36	O 36	0	0
2	H	35	Total 35	O 35	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcriptional regulator, MarR family

Chain A: 




- Molecule 1: Transcriptional regulator, MarR family

Chain B: 




- Molecule 1: Transcriptional regulator, MarR family

Chain C: 




- Molecule 1: Transcriptional regulator, MarR family

Chain D: 




- Molecule 1: Transcriptional regulator, MarR family

Chain E: 



- Molecule 1: Transcriptional regulator, MarR family


Chain F: 



- Molecule 1: Transcriptional regulator, MarR family



- Molecule 1: Transcriptional regulator, MarR family



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.25Å 159.70Å 90.27Å 90.00° 103.26° 90.00°	Depositor
Resolution (Å)	46.98 – 2.07 46.96 – 2.07	Depositor EDS
% Data completeness (in resolution range)	98.1 (46.98-2.07) 98.1 (46.96-2.07)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 2.07Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.239 , 0.293 0.243 , 0.290	Depositor DCC
R_{free} test set	3967 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	26.4	Xtriage
Anisotropy	0.472	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 60.0	EDS
Estimated twinning fraction	0.037 for h,-k,-h-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 79173 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8738	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.51 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.2438e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	1/1103 (0.1%)	0.63	0/1482
1	B	0.52	0/1081	0.62	0/1455
1	C	0.50	0/1056	0.61	0/1424
1	D	0.48	0/1035	0.59	0/1393
1	E	0.52	0/1052	0.63	0/1418
1	F	0.58	0/1035	0.66	0/1393
1	G	0.50	0/1038	0.61	1/1394 (0.1%)
1	H	0.48	0/1060	0.63	0/1430
All	All	0.51	1/8460 (0.0%)	0.62	1/11389 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	123	MSE	SE-CE	-5.55	1.62	1.95

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	119	MSE	CG-SE-CE	-5.74	86.28	98.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1094	0	1120	8	0
1	B	1075	0	1102	11	0
1	C	1041	0	1067	4	0
1	D	1032	0	1052	9	0
1	E	1047	0	1045	5	0
1	F	1032	0	1053	14	0
1	G	1036	0	1072	11	0
1	H	1048	0	1056	9	0
2	A	49	0	0	1	0
2	B	45	0	0	0	0
2	C	50	0	0	1	0
2	D	37	0	0	0	0
2	E	41	0	0	0	0
2	F	40	0	0	0	0
2	G	36	0	0	1	0
2	H	35	0	0	0	0
All	All	8738	0	8567	62	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:G:25:SER:HA	1:G:119:MSE:HE1	1.61	0.81
1:B:61:THR:HG22	1:B:71:ALA:CB	2.19	0.72
1:D:61:THR:HG22	1:D:71:ALA:CB	2.22	0.70
1:F:61:THR:HG22	1:F:71:ALA:CB	2.23	0.69
1:G:25:SER:CA	1:G:119:MSE:HE1	2.23	0.67

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	144/154 (94%)	142 (99%)	2 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	142/154 (92%)	140 (99%)	2 (1%)	0	100	100
1	C	138/154 (90%)	136 (99%)	2 (1%)	0	100	100
1	D	135/154 (88%)	134 (99%)	1 (1%)	0	100	100
1	E	140/154 (91%)	138 (99%)	2 (1%)	0	100	100
1	F	135/154 (88%)	132 (98%)	2 (2%)	1 (1%)	30	16
1	G	136/154 (88%)	135 (99%)	1 (1%)	0	100	100
1	H	142/154 (92%)	141 (99%)	0	1 (1%)	30	16
All	All	1112/1232 (90%)	1098 (99%)	12 (1%)	2 (0%)	56	47

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	65	ILE
1	F	65	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	116/127 (91%)	114 (98%)	2 (2%)	73	71
1	B	110/127 (87%)	110 (100%)	0	100	100
1	C	111/127 (87%)	111 (100%)	0	100	100
1	D	108/127 (85%)	107 (99%)	1 (1%)	87	88
1	E	105/127 (83%)	105 (100%)	0	100	100
1	F	109/127 (86%)	109 (100%)	0	100	100
1	G	108/127 (85%)	108 (100%)	0	100	100
1	H	107/127 (84%)	107 (100%)	0	100	100
All	All	874/1016 (86%)	871 (100%)	3 (0%)	96	96

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	67	THR
1	D	62	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	C	129	ASN
1	E	22	GLN
1	F	10	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	146/154 (94%)	0.15	2 (1%) 72 75	30, 41, 56, 71	0
1	B	144/154 (93%)	0.18	3 (2%) 60 62	30, 40, 55, 64	0
1	C	138/154 (89%)	0.11	1 (0%) 84 88	30, 40, 54, 57	0
1	D	139/154 (90%)	0.26	4 (2%) 49 50	31, 40, 54, 58	0
1	E	143/154 (92%)	0.18	2 (1%) 72 75	30, 40, 54, 61	0
1	F	139/154 (90%)	0.26	4 (2%) 49 50	29, 40, 54, 63	0
1	G	140/154 (90%)	0.18	2 (1%) 72 75	30, 40, 55, 62	0
1	H	143/154 (92%)	0.29	2 (1%) 72 75	29, 40, 53, 61	0
All	All	1132/1232 (91%)	0.20	20 (1%) 65 68	29, 40, 55, 71	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	87	LEU	3.8
1	H	87	LEU	3.5
1	H	109	LEU	3.5
1	D	6	THR	3.2
1	D	73	ALA	3.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.