



# Full wwPDB X-ray Structure Validation Report

Sep 10, 2014 – 10:54 AM EDT

PDB ID : 4QWC  
Title : TERNARY CRYSTAL STRUCTURES of A Y-FAMILY DNA POLYMERASE DPO4 FROM SULFOLOBUS SOLFATARICUS IN COMPLEX WITH DNA AND L-DCDP  
Authors : Vyas, R.; Suo, Z.  
Deposited on : 2014-07-16  
Resolution : 2.40 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1439  
EDS : stable23489  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable23489

# 1 Overall quality at a glance





The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	343	
1	D	343	
2	B	13	
2	E	13	
3	C	18	
3	F	18	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
7	DOC	B	101	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6888 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	0	0
			2752	1765	473	507	7			
1	D	342	Total	C	N	O	S	0	0	0
			2752	1765	473	507	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	343	ALA	-	EXPRESSION TAG	UNP Q97W02
D	343	ALA	-	EXPRESSION TAG	UNP Q97W02

- Molecule 2 is a DNA chain called DNA (5'-D(\*GP\*GP\*CP\*TP\*AP\*CP\*AP\*GP\*GP\*AP\*CP\*TP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	P	0	0	0
			245	117	48	69	11			
2	E	12	Total	C	N	O	P	0	0	0
			245	117	48	69	11			


- Molecule 3 is a DNA chain called DNA (5'-D(\*TP\*TP\*CP\*AP\*GP\*GP\*AP\*GP\*TP\*CP\*CP\*TP\*GP\*TP\*AP\*GP\*CP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	15	Total	C	N	O	P	0	0	0
			309	146	58	90	15			
3	F	15	Total	C	N	O	P	0	0	0
			309	146	58	90	15			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Ca 2 2	0	0
4	D	3	Total Ca 3 3	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0

- Molecule 6 is 4-AMINO-1-{2-DEOXY-5-O-[(R)-HYDROXY(PHOSPHONOOXY)PHOSPHORYL]-BETA-L-ERYTHRO-PENTOFURANOSYL}PYRIMIDIN-2(1H)-ONE (three-letter code: LTP) (formula:  $C_9H_{15}N_3O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			24	9	3	10	2		
6	D	1	Total	C	N	O	P	0	0
			24	9	3	10	2		

- Molecule 7 is 2',3'-DIDEOXYCYTIDINE-5'-MONOPHOSPHATE (three-letter code: DOC) (formula: C<sub>9</sub>H<sub>14</sub>N<sub>3</sub>O<sub>6</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	B	1	Total	C	N	O	P	0	0
			18	9	3	5	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	E	1	Total	C	N	O	P	0	0
			18	9	3	5	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	80	Total	O	0	0
			80	80		
8	B	9	Total	O	0	0
			9	9		
8	D	57	Total	O	0	0
			57	57		
8	E	5	Total	O	0	0
			5	5		
8	F	10	Total	O	0	0
			10	10		
8	C	14	Total	O	0	0
			14	14		



- Molecule 3: DNA (5'-D(\*TP\*TP\*CP\*AP\*GP\*GP\*AP\*GP\*TP\*CP\*CP\*TP\*GP\*TP\*AP\*GP\*CP\*C)-3')

Chain F: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.44Å 101.93Å 105.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.46 – 2.40 46.46 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.2 (46.46-2.40) 99.2 (46.46-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.50 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.227 , 0.277 0.235 , 0.286	Depositor DCC
$R_{free}$ test set	2110 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	58.0	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.3	EDS
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 41776 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6888	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 73.34 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.8724e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DOC, CA, LTP, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/2791	0.65	0/3748
1	D	0.41	0/2791	0.64	0/3748
2	B	0.35	0/275	0.77	0/423
2	E	0.32	0/275	0.81	0/423
3	C	0.41	0/346	0.85	1/532 (0.2%)
3	F	0.35	0/346	0.84	0/532
All	All	0.42	0/6824	0.69	1/9406 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	8	DG	C1'-O4'-C4'	-5.08	105.02	110.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2752	0	2895	21	0
1	D	2752	0	2895	19	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	245	0	135	7	0
2	E	245	0	135	4	0
3	C	309	0	169	2	0
3	F	309	0	169	4	0
4	A	2	0	0	0	0
4	D	3	0	0	0	0
5	A	8	0	6	3	0
5	D	4	0	3	3	0
6	A	24	0	0	1	0
6	D	24	0	0	0	0
7	B	18	0	12	2	0
7	E	18	0	12	1	0
8	A	80	0	0	5	0
8	B	9	0	0	0	0
8	C	14	0	0	0	0
8	D	57	0	0	3	0
8	E	5	0	0	1	0
8	F	10	0	0	1	0
All	All	6888	0	6431	57	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (57) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
6:A:405:LTP:OBB	8:A:559:HOH:O	1.84	0.94
2:E:4:DT:H2''	2:E:5:DA:H5''	1.52	0.89
2:B:12:DT:H2'	7:B:101:DOC:O4'	1.85	0.77
1:A:181:ALA:O	5:A:404:ACT:C	2.37	0.73
7:E:101:DOC:H2''	8:E:204:HOH:O	1.90	0.71
2:B:6:DC:H2'	2:B:7:DA:C8	2.31	0.65
3:F:9:DT:H2''	3:F:10:DC:C6	2.34	0.63
1:D:298:ARG:HD2	2:E:8:DG:OP1	1.98	0.63
1:A:166:ILE:HG23	1:A:170:GLU:HB3	1.80	0.62
1:D:250:THR:HG22	1:D:332:ARG:HG2	1.83	0.59
1:A:1:MET:HE1	8:A:503:HOH:O	2.02	0.58
1:D:285:HIS:HB3	8:D:537:HOH:O	2.04	0.57
1:A:32:VAL:HG11	3:C:6:DG:H5'	1.86	0.57
1:A:129:LYS:HE3	1:A:163:ILE:HG13	1.89	0.54
1:A:145:SER:OG	1:A:146:LYS:N	2.39	0.54
2:B:12:DT:H2'	7:B:101:DOC:C4'	2.36	0.54
1:D:4:LEU:CD1	1:D:142:VAL:HG13	2.37	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:195:LYS:O	1:A:197:LEU:N	2.40	0.54
1:A:188:ASN:N	1:A:188:ASN:OD1	2.41	0.53
1:D:4:LEU:HD13	1:D:142:VAL:HG13	1.90	0.53
2:B:11:DC:H2'	2:B:12:DT:H72	1.93	0.51
1:A:135:LYS:HD2	8:A:512:HOH:O	2.11	0.50
1:A:289:VAL:HB	1:A:332:ARG:HB2	1.93	0.49
1:D:186:ILE:O	5:D:404:ACT:OXT	2.32	0.48
3:F:5:DG:H1'	3:F:6:DG:H5''	1.95	0.47
1:D:93:ARG:HD2	1:D:99:ILE:HG21	1.97	0.47
1:A:12:TYR:HB2	1:A:45:THR:HG21	1.96	0.46
1:A:257:ASN:HB3	1:A:260:GLU:HG2	1.97	0.46
3:C:12:DT:H2''	3:C:13:DG:C8	2.51	0.46
1:D:112:SER:O	1:D:113:ASP:CB	2.64	0.45
1:D:181:ALA:O	5:D:404:ACT:C	2.64	0.45
2:B:11:DC:H2'	2:B:12:DT:C7	2.47	0.45
1:A:2:ILE:HG12	8:A:537:HOH:O	2.17	0.45
1:A:181:ALA:O	5:A:404:ACT:OXT	2.34	0.45
1:D:12:TYR:HB2	1:D:45:THR:CG2	2.46	0.45
1:A:46:ALA:HB1	1:A:50:ALA:HB3	1.99	0.44
1:D:341:ILE:HG22	1:D:342:GLU:N	2.32	0.44
1:A:12:TYR:HB2	1:A:45:THR:CG2	2.48	0.44
1:D:129:LYS:NZ	8:D:530:HOH:O	2.34	0.43
1:A:9:ASP:OD2	1:A:139:THR:HG23	2.18	0.43
2:B:11:DC:C6	2:B:12:DT:H72	2.54	0.42
2:B:3:DC:H2'	2:B:4:DT:C6	2.54	0.42
1:A:186:ILE:O	5:A:404:ACT:OXT	2.37	0.42
2:E:4:DT:C2'	2:E:5:DA:H5''	2.36	0.42
1:D:166:ILE:HG23	1:D:170:GLU:HB3	2.02	0.42
1:D:46:ALA:HB1	1:D:50:ALA:HB3	2.00	0.42
1:D:10:TYR:HB3	8:D:527:HOH:O	2.20	0.41
1:A:239:THR:HA	8:A:504:HOH:O	2.20	0.41
2:E:11:DC:N3	8:F:104:HOH:O	2.51	0.41
1:D:45:THR:OG1	1:D:46:ALA:N	2.53	0.41
1:D:181:ALA:O	5:D:404:ACT:OXT	2.39	0.41
1:D:247:ARG:NE	1:D:271:GLU:OE1	2.51	0.40
1:D:326:ASP:OD1	1:D:328:ARG:HG2	2.21	0.40
1:A:112:SER:O	1:A:113:ASP:HB2	2.21	0.40
1:A:282:LYS:HD2	1:A:341:ILE:HD13	2.04	0.40
3:F:11:DC:H2'	3:F:12:DT:C6	2.56	0.40
3:F:5:DG:C2'	3:F:6:DG:H5''	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/343 (99%)	328 (96%)	10 (3%)	2 (1%)	33	47
1	D	340/343 (99%)	328 (96%)	10 (3%)	2 (1%)	33	47
All	All	680/686 (99%)	656 (96%)	20 (3%)	4 (1%)	33	47

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	196	LYS
1	A	197	LEU
1	D	292	ASP
1	D	113	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/301 (100%)	279 (93%)	22 (7%)	20	30
1	D	301/301 (100%)	279 (93%)	22 (7%)	20	30
All	All	602/602 (100%)	558 (93%)	44 (7%)	20	30

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	VAL
1	A	17	GLU
1	A	20	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	55	VAL
1	A	63	GLU
1	A	76	MET
1	A	124	LEU
1	A	133	LEU
1	A	163	ILE
1	A	165	VAL
1	A	170	GLU
1	A	174	LEU
1	A	176	ARG
1	A	188	ASN
1	A	202	LEU
1	A	216	MET
1	A	235	GLU
1	A	247	ARG
1	A	248	ILE
1	A	253	ARG
1	A	276	LEU
1	A	323	LEU
1	D	4	LEU
1	D	23	LEU
1	D	26	LYS
1	D	55	VAL
1	D	76	MET
1	D	93	ARG
1	D	124	LEU
1	D	163	ILE
1	D	170	GLU
1	D	174	LEU
1	D	197	LEU
1	D	200	ASN
1	D	202	LEU
1	D	241	VAL
1	D	247	ARG
1	D	278	LYS
1	D	279	ARG
1	D	282	LYS
1	D	297	SER
1	D	298	ARG
1	D	328	ARG
1	D	335	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	D	20	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 5 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	ACT	A	403	-	1,3,3	1.90	0	0,3,3	0.00	-
5	ACT	A	404	-	1,3,3	1.43	0	0,3,3	0.00	-
6	LTP	A	405	4	25,25,25	2.20	4 (16%)	35,38,38	2.08	6 (17%)
7	DOC	B	101	2	17,19,20	1.58	3 (17%)	20,26,29	1.74	4 (20%)
5	ACT	D	404	-	1,3,3	0.94	0	0,3,3	0.00	-
6	LTP	D	405	4	25,25,25	2.06	4 (16%)	35,38,38	1.89	6 (17%)
7	DOC	E	101	2	17,19,20	1.73	3 (17%)	20,26,29	2.05	8 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ACT	A	403	-	-	0/0/0/0	0/0/0/0
5	ACT	A	404	-	-	0/0/0/0	0/0/0/0
6	LTP	A	405	4	-	0/13/28/28	0/2/2/2
7	DOC	B	101	2	-	0/5/18/19	0/2/2/2
5	ACT	D	404	-	-	0/0/0/0	0/0/0/0
6	LTP	D	405	4	-	0/13/28/28	0/2/2/2
7	DOC	E	101	2	-	0/5/18/19	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	405	LTP	C6-N1	8.48	1.48	1.35
6	D	405	LTP	C6-N1	7.85	1.47	1.35
7	B	101	DOC	P-OP1	4.80	1.52	1.46
7	E	101	DOC	C2-N1	4.27	1.43	1.38
7	E	101	DOC	P-OP1	4.23	1.51	1.46
6	A	405	LTP	O2-C2	4.18	1.27	1.21
6	A	405	LTP	C6-C5	3.88	1.46	1.38
6	D	405	LTP	C6-C5	3.77	1.46	1.38
6	D	405	LTP	O2-C2	3.76	1.26	1.21
6	D	405	LTP	C2-N1	-3.06	1.34	1.38
7	E	101	DOC	O2-C2	3.03	1.25	1.21
7	B	101	DOC	O2-C2	2.97	1.25	1.21
6	A	405	LTP	C2-N1	-2.86	1.35	1.38
7	B	101	DOC	C2-N1	2.58	1.41	1.38

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	405	LTP	C6-C5-C4	6.37	120.55	117.51
6	A	405	LTP	C2-N3-C4	5.80	124.00	115.65
6	A	405	LTP	C6-C5-C4	5.34	120.06	117.51
6	A	405	LTP	PAI-OAH-PAF	-5.33	117.16	131.93
6	D	405	LTP	C2-N3-C4	5.13	123.04	115.65
6	D	405	LTP	PAI-OAH-PAF	-4.02	120.80	131.93
7	E	101	DOC	C6-C5-C4	4.01	119.42	117.51
6	A	405	LTP	CAQ-CAR-N1	-4.01	104.13	114.14
7	B	101	DOC	C6-C5-C4	3.86	119.36	117.51
7	E	101	DOC	C4'-O4'-C1'	3.83	111.64	110.05
7	B	101	DOC	C2-N3-C4	3.77	121.07	115.65

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	E	101	DOC	C2-N3-C4	3.37	120.49	115.65
6	A	405	LTP	C5-C6-N1	-3.26	112.93	120.68
6	A	405	LTP	OAO-CAR-N1	3.15	113.34	107.66
6	D	405	LTP	C5-C6-N1	-3.04	113.44	120.68
7	B	101	DOC	C3'-C2'-C1'	3.04	106.17	102.80
6	D	405	LTP	CAQ-CAR-N1	-2.90	106.90	114.14
7	B	101	DOC	C4'-O4'-C1'	2.89	111.25	110.05
7	E	101	DOC	C3'-C2'-C1'	2.88	105.99	102.80
7	E	101	DOC	O4'-C4'-C5'	2.38	113.34	109.52
7	E	101	DOC	O4'-C1'-N1	2.33	111.86	107.66
6	D	405	LTP	OAO-CAR-N1	2.28	111.76	107.66
7	E	101	DOC	C2'-C3'-C4'	2.13	106.84	102.75
7	E	101	DOC	C2-N1-C1'	2.03	121.47	118.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	342/343 (99%)	0.16	7 (2%) 62 59	28, 61, 89, 112	0
1	D	342/343 (99%)	0.25	3 (0%) 81 81	35, 69, 103, 124	0
2	B	12/13 (92%)	0.06	0 100 100	55, 69, 92, 92	0
2	E	12/13 (92%)	0.33	1 (8%) 11 10	74, 87, 94, 98	0
3	C	15/18 (83%)	0.09	0 100 100	53, 74, 90, 139	0
3	F	15/18 (83%)	0.39	1 (6%) 17 16	64, 95, 117, 149	0
All	All	738/748 (98%)	0.20	12 (1%) 68 67	28, 66, 101, 149	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	37	PHE	3.1
1	A	292	ASP	2.8
1	A	212	LYS	2.6
1	A	295	ILE	2.5
1	D	293	LEU	2.5
1	D	163	ILE	2.4
1	A	293	LEU	2.1
1	A	216	MET	2.1
1	D	322	ILE	2.1
3	F	4	DA	2.1
1	A	207	SER	2.0
2	E	9	DG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
7	DOC	B	101	18/19	0.26	6.41	75,82,88,88	4
5	ACT	A	403	4/4	0.29	1.11	78,83,84,85	0
6	LTP	A	405	24/24	0.16	0.52	47,57,73,79	0
7	DOC	E	101	18/19	0.15	0.48	77,84,91,92	0
5	ACT	A	404	4/4	0.16	0.47	42,46,46,52	0
6	LTP	D	405	24/24	0.17	0.43	58,65,70,71	0
5	ACT	D	404	4/4	0.13	-2.14	59,62,64,65	0
4	CA	A	401	1/1	0.09	-2.51	43,43,43,43	0
4	CA	A	402	1/1	0.09	-2.71	88,88,88,88	0
4	CA	D	401	1/1	0.06	-3.15	96,96,96,96	0
4	CA	D	403	1/1	0.10	-3.50	52,52,52,52	0
4	CA	D	402	1/1	0.08	-8.30	86,86,86,86	0

### 6.5 Other polymers

There are no such residues in this entry.