



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 10:38 AM GMT

PDB ID : 2QX0
Title : Crystal Structure of Yersinia pestis HPPK (Ternary Complex)
Authors : Blaszczyk, J.; Cherry, S.; Tropea, J.E.; Waugh, D.S.; Ji, X.
Deposited on : 2007-08-10
Resolution : 1.80 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3513 (1.80-1.80)
Clashscore	79885	4461 (1.80-1.80)
Ramachandran outliers	78287	4404 (1.80-1.80)
Sidechain outliers	78261	4403 (1.80-1.80)
RSRZ outliers	66119	3515 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	159	
1	B	159	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	PH2	A	181	X	-
4	PH2	B	182	X	-

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 3089 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 7,8-dihydro-6-hydroxymethylpterin-pyrophosphokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	159	Total	C	N	O	S	0	0	0
			1265	804	225	230	6			
1	B	159	Total	C	N	O	S	0	0	0
			1265	804	225	230	6			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mg	0	0
			2	2		
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is DIPHOSPHOMETHYLPHOSPHONICACID ADENOSYL ESTER (three-letter code: APC) (formula: C₁₁H₁₈N₅O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
3	B	1	Total	C	N	O	P	0	0
			31	11	5	12	3		

- Molecule 4 is 2-AMINO-6-HYDROXYMETHYL-7,8-DIHYDRO-3H-PTERIDIN-4-ONE (three-letter code: PH2) (formula: C₇H₉N₅O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	7	5	2		
4	B	1	Total	C	N	O	0	0
			14	7	5	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	240	Total	O	0	0
			240	240		
5	B	225	Total	O	0	0
			225	225		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 7,8-dihydro-6-hydroxymethylpterin-pyrophosphokinase



- Molecule 1: 7,8-dihydro-6-hydroxymethylpterin-pyrophosphokinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.40Å 78.62Å 52.60Å 90.00° 100.72° 90.00°	Depositor
Resolution (Å)	28.90 – 1.80 28.90 – 1.74	Depositor EDS
% Data completeness (in resolution range)	95.7 (28.90-1.80) 94.1 (28.90-1.74)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 1.74Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.230 , 0.269 0.231 , 0.268	Depositor DCC
R_{free} test set	1525 reflections (4.95%)	DCC
Wilson B-factor (Å ²)	14.8	Xtriage
Anisotropy	0.522	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 33783 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3089	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.65 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.5366e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PH2, CSO, MG, APC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/1284	0.64	0/1743
1	B	0.30	0/1284	0.65	0/1743
All	All	0.30	0/2568	0.64	0/3486

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1265	0	1294	40	0
1	B	1265	0	1294	42	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	31	0	14	0	0
3	B	31	0	14	0	0
4	A	14	0	8	1	0
4	B	14	0	8	0	0
5	A	240	0	0	16	0
5	B	225	0	0	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3089	0	2632	83	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

All (83) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:100:MET:HG2	1:B:125:MET:HE3	1.47	0.97
1:A:71:LEU:HD22	1:A:112:LEU:HD13	1.63	0.78
1:B:100:MET:HG2	1:B:125:MET:CE	2.14	0.77
1:B:27:ALA:HB2	1:B:34:LEU:HD12	1.68	0.76
1:A:27:ALA:HB2	1:A:34:LEU:HD12	1.70	0.73
1:A:100:MET:HG2	1:A:125:MET:HE3	1.69	0.72
1:A:37:CSO:SG	5:A:205:HOH:O	2.47	0.71
1:A:100:MET:HG2	1:A:125:MET:CE	2.20	0.70
1:B:40:LEU:HD22	1:B:55:LEU:HD21	1.78	0.65
1:A:1:MET:HE3	5:A:383:HOH:O	1.96	0.65
1:B:123:GLU:HG3	1:B:153:LYS:O	2.02	0.60
1:A:58:VAL:HG23	1:A:128:PRO:HB2	1.84	0.59
1:B:19:VAL:HG21	1:B:55:LEU:HD22	1.86	0.56
1:B:17:GLN:HB3	5:B:402:HOH:O	2.05	0.56
1:B:119:LEU:HG	1:B:125:MET:HE2	1.87	0.56
1:B:80:ASN:O	1:B:84:VAL:HG23	2.06	0.56
1:B:34:LEU:HD21	1:B:37:CSO:OD	2.08	0.54
1:A:34:LEU:HD23	1:A:35:VAL:N	2.22	0.54
1:A:119:LEU:HD13	1:A:125:MET:HE1	1.90	0.54
1:B:79:ARG:O	1:B:83:ARG:HG2	2.07	0.54
1:B:1:MET:HG3	5:B:387:HOH:O	2.06	0.54
1:A:41:TYR:CD2	1:A:128:PRO:HG3	2.43	0.54
1:B:58:VAL:HG23	1:B:128:PRO:HB2	1.91	0.52
1:A:58:VAL:CG2	1:A:128:PRO:HB2	2.39	0.52
1:B:156:LEU:HD23	1:B:156:LEU:C	2.30	0.52
1:A:119:LEU:HD13	1:A:125:MET:CE	2.39	0.52
1:B:158:LEU:HD13	5:B:257:HOH:O	2.10	0.51
1:A:123:GLU:HG3	1:A:153:LYS:O	2.11	0.51
1:B:6:ILE:HD13	1:B:99:ILE:HA	1.92	0.51
1:B:51:GLN:HB2	1:B:52:PRO:CD	2.41	0.51
1:B:34:LEU:HD23	1:B:35:VAL:N	2.26	0.50
1:B:136:LEU:HD23	1:B:137:ILE:N	2.27	0.50
1:A:81:GLN:NE2	1:A:95:LEU:HD22	2.28	0.49
1:A:12:LEU:HD23	1:A:92:PRO:HB2	1.93	0.49
1:A:106:VAL:HG13	5:A:354:HOH:O	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:43:THR:HG21	1:B:124:PHE:HB3	1.94	0.49
1:A:23:ARG:HB3	5:A:225:HOH:O	2.12	0.49
1:B:127:TYR:HB2	1:B:128:PRO:HD3	1.95	0.48
1:B:41:TYR:CD2	1:B:128:PRO:HG3	2.49	0.47
1:A:113:THR:HG23	1:A:113:THR:O	2.15	0.47
1:A:23:ARG:NH2	5:A:219:HOH:O	2.37	0.47
1:B:41:TYR:CD1	1:B:156:LEU:HD21	2.49	0.47
1:A:51:GLN:HB2	1:A:52:PRO:CD	2.44	0.47
1:A:122:ARG:HD3	1:A:124:PHE:CZ	2.50	0.47
1:B:58:VAL:CG2	1:B:128:PRO:HB2	2.45	0.47
1:A:34:LEU:HG	5:A:255:HOH:O	2.14	0.47
1:B:138:PHE:HE2	1:B:147:CYS:HG	1.62	0.46
5:A:278:HOH:O	1:B:108:LYS:HE2	2.15	0.46
1:B:113:THR:O	1:B:113:THR:HG23	2.16	0.46
1:A:111:ARG:HD3	5:A:324:HOH:O	2.14	0.46
1:A:13:ALA:HB3	5:A:387:HOH:O	2.16	0.46
4:A:181:PH2:H112	5:A:316:HOH:O	2.15	0.46
1:A:114:ILE:HA	1:A:115:PRO:C	2.35	0.45
5:A:327:HOH:O	1:B:108:LYS:HG2	2.16	0.45
1:B:11:ASN:O	1:B:52:PRO:HB2	2.18	0.44
1:A:79:ARG:HD2	5:A:360:HOH:O	2.17	0.44
1:B:49:GLN:HB2	5:B:298:HOH:O	2.17	0.44
1:A:61:LEU:N	1:A:61:LEU:HD23	2.33	0.44
1:A:100:MET:HE2	1:A:100:MET:HB3	1.83	0.44
1:B:23:ARG:HB3	5:B:243:HOH:O	2.17	0.44
1:B:34:LEU:HA	1:B:61:LEU:HD12	2.00	0.44
1:B:23:ARG:HD3	5:B:243:HOH:O	2.18	0.44
1:A:16:LEU:HD21	1:A:159:TRP:CD1	2.53	0.44
1:A:19:VAL:HG21	1:A:55:LEU:HG	2.01	0.43
1:B:15:PRO:HG2	1:B:55:LEU:HD12	1.99	0.43
1:A:127:TYR:HB2	1:A:128:PRO:HD3	2.01	0.43
1:B:40:LEU:HD23	1:B:57:ALA:HB2	2.01	0.42
1:B:107:ILE:HB	1:B:114:ILE:HB	2.01	0.42
1:A:120:LYS:HB2	5:A:281:HOH:O	2.18	0.42
1:B:81:GLN:NE2	1:B:95:LEU:HD22	2.35	0.42
1:A:16:LEU:HD21	1:A:159:TRP:NE1	2.35	0.42
1:A:100:MET:HG2	1:A:125:MET:HE2	2.00	0.42
1:B:114:ILE:HA	1:B:115:PRO:C	2.40	0.41
1:B:71:LEU:O	1:B:75:GLN:HG3	2.20	0.41
1:A:16:LEU:HB2	5:A:247:HOH:O	2.18	0.41
1:A:158:LEU:HD22	5:A:329:HOH:O	2.20	0.41
1:B:12:LEU:O	1:B:15:PRO:HD3	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:11:ASN:HB3	1:B:54:PHE:CD1	2.55	0.41
1:A:145:SER:HB3	5:A:386:HOH:O	2.19	0.41
1:A:71:LEU:O	1:A:75:GLN:HG3	2.20	0.41
1:A:136:LEU:HD23	1:A:137:ILE:N	2.37	0.40
1:B:100:MET:HB3	1:B:100:MET:HE2	1.97	0.40
1:A:11:ASN:HB3	1:A:54:PHE:CG	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/159 (98%)	155 (99%)	1 (1%)	0	100	100
1	B	156/159 (98%)	156 (100%)	0	0	100	100
All	All	312/318 (98%)	311 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/137 (100%)	136 (99%)	1 (1%)	91	88
1	B	137/137 (100%)	135 (98%)	2 (2%)	76	66
All	All	274/274 (100%)	271 (99%)	3 (1%)	84	77

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	LEU
1	B	16	LEU
1	B	119	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	73	HIS
1	A	81	GLN
1	A	105	GLN
1	B	17	GLN
1	B	18	GLN
1	B	81	GLN
1	B	105	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	A	37	1	6,6,7	8.13	3 (50%)	3,6,8	1.76	1 (33%)
1	CSO	B	37	1	6,6,7	7.81	3 (50%)	3,6,8	1.82	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	37	1	-	0/2/5/7	0/0/0/0
1	CSO	B	37	1	-	0/2/5/7	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	37	CSO	O-C	19.07	1.24	1.11
1	B	37	CSO	O-C	18.49	1.24	1.11
1	B	37	CSO	OD-SG	4.02	1.79	1.62
1	A	37	CSO	OD-SG	4.00	1.79	1.62
1	A	37	CSO	CB-SG	-3.58	1.78	1.82
1	B	37	CSO	CB-SG	-2.01	1.80	1.82

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	37	CSO	C-CA-N	-2.96	110.87	113.83
1	B	37	CSO	C-CA-N	-2.95	110.88	113.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	APC	A	171	2	33,33,33	2.01	10 (30%)	52,52,52	1.92	10 (19%)
4	PH2	A	181	2	15,15,15	3.47	7 (46%)	19,21,21	4.65	13 (68%)
3	APC	B	172	2	33,33,33	2.00	7 (21%)	52,52,52	1.89	10 (19%)
4	PH2	B	182	2	15,15,15	3.40	9 (60%)	19,21,21	4.69	13 (68%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	APC	A	171	2	-	0/20/38/38	0/1/3/3
4	PH2	A	181	2	-	1/1/11/11	0/0/2/2
3	APC	B	172	2	-	0/20/38/38	0/1/3/3
4	PH2	B	182	2	-	1/1/11/11	0/0/2/2

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	181	PH2	C6-N7	7.46	1.48	1.36
4	B	182	PH2	C6-N7	7.04	1.48	1.36
3	B	172	APC	C8-N9	5.94	1.45	1.36
4	A	181	PH2	C3-N4	-5.92	1.33	1.45
4	B	182	PH2	C3-N4	-5.60	1.33	1.45
3	A	171	APC	C8-N9	5.36	1.44	1.36
3	A	171	APC	C2-N3	5.33	1.42	1.32
3	B	172	APC	C2-N3	5.32	1.42	1.32
4	B	182	PH2	C6-N6	-4.73	1.25	1.32
4	A	181	PH2	C6-N6	-4.38	1.25	1.32
4	A	181	PH2	C3-C2	4.21	1.57	1.49
4	B	182	PH2	C6-N5	-4.18	1.28	1.33
4	B	182	PH2	C3-C2	4.15	1.56	1.49
4	A	181	PH2	C6-N5	-4.14	1.28	1.33
3	A	171	APC	O4'-C1'	3.92	1.47	1.41
3	B	172	APC	O4'-C1'	3.79	1.47	1.41
4	A	181	PH2	C9-C10	3.68	1.52	1.42
3	A	171	APC	C4-N3	3.65	1.41	1.35
4	B	182	PH2	C9-C10	3.61	1.52	1.42
3	B	172	APC	C4-N3	3.31	1.40	1.35
4	A	181	PH2	C8-C9	-3.08	1.36	1.42
4	B	182	PH2	C8-C9	-2.84	1.37	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	172	APC	PA-O2A	-2.75	1.49	1.56
3	B	172	APC	PB-O2B	-2.72	1.49	1.56
3	A	171	APC	PB-O2B	-2.60	1.49	1.56
3	A	171	APC	PA-C3A	-2.49	1.77	1.79
3	A	171	APC	O3'-C3'	-2.41	1.37	1.43
3	A	171	APC	PA-O2A	-2.37	1.50	1.56
4	B	182	PH2	C2-N1	2.24	1.31	1.28
3	A	171	APC	PG-O3B	2.07	1.63	1.60
3	A	171	APC	C4-N9	-2.06	1.34	1.37
3	B	172	APC	O3'-C3'	-2.04	1.38	1.43
4	B	182	PH2	O8-C8	2.03	1.28	1.24

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	182	PH2	C3-N4-C10	10.29	133.06	121.36
4	A	181	PH2	C3-N4-C10	10.19	132.94	121.36
4	B	182	PH2	N4-C10-N5	8.55	128.36	115.82
4	A	181	PH2	N4-C10-N5	8.49	128.27	115.82
4	B	182	PH2	N6-C6-N5	7.71	130.72	120.31
4	A	181	PH2	N6-C6-N5	7.66	130.65	120.31
3	A	171	APC	C4'-O4'-C1'	-7.32	101.80	109.75
3	B	172	APC	C4'-O4'-C1'	-7.26	101.86	109.75
4	B	182	PH2	C6-N5-C10	7.15	127.78	117.61
4	A	181	PH2	C6-N5-C10	7.12	127.73	117.61
4	A	181	PH2	N7-C6-N5	-5.55	114.01	121.78
4	B	182	PH2	N7-C6-N5	-5.42	114.18	121.78
4	A	181	PH2	C9-C10-N4	-5.19	113.12	119.17
4	B	182	PH2	C9-C10-N4	-5.11	113.21	119.17
4	B	182	PH2	C9-N1-C2	4.35	122.40	117.95
4	A	181	PH2	C9-N1-C2	4.08	122.13	117.95
3	B	172	APC	PA-C3A-PB	-3.95	111.94	117.62
3	A	171	APC	PA-C3A-PB	-3.83	112.11	117.62
3	B	172	APC	C5-C6-N6	3.74	129.17	120.72
3	A	171	APC	O5'-PA-O1A	-3.73	104.88	114.21
4	B	182	PH2	C10-C9-N1	-3.59	119.49	123.15
3	A	171	APC	C5-C6-N6	3.57	128.79	120.72
3	B	172	APC	C8-N9-C1'	-3.49	119.50	126.38
4	A	181	PH2	C10-C9-N1	-3.43	119.65	123.15
3	A	171	APC	C8-N9-C1'	-3.43	119.63	126.38
3	B	172	APC	C3'-C2'-C1'	3.34	106.13	100.91
3	B	172	APC	C1'-N9-C4	3.32	132.36	126.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	172	APC	O5'-PA-O1A	-3.23	106.14	114.21
3	A	171	APC	C3'-C2'-C1'	3.17	105.88	100.91
3	A	171	APC	C1'-N9-C4	3.13	132.04	126.64
3	A	171	APC	O3B-PB-O1B	-3.03	105.10	111.51
4	A	181	PH2	C8-C9-C10	2.79	116.46	114.55
4	B	182	PH2	C8-C9-N1	2.69	124.16	120.39
3	B	172	APC	N6-C6-N1	-2.64	114.18	119.36
4	B	182	PH2	C8-C9-C10	2.62	116.34	114.55
3	A	171	APC	O2B-PB-O1B	2.59	118.99	110.93
3	A	171	APC	N6-C6-N1	-2.52	114.42	119.36
4	B	182	PH2	N6-C6-N7	-2.50	115.10	117.86
4	A	181	PH2	C8-N7-C6	2.50	123.88	119.51
4	B	182	PH2	C8-N7-C6	2.50	123.88	119.51
4	A	181	PH2	C8-C9-N1	2.49	123.89	120.39
3	B	172	APC	O2B-PB-O1B	2.39	118.36	110.93
3	B	172	APC	O3B-PB-O1B	-2.32	106.59	111.51
4	A	181	PH2	N6-C6-N7	-2.29	115.34	117.86
4	B	182	PH2	O4-C11-C2	2.15	116.48	110.83
4	A	181	PH2	O4-C11-C2	2.15	116.48	110.83

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	181	PH2	O4-C11-C2-C3
4	B	182	PH2	O4-C11-C2-C3

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	159/159 (100%)	1.07	17 (10%) 6 5	10, 19, 41, 60	0
1	B	159/159 (100%)	0.95	18 (11%) 6 4	9, 18, 39, 57	0
All	All	318/318 (100%)	1.01	35 (11%) 6 5	9, 18, 39, 60	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	48	PRO	7.0
1	A	84	VAL	4.5
1	A	49	GLN	4.3
1	B	48	PRO	4.0
1	A	50	ASP	4.0
1	A	34	LEU	3.7
1	A	83	ARG	3.7
1	A	104	ASP	3.6
1	B	34	LEU	3.5
1	B	159	TRP	3.2
1	B	50	ASP	3.0
1	B	17	GLN	2.9
1	B	104	ASP	2.9
1	A	146	GLU	2.8
1	B	158	LEU	2.8
1	B	49	GLN	2.8
1	A	86	LYS	2.8
1	A	158	LEU	2.7
1	B	14	MET	2.7
1	B	132	ILE	2.6
1	A	159	TRP	2.5
1	B	58	VAL	2.5
1	B	147	CYS	2.4
1	B	146	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2	ILE	2.3
1	A	152	ASP	2.2
1	A	145	SER	2.2
1	B	7	ALA	2.2
1	B	21	ALA	2.2
1	B	156	LEU	2.2
1	B	37	CSO	2.1
1	A	24	GLU	2.1
1	B	20	SER	2.1
1	A	58	VAL	2.1
1	A	14	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CSO	B	37	7/8	0.18	0.00	12,12,25,26	0
1	CSO	A	37	7/8	0.16	-0.56	10,14,23,26	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MG	B	163	1/1	0.14	-0.07	14,14,14,14	0
4	PH2	A	181	14/14	0.14	-0.43	11,13,14,14	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	PH2	B	182	14/14	0.13	-0.44	11,13,15,16	0
2	MG	A	161	1/1	0.13	-0.96	14,14,14,14	0
2	MG	A	162	1/1	0.12	-1.22	13,13,13,13	0
3	APC	A	171	31/31	0.12	-1.54	4,10,12,13	0
2	MG	B	164	1/1	0.10	-1.64	9,9,9,9	0
3	APC	B	172	31/31	0.11	-1.88	5,9,12,12	0

6.5 Other polymers ⓘ

There are no such residues in this entry.