



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 12:39 PM GMT

PDB ID : 1R0L
Title : 1-deoxy-D-xylulose 5-phosphate reductoisomerase from zymomonas mobilis in complex with NADPH
Authors : Ricagno, S.; Grolle, S.; Bringer-Meyer, S.; Sahm, H.; Lindqvist, Y.; Schneider, G.
Deposited on : 2003-09-22
Resolution : 2.70 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1557 (2.70-2.70)
Clashscore	79885	1939 (2.70-2.70)
Ramachandran outliers	78287	1905 (2.70-2.70)
Sidechain outliers	78261	1905 (2.70-2.70)
RSRZ outliers	66119	1559 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	388	
1	B	388	
1	C	388	
1	D	388	

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 11652 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-deoxy-D-xylulose 5-phosphate reductoisomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	378	Total	C	N	O	S	0	0	0
			2854	1798	496	544	16			
1	B	379	Total	C	N	O	S	0	0	0
			2867	1807	499	545	16			
1	C	378	Total	C	N	O	S	0	0	0
			2855	1799	495	544	17			
1	D	378	Total	C	N	O	S	0	0	0
			2860	1802	498	544	16			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	30	Total	O	0	0
			30	30		
3	B	36	Total	O	0	0
			36	36		
3	C	22	Total	O	0	0
			22	22		
3	D	20	Total	O	0	0
			20	20		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 1-deoxy-D-xylulose 5-phosphate reductoisomerase

Chain A:




- Molecule 1: 1-deoxy-D-xylulose 5-phosphate reductoisomerase

Chain B:




• Molecule 1: 1-deoxy-D-xylulose 5-phosphate reductoisomerase

Chain C:



• Molecule 1: 1-deoxy-D-xylulose 5-phosphate reductoisomerase

Chain D:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.70Å 93.20Å 98.60Å 90.00° 90.50° 90.00°	Depositor
Resolution (Å)	29.63 – 2.70 29.63 – 2.69	Depositor EDS
% Data completeness (in resolution range)	98.9 (29.63-2.70) 98.3 (29.63-2.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.68Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.249 , 0.269 0.361 , 0.363	Depositor DCC
R_{free} test set	492 reflections (1.14%)	DCC
Wilson B-factor (Å ²)	42.4	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 15.3	EDS
Estimated twinning fraction	0.000 for h,-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	3 of 43474 reflections (0.007%)	Xtriage
F_o, F_c correlation	0.82	EDS
Total number of atoms	11652	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.88 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.9583e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/2901	0.65	0/3933
1	B	0.46	1/2916 (0.0%)	0.73	2/3955 (0.1%)
1	C	0.41	0/2902	0.62	0/3934
1	D	0.41	0/2908	0.64	2/3943 (0.1%)
All	All	0.43	1/11627 (0.0%)	0.66	4/15765 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	199	HIS	C-O	-5.95	1.12	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	199	HIS	C-N-CD	-17.40	82.32	120.60
1	B	199	HIS	C-N-CA	6.81	150.59	122.00
1	D	40	ARG	NE-CZ-NH1	-5.19	117.70	120.30
1	D	126	GLU	CA-CB-CG	-5.09	102.19	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2854	0	2891	113	0
1	B	2867	0	2902	98	14
1	C	2855	0	2897	103	15
1	D	2860	0	2895	101	0
2	A	27	0	11	1	0
2	B	27	0	11	3	0
2	C	27	0	11	1	0
2	D	27	0	11	3	0
3	A	30	0	0	3	0
3	B	36	0	0	6	1
3	C	22	0	0	4	0
3	D	20	0	0	2	0
All	All	11652	0	11629	407	15

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 18.

The worst 5 of 407 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:125:LYS:HB2	1:C:224:GLU:OE2	1.50	1.11
1:D:40:ARG:HG2	1:D:64:LEU:HD11	1.48	0.95
1:C:126:GLU:OE1	1:C:126:GLU:HA	1.74	0.87
1:A:3:GLN:HB3	1:A:4:PRO:HD3	1.59	0.84
1:A:247:HIS:HD2	1:A:260:GLN:HE22	1.25	0.82

The worst 5 of 15 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:164:ASN:N	1:C:1:MET:O[2_656]	0.73	1.47
1:B:164:ASN:N	1:C:1:MET:C[2_656]	1.46	0.74
1:B:163:HIS:C	1:C:1:MET:C[2_656]	1.50	0.70
1:B:164:ASN:CA	1:C:1:MET:O[2_656]	1.55	0.65
1:B:163:HIS:N	1:C:1:MET:CA[2_656]	1.59	0.61

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/388 (96%)	355 (95%)	17 (4%)	2 (0%)	38	70
1	B	375/388 (97%)	355 (95%)	19 (5%)	1 (0%)	50	82
1	C	374/388 (96%)	353 (94%)	19 (5%)	2 (0%)	38	70
1	D	374/388 (96%)	360 (96%)	12 (3%)	2 (0%)	38	70
All	All	1497/1552 (96%)	1423 (95%)	67 (4%)	7 (0%)	38	70

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	248	SER
1	B	248	SER
1	C	248	SER
1	D	248	SER
1	D	293	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/308 (97%)	281 (94%)	18 (6%)	27	56
1	B	301/308 (98%)	287 (95%)	14 (5%)	36	69
1	C	300/308 (97%)	292 (97%)	8 (3%)	57	87
1	D	300/308 (97%)	288 (96%)	12 (4%)	42	75
All	All	1200/1232 (97%)	1148 (96%)	52 (4%)	40	72

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	151	SER
1	B	287	LEU
1	D	260	GLN
1	B	166	ASP
1	B	234	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	294	GLN
1	C	260	GLN
1	D	142	HIS
1	B	272	HIS
1	D	153	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	A	400	-	29,29,52	1.64	9 (31%)	45,45,80	2.87	18 (40%)
2	NDP	B	401	-	29,29,52	1.65	6 (20%)	45,45,80	2.86	17 (37%)
2	NDP	C	402	-	29,29,52	1.47	3 (10%)	45,45,80	1.81	10 (22%)
2	NDP	D	403	-	29,29,52	3.24	12 (41%)	45,45,80	3.87	26 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	400	-	-	0/15/31/77	0/1/3/5
2	NDP	B	401	-	-	0/15/31/77	0/1/3/5
2	NDP	C	402	-	-	0/15/31/77	0/1/3/5
2	NDP	D	403	-	-	0/15/31/77	0/1/3/5

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	403	NDP	O4B-C1B	8.13	1.53	1.41
2	D	403	NDP	O5B-C5B	-7.24	1.14	1.44
2	D	403	NDP	C3B-C4B	-6.72	1.34	1.53
2	D	403	NDP	C5B-C4B	-5.60	1.33	1.51
2	D	403	NDP	C8A-N7A	-5.52	1.23	1.34

The worst 5 of 71 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	403	NDP	O2A-PA-O3	-10.36	69.50	110.73
2	B	401	NDP	O3X-P2B-O1X	-8.96	81.14	110.44
2	A	400	NDP	O3X-P2B-O1X	-8.92	81.30	110.44
2	D	403	NDP	O3X-P2B-O1X	-8.37	83.07	110.44
2	A	400	NDP	O3X-P2B-O2B	-8.31	83.15	107.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	378/388 (97%)	1.72	131 (34%) 1 1	16, 43, 87, 104	0
1	B	379/388 (97%)	1.06	56 (14%) 3 3	17, 35, 63, 88	0
1	C	378/388 (97%)	1.85	137 (36%) 1 1	22, 44, 69, 94	0
1	D	378/388 (97%)	1.45	88 (23%) 1 1	22, 45, 74, 107	0
All	All	1513/1552 (97%)	1.52	412 (27%) 1 1	16, 41, 75, 107	0

The worst 5 of 412 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	MET	9.9
1	C	2	SER	9.0
1	C	68	LEU	8.9
1	A	365	LEU	7.7
1	C	365	LEU	7.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NDP	C	402	27/48	0.30	-0.12	74,76,99,101	0
2	NDP	B	401	27/48	0.20	-0.48	41,53,88,96	0
2	NDP	A	400	27/48	0.25	-0.70	61,63,89,95	0
2	NDP	D	403	27/48	0.20	-0.82	51,55,75,78	0

6.5 Other polymers ⓘ

There are no such residues in this entry.