



Full wwPDB X-ray Structure Validation Report

Aug 18, 2014 – 05:27 PM EDT

PDB ID : 4R1M
Title : Crystal structure of a Hypothetical Acyl-CoA ligase (BT_0428) from *Bacteroides thetaiotaomicron* VPI-5482 at 2.48 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2014-08-06
Resolution : 2.48 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable23489
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable23489

1 Overall quality at a glance

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3277 (2.50-2.46)
Clashscore	79885	4136 (2.50-2.46)
Ramachandran outliers	78287	4052 (2.50-2.46)
Sidechain outliers	78261	4054 (2.50-2.46)
RSRZ outliers	66119	3279 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	436	
1	B	436	
1	C	436	
1	D	436	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	AMP	A	502	-	X
5	EDO	A	504	-	X
5	EDO	A	505	-	X
5	EDO	A	506	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
5	EDO	A	507	-	X
5	EDO	A	509	-	X
5	EDO	A	510	-	X
5	EDO	A	511	-	X
5	EDO	A	512	-	X
5	EDO	B	504	-	X
5	EDO	C	506	-	X
5	EDO	C	508	-	X
5	EDO	D	503	-	X
5	EDO	D	504	-	X
6	CL	C	505	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 13860 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylacetate-coenzymeA ligase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	435	Total	C	N	O	S	Se	0	2	0
			3397	2144	589	644	6	14			
1	B	426	Total	C	N	O	S	Se	0	1	0
			3310	2088	577	625	6	14			
1	C	431	Total	C	N	O	S	Se	0	3	0
			3392	2142	596	634	6	14			
1	D	428	Total	C	N	O	S	Se	0	3	0
			3341	2108	578	635	6	14			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP Q8AAN6
B	0	GLY	-	LEADER SEQUENCE	UNP Q8AAN6
C	0	GLY	-	LEADER SEQUENCE	UNP Q8AAN6
D	0	GLY	-	LEADER SEQUENCE	UNP Q8AAN6

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		
2	D	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	D	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		
4	A	1	Total	Na	0	0
			1	1		
4	C	2	Total	Na	0	0
			2	2		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	D	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	2	Total	Cl	0	0
			2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	97	Total	O	0	0
			97	97		
7	B	72	Total	O	0	0
			72	72		
7	C	54	Total	O	0	0
			54	54		
7	D	58	Total	O	0	0
			58	58		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phenylacetate-coenzymeA ligase




- Molecule 1: Phenylacetate-coenzymeA ligase



- Molecule 1: Phenylacetate-coenzymeA ligase



- Molecule 1: Phenylacetate-coenzymeA ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	127.06Å 211.87Å 72.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.89 – 2.48 48.89 – 2.48	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.89-2.48) 99.5 (48.89-2.48)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 2.48Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.175 , 0.200 0.181 , 0.208	Depositor DCC
R_{free} test set	3483 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	52.8	Xtriage
Anisotropy	0.492	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 69658 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13860	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ZN, AMP, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/3447	0.67	0/4636
1	B	0.50	0/3357	0.66	0/4519
1	C	0.51	0/3444	0.67	0/4627
1	D	0.50	0/3394	0.65	0/4571
All	All	0.51	0/13642	0.66	0/18353

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3397	0	3350	9	0
1	B	3310	0	3237	11	0
1	C	3392	0	3385	11	0
1	D	3341	0	3261	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	23	0	12	0	0
3	B	23	0	12	0	0
3	D	23	0	12	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	2	0	0	0	0
5	A	36	0	54	2	0
5	B	4	0	6	0	0
5	C	12	0	18	0	0
5	D	8	0	12	0	0
6	C	2	0	0	0	0
7	A	97	0	0	0	0
7	B	72	0	0	0	0
7	C	54	0	0	0	0
7	D	58	0	0	0	0
All	All	13860	0	13359	43	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (43) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:90:ILE:HG23	1:D:102:VAL:HG13	1.69	0.75
1:C:185:PRO:HD2	1:C:215:GLU:HG2	1.70	0.74
1:C:335:ILE:HD11	1:C:370:ASP:HB3	1.80	0.63
1:B:332:MSE:HE3	1:B:341:PHE:CE2	2.36	0.61
1:B:90:ILE:HG23	1:B:102:VAL:HG13	1.83	0.61
1:C:366:VAL:HG13	1:C:367:ASN:H	1.65	0.60
1:A:90:ILE:HG23	1:A:102:VAL:HG13	1.85	0.59
1:D:332:MSE:HE3	1:D:341:PHE:CE2	2.38	0.57
1:D:348:ILE:HD12	1:D:400:LEU:HD23	1.89	0.55
1:C:185:PRO:HD2	1:C:215:GLU:CG	2.35	0.54
1:A:291:ASP:HA	5:A:512:EDO:H21	1.90	0.53
1:C:90:ILE:HG23	1:C:102:VAL:HG13	1.89	0.52
1:C:335:ILE:CD1	1:C:370:ASP:HB3	2.39	0.52
1:A:354:GLU:HG2	1:A:392:ILE:HD13	1.91	0.52
1:D:237:PHE:HB3	1:D:248:ALA:HB3	1.93	0.51
1:C:166:SER:HB3	1:C:194:VAL:HG21	1.93	0.50
1:D:214:ALA:HB1	1:D:424:LYS:HE2	1.94	0.49
1:C:335:ILE:HG22	1:C:340:ILE:HD12	1.95	0.48
1:D:389:LEU:HA	1:D:392:ILE:HD12	1.95	0.47
1:B:92:SER:HB2	1:B:140:MSE:HE3	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:30:ILE:HD13	1:A:53:ILE:HG13	1.99	0.45
1:D:136:SER:O	1:D:162:ALA:HA	2.17	0.45
1:B:237:PHE:HB3	1:B:248:ALA:HB3	1.98	0.44
1:D:237:PHE:CZ	1:D:239:MSE:HE3	2.53	0.43
1:A:136:SER:O	1:A:162:ALA:HA	2.19	0.43
1:D:239:MSE:HE1	1:D:302:THR:HB	2.01	0.43
1:B:332:MSE:HE2	1:B:339:ASN:HB3	2.01	0.42
1:C:136:SER:O	1:C:162:ALA:HA	2.19	0.42
1:C:366:VAL:HG13	1:C:367:ASN:N	2.34	0.42
1:D:90:ILE:HG23	1:D:102:VAL:CG1	2.43	0.42
1:B:348:ILE:HD12	1:B:400:LEU:HD23	2.01	0.42
1:B:30:ILE:HD13	1:B:53:ILE:HG13	2.02	0.42
1:A:102:VAL:H	5:A:506:EDO:H12	1.85	0.42
1:D:348:ILE:HD11	1:D:403:GLU:CB	2.50	0.41
1:D:400:LEU:O	1:D:404:ILE:HG12	2.21	0.41
1:A:90:ILE:HG13	1:B:157:LEU:HD11	2.02	0.41
1:D:30:ILE:HD13	1:D:53:ILE:HG13	2.03	0.41
1:A:185:PRO:HD2	1:A:215:GLU:HG2	2.03	0.41
1:B:214:ALA:HB1	1:B:424:LYS:HE2	2.01	0.41
1:D:332:MSE:HE2	1:D:339:ASN:HB3	2.03	0.40
1:A:129:THR:HB	1:B:84:LYS:HE2	2.02	0.40
1:B:282:ILE:HD13	1:B:326:LYS:HD2	2.02	0.40
1:C:219:ASP:O	1:C:223:ARG:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	435/436 (100%)	419 (96%)	15 (3%)	1 (0%)	56 77
1	B	423/436 (97%)	408 (96%)	12 (3%)	3 (1%)	30 48
1	C	430/436 (99%)	410 (95%)	18 (4%)	2 (0%)	38 59

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	427/436 (98%)	410 (96%)	15 (4%)	2 (0%)	38	59
All	All	1715/1744 (98%)	1647 (96%)	60 (4%)	8 (0%)	38	59

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	243	ASN
1	B	243	ASN
1	C	84	LYS
1	C	243	ASN
1	D	243	ASN
1	B	382	SER
1	B	367	ASN
1	D	366	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/370 (98%)	360 (99%)	2 (1%)	92	98
1	B	350/370 (95%)	347 (99%)	3 (1%)	87	97
1	C	365/370 (99%)	362 (99%)	3 (1%)	89	97
1	D	355/370 (96%)	351 (99%)	4 (1%)	84	95
All	All	1432/1480 (97%)	1420 (99%)	12 (1%)	89	97

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	210	LEU
1	A	259	PHE
1	B	210	LEU
1	B	259	PHE
1	B	330	ASP
1	C	210	LEU
1	C	221	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	259	PHE
1	D	210	LEU
1	D	249	PHE
1	D	259	PHE
1	D	369	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 28 ligands modelled in this entry, 10 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	AMP	A	502	-	25,25,25	0.64	0	38,38,38	0.81	0
5	EDO	A	504	-	3,3,3	0.69	0	2,2,2	0.32	0
5	EDO	A	505	-	3,3,3	0.68	0	2,2,2	0.35	0
5	EDO	A	506	-	3,3,3	0.54	0	2,2,2	0.64	0
5	EDO	A	507	-	3,3,3	0.82	0	2,2,2	0.11	0
5	EDO	A	508	-	3,3,3	0.68	0	2,2,2	0.26	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	A	509	-	3,3,3	0.69	0	2,2,2	0.40	0
5	EDO	A	510	-	3,3,3	0.86	0	2,2,2	0.11	0
5	EDO	A	511	-	3,3,3	0.77	0	2,2,2	0.23	0
5	EDO	A	512	-	3,3,3	0.66	0	2,2,2	0.31	0
3	AMP	B	502	-	25,25,25	0.59	0	38,38,38	0.79	1 (2%)
5	EDO	B	504	-	3,3,3	0.62	0	2,2,2	0.43	0
5	EDO	C	506	-	3,3,3	0.63	0	2,2,2	0.44	0
5	EDO	C	507	-	3,3,3	0.66	0	2,2,2	0.39	0
5	EDO	C	508	-	3,3,3	0.70	0	2,2,2	0.35	0
3	AMP	D	502	-	25,25,25	0.59	0	38,38,38	0.85	1 (2%)
5	EDO	D	503	-	3,3,3	0.80	0	2,2,2	0.33	0
5	EDO	D	504	-	3,3,3	0.74	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AMP	A	502	-	-	0/10/26/26	0/3/3/3
5	EDO	A	504	-	-	0/1/1/1	0/0/0/0
5	EDO	A	505	-	-	0/1/1/1	0/0/0/0
5	EDO	A	506	-	-	0/1/1/1	0/0/0/0
5	EDO	A	507	-	-	0/1/1/1	0/0/0/0
5	EDO	A	508	-	-	0/1/1/1	0/0/0/0
5	EDO	A	509	-	-	0/1/1/1	0/0/0/0
5	EDO	A	510	-	-	0/1/1/1	0/0/0/0
5	EDO	A	511	-	-	0/1/1/1	0/0/0/0
5	EDO	A	512	-	-	0/1/1/1	0/0/0/0
3	AMP	B	502	-	-	0/10/26/26	0/3/3/3
5	EDO	B	504	-	-	0/1/1/1	0/0/0/0
5	EDO	C	506	-	-	0/1/1/1	0/0/0/0
5	EDO	C	507	-	-	0/1/1/1	0/0/0/0
5	EDO	C	508	-	-	0/1/1/1	0/0/0/0
3	AMP	D	502	-	-	0/10/26/26	0/3/3/3
5	EDO	D	503	-	-	0/1/1/1	0/0/0/0
5	EDO	D	504	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	D	502	AMP	P-O5'-C5'	2.58	125.24	118.63
3	B	502	AMP	P-O5'-C5'	2.12	124.06	118.63

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	435/436 (99%)	0.02	19 (4%)	33 34	36, 57, 105, 138	0
1	B	426/436 (97%)	0.34	33 (7%)	13 13	40, 63, 108, 148	0
1	C	431/436 (98%)	0.05	11 (2%)	53 55	41, 67, 103, 139	0
1	D	428/436 (98%)	0.07	19 (4%)	33 34	41, 64, 110, 142	0
All	All	1720/1744 (98%)	0.12	82 (4%)	29 30	36, 63, 107, 148	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	383	THR	6.6
1	B	410	VAL	5.7
1	A	382	SER	5.4
1	B	340	ILE	5.3
1	B	400	LEU	5.2
1	A	381	LEU	5.0
1	A	387	ILE	4.8
1	B	355	LEU	4.8
1	A	383	THR	4.7
1	B	396	ILE	4.7
1	B	412	LEU	4.4
1	A	421	SER	4.2
1	B	386	TYR	4.1
1	B	411	LYS	4.1
1	B	405	LEU	4.0
1	C	1	MSE	4.0
1	A	366	VAL	3.9
1	B	378	LEU	3.9
1	D	380	ASP	3.8
1	B	333	PHE	3.8
1	B	334	ILE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	0	GLY	3.5
1	D	410	VAL	3.4
1	B	389	LEU	3.4
1	B	392	ILE	3.3
1	A	425	ALA	3.3
1	B	382	SER	3.2
1	B	408	PRO	3.2
1	C	386	TYR	3.2
1	C	59	ILE	3.2
1	D	373	ILE	3.2
1	C	0	GLY	3.1
1	D	405	LEU	3.1
1	D	338	VAL	3.0
1	D	389	LEU	3.0
1	A	384	ASP	3.0
1	A	373	ILE	2.9
1	B	387	ILE	2.9
1	C	381	LEU	2.9
1	B	374	VAL	2.9
1	A	388	GLU	2.8
1	A	427	ARG	2.7
1	A	426	VAL	2.7
1	D	430	ASP	2.7
1	D	369	GLN	2.6
1	C	380	ASP	2.6
1	D	396	ILE	2.6
1	B	98	GLY	2.6
1	C	53	ILE	2.6
1	C	387	ILE	2.5
1	A	412	LEU	2.5
1	B	413	VAL	2.5
1	A	386	TYR	2.4
1	A	355	LEU	2.4
1	D	382	SER	2.4
1	D	334	ILE	2.4
1	D	392	ILE	2.4
1	B	351[A]	GLN	2.4
1	A	392	ILE	2.4
1	B	93	SER	2.4
1	B	341	PHE	2.4
1	C	421	SER	2.3
1	D	374	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	354	GLU	2.3
1	B	384	ASP	2.3
1	D	378	LEU	2.3
1	A	2	SER	2.3
1	D	404	ILE	2.2
1	B	407	THR	2.2
1	D	340	ILE	2.2
1	A	376	VAL	2.2
1	B	363	LEU	2.2
1	C	422	GLU	2.1
1	B	404	ILE	2.1
1	B	418	LEU	2.1
1	D	406	VAL	2.1
1	B	431	LEU	2.1
1	B	345	VAL	2.1
1	D	339	ASN	2.1
1	C	3	THR	2.0
1	D	2	SER	2.0
1	B	9	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	EDO	A	512	4/4	0.64	29.05	84,84,85,85	0
5	EDO	A	509	4/4	0.43	12.68	77,77,78,78	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	EDO	C	508	4/4	0.41	9.14	73,76,79,81	0
5	EDO	A	505	4/4	0.34	8.66	75,76,77,78	0
5	EDO	A	504	4/4	0.33	8.58	61,62,62,62	0
5	EDO	A	507	4/4	0.34	8.42	50,52,54,54	0
5	EDO	A	506	4/4	0.29	6.68	81,82,83,85	0
6	CL	C	505	1/1	0.20	5.41	90,90,90,90	0
5	EDO	D	504	4/4	0.22	5.03	63,65,68,71	0
5	EDO	B	504	4/4	0.24	3.96	85,85,85,85	0
5	EDO	D	503	4/4	0.20	3.71	61,62,62,63	0
5	EDO	A	510	4/4	0.18	3.58	64,66,69,71	0
3	AMP	A	502	23/23	0.20	2.96	78,87,103,106	0
5	EDO	A	511	4/4	0.25	2.88	68,72,75,76	0
5	EDO	C	506	4/4	0.15	2.08	87,87,88,88	0
5	EDO	A	508	4/4	0.19	0.23	78,81,83,83	0
5	EDO	C	507	4/4	0.14	0.04	83,84,86,87	0
2	ZN	A	501	1/1	0.11	-0.27	52,52,52,52	0
2	ZN	C	501	1/1	0.09	-0.45	66,66,66,66	0
3	AMP	B	502	23/23	0.14	-0.56	50,52,62,65	0
2	ZN	B	501	1/1	0.13	-0.87	55,55,55,55	0
4	NA	A	503	1/1	0.12	-1.06	51,51,51,51	0
6	CL	C	504	1/1	0.07	-1.20	59,59,59,59	0
2	ZN	D	501	1/1	0.10	-1.29	55,55,55,55	0
4	NA	C	502	1/1	0.07	-1.43	53,53,53,53	0
3	AMP	D	502	23/23	0.11	-1.53	62,71,86,88	0
4	NA	C	503	1/1	0.07	-1.61	52,52,52,52	0
4	NA	B	503	1/1	0.07	-2.06	51,51,51,51	0

6.5 Other polymers ⓘ

There are no such residues in this entry.