



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 12:38 AM GMT

PDB ID : 3R2G
Title : Crystal structure of Inosine 5' monophosphate dehydrogenase from Legionella pneumophila
Authors : Agarwal, R.; Almo, S.C.; Swaminathan, S.; New York Structural Genomics Research Consortium (NYSGRC)
Deposited on : 2011-03-14
Resolution : 1.94 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2024 (1.96-1.92)
Clashscore	79885	2281 (1.96-1.92)
Ramachandran outliers	78287	2255 (1.96-1.92)
Sidechain outliers	78261	2255 (1.96-1.92)
RSRZ outliers	66119	2024 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	361	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2622 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inosine 5'-monophosphate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	323	2405	1500	425	461	19	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP Q5ZRN7
A	2	VAL	-	EXPRESSION TAG	UNP Q5ZRN7
A	340	ALA	-	EXPRESSION TAG	UNP Q5ZRN7
A	341	GLU	-	EXPRESSION TAG	UNP Q5ZRN7
A	342	ASN	-	EXPRESSION TAG	UNP Q5ZRN7
A	343	LEU	-	EXPRESSION TAG	UNP Q5ZRN7
A	344	TYR	-	EXPRESSION TAG	UNP Q5ZRN7
A	345	PHE	-	EXPRESSION TAG	UNP Q5ZRN7
A	346	GLN	-	EXPRESSION TAG	UNP Q5ZRN7
A	347	SER	-	EXPRESSION TAG	UNP Q5ZRN7
A	348	HIS	-	EXPRESSION TAG	UNP Q5ZRN7
A	349	HIS	-	EXPRESSION TAG	UNP Q5ZRN7
A	350	HIS	-	EXPRESSION TAG	UNP Q5ZRN7
A	351	HIS	-	EXPRESSION TAG	UNP Q5ZRN7
A	352	HIS	-	EXPRESSION TAG	UNP Q5ZRN7
A	353	HIS	-	EXPRESSION TAG	UNP Q5ZRN7
A	354	TRP	-	EXPRESSION TAG	UNP Q5ZRN7
A	355	SER	-	EXPRESSION TAG	UNP Q5ZRN7
A	356	HIS	-	EXPRESSION TAG	UNP Q5ZRN7
A	357	PRO	-	EXPRESSION TAG	UNP Q5ZRN7
A	358	GLN	-	EXPRESSION TAG	UNP Q5ZRN7
A	359	PHE	-	EXPRESSION TAG	UNP Q5ZRN7
A	360	GLU	-	EXPRESSION TAG	UNP Q5ZRN7
A	361	LYS	-	EXPRESSION TAG	UNP Q5ZRN7

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	217	Total 217	O 217	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Inosine 5'-monophosphate dehydrogenase

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	115.13Å 115.13Å 63.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.70 – 1.94 40.70 – 1.94	Depositor EDS
% Data completeness (in resolution range)	97.6 (40.70-1.94) 97.6 (40.70-1.94)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.179 , 0.207 0.181 , 0.175	Depositor DCC
R_{free} test set	1516 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	27.2	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 41.5	EDS
Estimated twinning fraction	0.027 for -k,-h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 29904 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2622	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/2441	0.70	0/3289

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2405	0	2399	28	0
2	A	217	0	0	3	0
All	All	2622	0	2399	28	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (28) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:80:ASN:HD22	1:A:104:ARG:HH11	1.41	0.68
1:A:45:VAL:CG1	1:A:227:ILE:HD12	2.25	0.67
1:A:35:ARG:HG3	1:A:35:ARG:NH1	2.09	0.66
1:A:35:ARG:CG	1:A:35:ARG:HH11	2.14	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:35:ARG:HG3	1:A:35:ARG:HH11	1.66	0.60
1:A:150:ALA:HB2	1:A:170:ILE:HG12	1.84	0.59
1:A:45:VAL:HG13	1:A:227:ILE:HD12	1.87	0.56
1:A:35:ARG:CG	1:A:35:ARG:NH1	2.71	0.53
1:A:175:VAL:HG22	1:A:178:THR:H	1.73	0.53
1:A:245:ASP:O	1:A:246:ASP:CB	2.57	0.52
1:A:31:THR:HG22	2:A:448:HOH:O	2.10	0.51
1:A:171:GLY:HA3	1:A:175:VAL:HG13	1.95	0.49
1:A:35:ARG:NH2	2:A:415:HOH:O	2.46	0.48
1:A:172:GLY:HA2	1:A:179:ARG:CD	2.44	0.47
1:A:173:GLY:O	1:A:174:SER:HB2	2.17	0.45
1:A:45:VAL:CG1	1:A:227:ILE:CD1	2.94	0.45
1:A:143:CYS:C	1:A:144:ILE:HD13	2.37	0.45
1:A:227:ILE:H	1:A:227:ILE:HD13	1.81	0.44
1:A:188:MET:O	1:A:192:ILE:HG12	2.17	0.44
1:A:32:SER:HB3	1:A:44:PRO:HB3	1.99	0.44
1:A:75:MET:CE	1:A:83:GLU:HG2	2.47	0.44
1:A:51:ASP:HA	1:A:72:HIS:CD2	2.52	0.44
1:A:81:ILE:CD1	1:A:108:LEU:HD23	2.49	0.42
1:A:126:LYS:HE3	1:A:130:LYS:HE3	2.02	0.42
1:A:150:ALA:HB2	1:A:170:ILE:CG1	2.51	0.40
1:A:144:ILE:N	1:A:144:ILE:HD13	2.36	0.40
1:A:181:LYS:HE3	2:A:442:HOH:O	2.21	0.40
1:A:8:ILE:HG21	1:A:322:ILE:HD12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	321/361 (89%)	303 (94%)	14 (4%)	4 (1%)	19 6

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	174	SER
1	A	246	ASP
1	A	171	GLY
1	A	172	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	251/289 (87%)	243 (97%)	8 (3%)	51 36

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	ASP
1	A	35	ARG
1	A	144	ILE
1	A	177	SER
1	A	227	ILE
1	A	245	ASP
1	A	246	ASP
1	A	273	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	ASN
1	A	262	GLN
1	A	268	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/361 (89%)	0.20	22 (6%) 17 20	16, 25, 50, 64	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	173	GLY	11.4
1	A	174	SER	7.5
1	A	175	VAL	5.4
1	A	25	ARG	4.7
1	A	171	GLY	4.4
1	A	27	VAL	4.4
1	A	247	GLY	4.2
1	A	274	THR	4.2
1	A	248	SER	4.1
1	A	246	ASP	3.8
1	A	172	GLY	3.7
1	A	275	ALA	3.1
1	A	273	LYS	3.1
1	A	245	ASP	2.9
1	A	243	GLN	2.9
1	A	278	VAL	2.7
1	A	286	GLU	2.7
1	A	88	LYS	2.6
1	A	271	GLU	2.6
1	A	170	ILE	2.4
1	A	176	CYS	2.2
1	A	241	VAL	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.