



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 08:25 AM GMT

PDB ID : 2R4U
Title : Crystal Structure of Wild-type E.coli GS in complex with ADP and Glucose(wtGSd)
Authors : Sheng, F.; Geiger, J.
Deposited on : 2007-09-01
Resolution : 2.37 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3327 (2.40-2.32)
Clashscore	79885	1064 (2.38-2.34)
Ramachandran outliers	78287	1048 (2.38-2.34)
Sidechain outliers	78261	1049 (2.38-2.34)
RSRZ outliers	66119	3330 (2.40-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	485	

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 4014 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycogen synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	477	3735	2401	652	669	13	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	478	LEU	-	EXPRESSION TAG	UNP P0A6U8
A	479	GLU	-	EXPRESSION TAG	UNP P0A6U8
A	480	HIS	-	EXPRESSION TAG	UNP P0A6U8
A	481	HIS	-	EXPRESSION TAG	UNP P0A6U8
A	482	HIS	-	EXPRESSION TAG	UNP P0A6U8
A	483	HIS	-	EXPRESSION TAG	UNP P0A6U8
A	484	HIS	-	EXPRESSION TAG	UNP P0A6U8
A	485	HIS	-	EXPRESSION TAG	UNP P0A6U8

- Molecule 2 is SUGAR (GLUCOSE) (three-letter code: GLC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is (2R)-2-HYDROXY-3-[4-(2-HYDROXYETHYL)PIPERAZIN-1-YL]PROPAN E-1-SULFONICACID (three-letter code: 250) (formula: $C_9H_{20}N_2O_5S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			17	9	2	5	1		

- Molecule 5 is 3,6,9,12,15,18,21,24,27,30,33,36,39-TRIDECAXAHENTETRACONTANE-1,41-DIOL (three-letter code: PE3) (formula: $C_{28}H_{58}O_{15}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			13	9	4		
5	A	1	Total	C	O	0	0
			5	3	2		
5	A	1	Total	C	O	0	0
			6	4	2		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	199	Total	O	0	0
			199	199		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycogen synthase

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, α , β , γ	126.17Å 126.17Å 151.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	97.13 – 2.37 40.53 – 2.37	Depositor EDS
% Data completeness (in resolution range)	97.7 (97.13-2.37) 97.7 (40.53-2.37)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.37Å)	Xtriage
Refinement program	REFMAC 5.2	Depositor
R, R_{free}	0.182 , 0.203 0.181 , 0.200	Depositor DCC
R_{free} test set	2386 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	44.5	Xtriage
Anisotropy	0.308	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.1	EDS
Estimated twinning fraction	0.021 for -h,k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 47143 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4014	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 250, GLC, ADP, PE3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/3834	0.73	6/5213 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	396	ARG	NE-CZ-NH2	-7.69	116.45	120.30
1	A	470	ARG	NE-CZ-NH2	-5.96	117.32	120.30
1	A	48	ARG	NE-CZ-NH2	-5.91	117.35	120.30
1	A	470	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	A	48	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	A	267	ARG	NE-CZ-NH2	-5.12	117.74	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3735	0	3679	67	0
2	A	12	0	12	0	0
3	A	27	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	17	0	20	0	0
5	A	24	0	25	0	0
6	A	199	0	0	11	0
All	All	4014	0	3748	67	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (67) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:159:THR:HB	1:A:209:THR:HG22	1.50	0.90
1:A:1:MET:HE1	1:A:132:VAL:HG23	1.52	0.89
1:A:364:MET:O	1:A:391:THR:HG21	1.77	0.83
1:A:214:THR:HG23	1:A:403:ASP:OD1	1.79	0.82
1:A:62:THR:HG22	1:A:64:ALA:H	1.43	0.82
1:A:211:VAL:HG13	1:A:379:CYS:HB2	1.63	0.80
1:A:383:GLN:H	1:A:383:GLN:HE21	1.31	0.76
1:A:150:ARG:HG3	6:A:528:HOH:O	1.84	0.76
1:A:134:HIS:NE2	1:A:159:THR:CG2	2.48	0.75
1:A:209:THR:CG2	6:A:507:HOH:O	2.34	0.74
1:A:62:THR:CG2	1:A:64:ALA:H	1.99	0.74
1:A:163:LEU:HD23	1:A:225:ALA:HA	1.72	0.72
1:A:139:HIS:ND1	6:A:625:HOH:O	2.22	0.72
1:A:273:LYS:NZ	1:A:391:THR:HG22	2.06	0.71
1:A:258:ASP:HB3	1:A:261:LEU:HD22	1.74	0.70
1:A:62:THR:HG21	1:A:112:LEU:HD13	1.73	0.69
1:A:90:ARG:HD3	6:A:523:HOH:O	1.94	0.68
1:A:69:LEU:HD22	1:A:124:LEU:HD13	1.77	0.66
1:A:211:VAL:CG1	1:A:379:CYS:HB2	2.26	0.65
1:A:1:MET:CE	1:A:132:VAL:HG23	2.27	0.62
1:A:273:LYS:HZ2	1:A:391:THR:HG22	1.63	0.62
1:A:380:GLY:C	1:A:383:GLN:HE22	2.03	0.62
1:A:471:GLU:HG3	6:A:543:HOH:O	1.99	0.62
1:A:209:THR:HG23	6:A:507:HOH:O	1.95	0.61
1:A:62:THR:HG22	1:A:64:ALA:N	2.14	0.61
1:A:115:TRP:CH2	1:A:119:GLU:HG3	2.36	0.60
1:A:60:ARG:NH1	6:A:676:HOH:O	2.37	0.58
1:A:252:ILE:HG13	1:A:253:TRP:CD1	2.40	0.57
1:A:199:LYS:HG2	1:A:228:MET:HE1	1.88	0.56
1:A:214:THR:HG23	1:A:403:ASP:CG	2.26	0.55
1:A:98:THR:HG23	6:A:684:HOH:O	2.07	0.55
1:A:62:THR:HB	1:A:65:GLY:O	2.08	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:26:LEU:HD22	1:A:30:GLN:HG3	1.88	0.54
1:A:134:HIS:NE2	1:A:159:THR:HG21	2.23	0.54
1:A:3:VAL:HG21	1:A:473:TYR:CE2	2.44	0.53
1:A:383:GLN:NE2	1:A:383:GLN:H	2.04	0.53
1:A:374:SER:O	1:A:396:ARG:HD3	2.09	0.53
1:A:214:THR:CG2	1:A:403:ASP:OD1	2.53	0.52
1:A:199:LYS:HG2	1:A:228:MET:CE	2.39	0.52
1:A:134:HIS:NE2	1:A:159:THR:HG23	2.25	0.51
1:A:184:PHE:CZ	1:A:190:GLU:HG3	2.46	0.51
1:A:207:HIS:HE1	1:A:242:SER:OG	1.94	0.50
1:A:219:ILE:O	1:A:225:ALA:HB2	2.12	0.49
1:A:267:ARG:HD2	1:A:453:ARG:O	2.12	0.49
1:A:90:ARG:NH2	1:A:105:ASP:OD2	2.42	0.48
1:A:211:VAL:HG22	1:A:246:ASN:HD21	1.79	0.47
1:A:26:LEU:HB3	1:A:27:PRO:HD3	1.97	0.47
1:A:383:GLN:N	1:A:383:GLN:HE21	2.04	0.46
1:A:163:LEU:CD2	1:A:225:ALA:HA	2.46	0.44
1:A:33:ASP:OD2	1:A:470:ARG:HD3	2.17	0.44
1:A:273:LYS:HZ1	1:A:391:THR:HG22	1.82	0.44
1:A:252:ILE:HD12	1:A:361:HIS:CE1	2.53	0.43
1:A:159:THR:HB	1:A:209:THR:CG2	2.36	0.42
1:A:458:MET:HE2	6:A:559:HOH:O	2.19	0.42
1:A:181:TRP:NE1	6:A:666:HOH:O	2.29	0.42
1:A:320:GLN:HE22	1:A:436:ARG:HG2	1.85	0.42
1:A:160:VAL:HG21	1:A:163:LEU:HD13	2.02	0.41
1:A:147:LEU:HD23	1:A:156:SER:HB3	2.03	0.41
1:A:38:ARG:HD2	6:A:508:HOH:O	2.20	0.41
1:A:215:TYR:CG	1:A:378:PRO:HA	2.56	0.41
1:A:1:MET:HE2	1:A:473:TYR:HB3	2.03	0.41
1:A:3:VAL:HG21	1:A:473:TYR:CD2	2.56	0.41
1:A:228:MET:HE2	1:A:228:MET:HB3	1.64	0.41
1:A:90:ARG:HH22	1:A:105:ASP:CG	2.25	0.40
1:A:196:SER:HB3	1:A:199:LYS:HB2	2.02	0.40
1:A:396:ARG:HG3	1:A:423:PHE:CE1	2.56	0.40
1:A:394:LEU:C	1:A:394:LEU:HD23	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	475/485 (98%)	462 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	382/390 (98%)	347 (91%)	35 (9%)	13	13

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LEU
1	A	26	LEU
1	A	55	GLN
1	A	62	THR
1	A	77	VAL
1	A	90	ARG
1	A	98	THR
1	A	100	LEU
1	A	108	LEU
1	A	113	LEU
1	A	124	LEU
1	A	147	LEU
1	A	157	VAL
1	A	159	THR
1	A	160	VAL
1	A	178	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	198	LEU
1	A	209	THR
1	A	211	VAL
1	A	214	THR
1	A	228	MET
1	A	231	LEU
1	A	252	ILE
1	A	259	LEU
1	A	261	LEU
1	A	292	VAL
1	A	301	LEU
1	A	383	GLN
1	A	388	LYS
1	A	391	THR
1	A	396	ARG
1	A	411	GLU
1	A	417	VAL
1	A	436	ARG
1	A	472	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	HIS
1	A	192	ASN
1	A	207	HIS
1	A	223	GLN
1	A	233	GLN
1	A	246	ASN
1	A	279	GLN
1	A	281	GLN
1	A	320	GLN
1	A	383	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ADP	A	501	-	29,29,29	0.96	1 (3%)	45,45,45	2.00	9 (20%)
2	GLC	A	502	-	12,12,12	0.66	0	17,17,17	1.04	1 (5%)
4	250	A	503	-	17,17,17	2.28	1 (5%)	23,23,23	2.34	10 (43%)
5	PE3	A	504	-	11,12,42	1.54	1 (9%)	9,11,41	0.22	0
5	PE3	A	505	-	3,4,42	1.01	0	2,3,41	1.09	0
5	PE3	A	506	-	5,5,42	2.29	1 (20%)	3,4,41	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	501	-	-	0/16/32/32	0/1/3/3
2	GLC	A	502	-	-	0/2/22/22	0/1/1/1
4	250	A	503	-	-	0/12/22/22	0/1/1/1
5	PE3	A	504	-	-	0/10/10/40	0/0/0/0
5	PE3	A	505	-	-	0/2/2/40	0/0/0/0
5	PE3	A	506	-	-	0/3/3/40	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	250	C1-S1	-9.03	1.67	1.77
5	A	506	PE3	C6-C5	-5.05	1.53	1.55
5	A	504	PE3	C21-C20	-4.75	1.53	1.55
3	A	501	ADP	C4-N9	-3.14	1.33	1.37

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	ADP	N3-C2-N1	-9.75	120.56	128.71
4	A	503	250	C6-N2-C5	5.39	122.24	108.86
4	A	503	250	C7-N1-C4	4.37	119.70	108.86
3	A	501	ADP	C8-N9-C4	4.12	110.04	106.90
3	A	501	ADP	N3-C4-N9	3.79	132.27	125.43
4	A	503	250	C8-N2-C6	3.06	119.21	111.32
4	A	503	250	O3-S1-C1	2.98	109.47	105.64
3	A	501	ADP	O2B-PB-O1B	2.87	119.81	110.44
4	A	503	250	O1-S1-C1	2.87	112.05	107.03
4	A	503	250	C8-N2-C5	2.79	118.53	111.32
4	A	503	250	C4-N1-C3	2.76	120.76	111.13
3	A	501	ADP	C1'-N9-C4	-2.67	122.02	126.64
2	A	502	GLC	O5-C5-C4	-2.44	105.25	109.76
4	A	503	250	C7-N1-C3	2.40	119.50	111.13
3	A	501	ADP	N7-C8-N9	-2.31	107.81	114.36
3	A	501	ADP	C5-C4-N3	-2.13	121.07	125.70
3	A	501	ADP	C4-C5-N7	-2.08	107.74	109.52
4	A	503	250	C9-C8-N2	-2.07	106.83	113.62
3	A	501	ADP	C2-N3-C4	2.04	119.81	114.01
4	A	503	250	C4-C5-N2	-2.00	106.67	110.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	477/485 (98%)	-0.07	12 (2%) 54 58	30, 40, 56, 65	29 (6%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	343	ALA	3.9
1	A	101	PHE	3.7
1	A	170	TYR	3.0
1	A	181	TRP	2.9
1	A	98	THR	2.7
1	A	237	ARG	2.6
1	A	99	ASN	2.5
1	A	97	ASP	2.3
1	A	346	PRO	2.1
1	A	192	ASN	2.1
1	A	186	ILE	2.1
1	A	127	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	PE3	A	505	5/43	0.18	1.88	76,78,80,80	0
4	250	A	503	17/17	0.18	1.79	68,76,81,81	0
2	GLC	A	502	12/12	0.22	1.60	28,34,37,40	12
5	PE3	A	504	13/43	0.14	1.24	79,83,87,87	0
5	PE3	A	506	6/43	0.17	0.98	77,80,81,81	0
3	ADP	A	501	27/27	0.11	-0.50	34,40,42,42	0

6.5 Other polymers ⓘ

There are no such residues in this entry.