



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 09:36 AM GMT

PDB ID : 3R6X  
Title : CDK2 in complex with inhibitor KVR-1-158  
Authors : Betzi, S.; Alam, R.; Han, H.; Becker, A.; Schonbrunn, E.  
Deposited on : 2011-03-22  
Resolution : 1.75 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1134 (1.76-1.76)
Clashscore	79885	1304 (1.76-1.76)
Ramachandran outliers	78287	1288 (1.76-1.76)
Sidechain outliers	78261	1288 (1.76-1.76)
RSRZ outliers	66119	1135 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	306	

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2625 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cyclin-dependent kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	294	2379	1549	407	414	9	0	3	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	EXPRESSION TAG	UNP P24941
A	-6	PRO	-	EXPRESSION TAG	UNP P24941
A	-5	LEU	-	EXPRESSION TAG	UNP P24941
A	-4	GLY	-	EXPRESSION TAG	UNP P24941
A	-3	SER	-	EXPRESSION TAG	UNP P24941
A	-2	PRO	-	EXPRESSION TAG	UNP P24941
A	-1	GLU	-	EXPRESSION TAG	UNP P24941
A	0	PHE	-	EXPRESSION TAG	UNP P24941

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is 4-[(3-HYDROXYPROPYL)AMINO]-5-NITRO-2-[(PYRIDIN-3-YLMETHYL)AMINO]BENZAMIDE (three-letter code: X84) (formula: C<sub>16</sub>H<sub>19</sub>N<sub>5</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			25	16	5	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	217	Total	O	0	0
			217	217		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.61Å 71.59Å 72.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	18.79 – 1.75 18.79 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.3 (18.79-1.75) 98.4 (18.79-1.75)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.28 (at 1.75Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.198 , 0.235 0.200 , 0.235	Depositor DCC
$R_{free}$ test set	1100 reflections (3.87%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.0	Xtriage
Anisotropy	0.336	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 42.4	EDS
Estimated twinning fraction	0.014 for -h,l,k	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 149370 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2625	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, X84

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.40	0/2452	0.63	1/3324 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	20	LYS	N-CA-C	-5.10	97.23	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2379	0	2438	64	0
2	A	4	0	6	0	0
3	A	25	0	19	1	0
4	A	217	0	0	4	0
All	All	2625	0	2463	65	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

All (65) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:115[B]:LEU:HD23	1:A:146:PHE:HZ	1.38	0.88
1:A:15:TYR:HB2	1:A:33:LYS:HE3	1.57	0.84
1:A:125:HIS:HD2	1:A:127:ASP:H	1.22	0.81
1:A:129:LYS:HG3	1:A:131:GLN:HG2	1.63	0.80
1:A:60:HIS:HD2	1:A:62:ASN:H	1.26	0.80
1:A:14:THR:HA	1:A:158:THR:HG22	1.61	0.80
1:A:60:HIS:CD2	1:A:62:ASN:H	2.04	0.74
1:A:115[B]:LEU:CD2	1:A:146:PHE:HZ	2.01	0.72
1:A:115[B]:LEU:HD23	1:A:146:PHE:CZ	2.23	0.70
1:A:7:VAL:HG12	1:A:8:GLU:HG3	1.74	0.69
1:A:125:HIS:HE1	1:A:144:ALA:O	1.77	0.68
1:A:157:ARG:HD2	1:A:162:GLU:O	1.97	0.64
1:A:125:HIS:CD2	1:A:127:ASP:H	2.11	0.64
1:A:1:MET:HB2	1:A:4:PHE:O	1.96	0.64
1:A:230:VAL:O	1:A:233:MET:HG3	1.99	0.62
1:A:155:PRO:HG2	1:A:156:VAL:H	1.68	0.59
1:A:177[B]:CYS:SG	1:A:233:MET:CE	2.93	0.57
1:A:246:GLN:NE2	4:A:527:HOH:O	2.38	0.56
1:A:28:GLU:HG3	1:A:30:VAL:HG13	1.88	0.55
1:A:124:LEU:HD22	1:A:182:THR:HA	1.90	0.54
1:A:200:ARG:HD2	4:A:749:HOH:O	2.07	0.53
1:A:200:ARG:HG2	4:A:646:HOH:O	2.08	0.53
1:A:17:VAL:CG1	1:A:34:LYS:HD3	2.39	0.52
1:A:12:GLU:HB3	1:A:159:TYR:HE1	1.74	0.52
1:A:97:THR:HG23	4:A:483:HOH:O	2.09	0.52
1:A:12:GLU:HB3	1:A:159:TYR:CE1	2.45	0.51
1:A:34:LYS:HG2	1:A:77:TYR:HE1	1.75	0.51
1:A:34:LYS:HG2	1:A:77:TYR:CE1	2.45	0.51
1:A:56:LYS:NZ	1:A:56:LYS:HB3	2.26	0.51
1:A:18:VAL:HG22	1:A:33:LYS:HG3	1.93	0.51
1:A:48:ALA:O	1:A:52:ILE:HG12	2.10	0.51
1:A:1:MET:HG3	1:A:4:PHE:HB2	1.92	0.50
1:A:169[A]:ARG:HG3	1:A:169[A]:ARG:HH11	1.77	0.50
1:A:15:TYR:HB2	1:A:33:LYS:CE	2.33	0.50
1:A:154:VAL:HG22	1:A:156:VAL:HG23	1.95	0.48
1:A:5:GLN:HG3	1:A:5:GLN:O	2.14	0.48
1:A:9:LYS:HA	1:A:19:TYR:CD1	2.48	0.48
1:A:13:GLY:HA3	1:A:33:LYS:NZ	2.28	0.47
1:A:34:LYS:NZ	1:A:34:LYS:HB3	2.30	0.47
1:A:169[B]:ARG:HD2	1:A:174:LEU:CD2	2.45	0.47
1:A:12:GLU:HG2	1:A:159:TYR:OH	2.14	0.46
1:A:5:GLN:NE2	1:A:22:ARG:HD2	2.31	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:143:LEU:HD22	1:A:146:PHE:HE1	1.81	0.46
1:A:177[B]:CYS:SG	1:A:233:MET:HE1	2.55	0.46
1:A:155:PRO:O	1:A:156:VAL:HB	2.17	0.45
1:A:230:VAL:O	1:A:233:MET:CG	2.64	0.45
1:A:84:HIS:HB3	1:A:298:LEU:HD22	1.99	0.45
3:A:778:X84:N04	3:A:778:X84:O24	2.50	0.45
1:A:15:TYR:HD2	1:A:33:LYS:HE2	1.82	0.44
1:A:125:HIS:O	1:A:126:ARG:HB2	2.17	0.44
1:A:14:THR:O	1:A:158:THR:HA	2.17	0.44
1:A:7:VAL:O	1:A:8:GLU:HG2	2.18	0.44
1:A:245:ARG:NH2	1:A:248:PHE:HE1	2.16	0.44
1:A:177[B]:CYS:SG	1:A:233:MET:HE2	2.59	0.43
1:A:144:ALA:O	1:A:145:ASP:HB2	2.19	0.43
1:A:60:HIS:CG	1:A:61:PRO:HD2	2.54	0.42
1:A:125:HIS:CE1	1:A:144:ALA:O	2.65	0.42
1:A:157:ARG:HH11	1:A:157:ARG:HG3	1.85	0.42
1:A:1:MET:CB	1:A:4:PHE:HB2	2.50	0.42
1:A:160:THR:OG1	1:A:162:GLU:HG3	2.20	0.41
1:A:85:GLN:CD	1:A:296:LEU:HD11	2.41	0.41
1:A:36:ARG:O	1:A:41:THR:HA	2.21	0.40
1:A:157:ARG:HG3	1:A:157:ARG:NH1	2.36	0.40
1:A:46:SER:O	1:A:50:ARG:HG3	2.20	0.40
1:A:129:LYS:CG	1:A:131:GLN:HG2	2.43	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	293/306 (96%)	279 (95%)	11 (4%)	3 (1%)	22 6

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	159	TYR
1	A	156	VAL
1	A	155	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	262/269 (97%)	257 (98%)	5 (2%)	69 47

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	GLU
1	A	68	ASP
1	A	146	PHE
1	A	233	MET
1	A	254	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	60	HIS
1	A	85	GLN
1	A	125	HIS
1	A	268	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	A	777	-	3,3,3	0.58	0	2,2,2	0.32	0
3	X84	A	778	-	26,26,26	3.61	12 (46%)	34,34,34	1.50	6 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	777	-	-	0/1/1/1	0/0/0/0
3	X84	A	778	-	-	0/18/18/18	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	778	X84	C07-C12	-10.53	1.38	1.50
3	A	778	X84	C11-N05	-10.22	1.30	1.46
3	A	778	X84	O25-N05	5.56	1.32	1.23
3	A	778	X84	C10-C11	4.24	1.45	1.40
3	A	778	X84	C10-N04	-3.11	1.29	1.38
3	A	778	X84	O24-N05	3.06	1.31	1.25
3	A	778	X84	C09-C10	2.63	1.43	1.39
3	A	778	X84	C09-C08	2.56	1.43	1.39
3	A	778	X84	C08-N01	-2.42	1.31	1.38
3	A	778	X84	C06-C07	2.29	1.43	1.39
3	A	778	X84	C07-C08	2.17	1.44	1.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	778	X84	C12-N02	2.07	1.37	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	778	X84	C07-C06-C11	4.17	121.74	118.59
3	A	778	X84	C06-C11-N05	3.38	119.22	115.82
3	A	778	X84	C06-C11-C10	-3.30	118.69	121.69
3	A	778	X84	O22-C12-N02	-2.47	119.02	122.59
3	A	778	X84	C09-C10-C11	2.33	119.66	116.68
3	A	778	X84	C19-C20-C21	-2.05	109.52	113.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	294/306 (96%)	1.37	60 (20%) <b>1</b> <b>2</b>	8, 20, 70, 93	1 (0%)

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	44	VAL	20.8
1	A	156	VAL	17.5
1	A	159	TYR	14.9
1	A	155	PRO	14.8
1	A	157	ARG	13.3
1	A	154	VAL	12.8
1	A	160	THR	11.8
1	A	158	THR	10.7
1	A	42	GLU	9.3
1	A	43	GLY	8.6
1	A	2	GLU	8.3
1	A	41	THR	7.5
1	A	25	LEU	7.5
1	A	13	GLY	6.9
1	A	45	PRO	6.9
1	A	12	GLU	6.6
1	A	161	HIS	6.2
1	A	1	MET	6.0
1	A	14	THR	5.7
1	A	36	ARG	5.4
1	A	73	GLU	5.1
1	A	74	ASN	4.8
1	A	15	TYR	4.6
1	A	163	VAL	4.6
1	A	162	GLU	4.6
1	A	153	GLY	4.6
1	A	152	PHE	4.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	5	GLN	4.2
1	A	96	LEU	4.1
1	A	164	VAL	4.1
1	A	179	TYR	4.0
1	A	10	ILE	3.9
1	A	122	ARG	3.8
1	A	72	THR	3.8
1	A	146	PHE	3.7
1	A	3	ASN	3.6
1	A	4	PHE	3.6
1	A	75	LYS	3.6
1	A	297	ARG	3.5
1	A	19	TYR	3.4
1	A	7	VAL	3.4
1	A	47	THR	3.3
1	A	180	TYR	3.2
1	A	46	SER	2.9
1	A	147	GLY	2.8
1	A	148	LEU	2.7
1	A	35	ILE	2.6
1	A	24	LYS	2.6
1	A	34	LYS	2.6
1	A	6	LYS	2.5
1	A	247	ASP	2.5
1	A	246	GLN	2.4
1	A	151	ALA	2.2
1	A	57	GLU	2.2
1	A	126	ARG	2.2
1	A	296	LEU	2.1
1	A	26	THR	2.1
1	A	71	HIS	2.1
1	A	76	LEU	2.0
1	A	150	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EDO	A	777	4/4	0.13	0.92	22,26,27,27	0
3	X84	A	778	25/25	0.17	0.22	18,27,39,43	0

## 6.5 Other polymers

There are no such residues in this entry.