



# Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 07:50 AM GMT

PDB ID : 2RCM  
Title : Crystal Structure of Arabidopsis thaliana Allene Oxide Synthase variant (F137L) (At-AOS(F137L), cytochrome P450 74A) at 1.73 Å Resolution  
Authors : Lee, D.S.; Nioche, P.; Raman, C.S.  
Deposited on : 2007-09-20  
Resolution : 1.73 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1657 (1.76-1.72)
Clashscore	79885	1881 (1.76-1.72)
Ramachandran outliers	78287	1859 (1.76-1.72)
Sidechain outliers	78261	1859 (1.76-1.72)
RSRZ outliers	66119	1658 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	495	
1	B	495	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	GOL	B	1	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8384 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 74A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	465	Total	C	N	O	S	0	9	0
			3778	2435	638	695	10			
1	B	465	Total	C	N	O	S	0	5	0
			3746	2421	628	687	10			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	MET	-	EXPRESSION TAG	UNP Q96242
A	29	ALA	-	EXPRESSION TAG	UNP Q96242
A	30	LYS	-	EXPRESSION TAG	UNP Q96242
A	31	LYS	-	EXPRESSION TAG	UNP Q96242
A	32	THR	-	EXPRESSION TAG	UNP Q96242
A	33	SER	-	EXPRESSION TAG	UNP Q96242
A	137	LEU	PHE	ENGINEERED	UNP Q96242
A	519	HIS	-	EXPRESSION TAG	UNP Q96242
A	520	HIS	-	EXPRESSION TAG	UNP Q96242
A	521	HIS	-	EXPRESSION TAG	UNP Q96242
A	522	HIS	-	EXPRESSION TAG	UNP Q96242
B	28	MET	-	EXPRESSION TAG	UNP Q96242
B	29	ALA	-	EXPRESSION TAG	UNP Q96242
B	30	LYS	-	EXPRESSION TAG	UNP Q96242
B	31	LYS	-	EXPRESSION TAG	UNP Q96242
B	32	THR	-	EXPRESSION TAG	UNP Q96242
B	33	SER	-	EXPRESSION TAG	UNP Q96242
B	137	LEU	PHE	ENGINEERED	UNP Q96242
B	519	HIS	-	EXPRESSION TAG	UNP Q96242
B	520	HIS	-	EXPRESSION TAG	UNP Q96242
B	521	HIS	-	EXPRESSION TAG	UNP Q96242
B	522	HIS	-	EXPRESSION TAG	UNP Q96242

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	398	Total 398	O 398	0	0
4	B	370	Total 370	O 370	0	0

### 3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Cytochrome P450 74A

Chain A:



#### • Molecule 1: Cytochrome P450 74A

Chain B:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.45Å 105.28Å 162.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.39 – 1.73 32.54 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.2 (88.39-1.73) 99.2 (32.54-1.73)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 1.73Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.172 , 0.198 0.173 , 0.197	Depositor DCC
$R_{free}$ test set	5714 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.4	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 51.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 114097 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8384	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.61	0/3870	0.63	0/5232
1	B	0.59	0/3838	0.65	2/5192 (0.0%)
All	All	0.60	0/7708	0.64	2/10424 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	353	ARG	NE-CZ-NH1	10.83	125.72	120.30
1	B	353	ARG	NE-CZ-NH2	-7.95	116.32	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3778	0	3773	12	0
1	B	3746	0	3749	28	0
2	A	43	0	30	0	0
2	B	43	0	30	4	0
3	B	6	0	8	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	398	0	0	3	0
4	B	370	0	0	4	0
All	All	8384	0	7590	42	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (42) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:72:ARG:HH22	1:B:105:ALA:HB2	1.22	1.02
1:B:72:ARG:NH2	1:B:105:ALA:HB2	1.90	0.85
1:A:432[A]:ARG:HH11	1:A:437:VAL:HG21	1.44	0.80
1:A:291:GLU:HG3	4:A:958:HOH:O	1.84	0.77
1:A:432[A]:ARG:NH1	1:A:437:VAL:HG21	2.02	0.74
1:B:93[B]:ASN:ND2	4:B:758:HOH:O	2.21	0.74
2:B:600:HEM:HBC2	2:B:600:HEM:HHD	1.70	0.73
1:B:425:ARG:NH2	4:B:736:HOH:O	2.22	0.71
1:B:349:ALA:O	1:B:353:ARG:HG3	1.90	0.70
1:B:154:LEU:H	1:B:313:ASN:HD21	1.41	0.68
1:B:255:LEU:O	1:B:269:HIS:HE1	1.79	0.65
1:A:339:ARG:HH21	1:A:499:VAL:HG21	1.63	0.64
1:A:72:ARG:HH12	1:A:105:ALA:HB2	1.62	0.64
1:B:105:ALA:HB3	4:B:927:HOH:O	1.98	0.64
1:A:339:ARG:NH2	1:A:499:VAL:HG21	2.13	0.63
1:B:282:TYR:OH	1:B:312:HIS:HD2	1.82	0.63
1:B:60:ASN:HD22	1:B:62:GLY:H	1.50	0.59
1:B:353:ARG:HD3	1:B:490:ARG:HA	1.85	0.59
1:B:164:HIS:HE1	2:B:600:HEM:O2D	1.86	0.58
1:B:154:LEU:H	1:B:313:ASN:ND2	2.03	0.57
1:B:293:ALA:HB1	1:B:296[A]:ILE:CG2	2.40	0.51
1:B:164:HIS:HD2	4:B:652:HOH:O	1.94	0.51
1:B:257:ILE:HG22	1:B:258:GLY:H	1.77	0.50
1:B:174:LEU:HD13	1:B:296[B]:ILE:HG13	1.95	0.48
1:B:109:GLU:CD	1:B:109:GLU:H	2.16	0.48
1:B:173:PHE:HD2	1:B:296[A]:ILE:CD1	2.28	0.47
1:A:149:GLY:HA3	1:A:312:HIS:CE1	2.51	0.46
1:B:239:ALA:HB3	1:B:240:PRO:HD3	1.98	0.45
1:A:128:VAL:HG13	4:A:764:HOH:O	2.16	0.45
1:A:315:LEU:C	1:A:315:LEU:HD23	2.37	0.44
1:A:187[B]:GLN:HG2	4:A:966:HOH:O	2.15	0.44
1:B:259:LEU:HB2	1:B:264:GLU:CG	2.47	0.44
2:B:600:HEM:CHD	2:B:600:HEM:HBC2	2.44	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:377[A]:VAL:HG12	1:B:438:PRO:HB3	1.99	0.43
1:B:293:ALA:O	1:B:296[A]:ILE:HG22	2.19	0.43
1:B:245:LYS:HD3	1:B:281:ASP:OD2	2.18	0.43
1:B:65:ILE:O	1:B:69:ILE:HG12	2.19	0.43
1:B:255:LEU:O	1:B:269:HIS:CE1	2.66	0.42
1:A:169:ASN:HD22	1:A:169:ASN:HA	1.77	0.41
1:B:259:LEU:HB2	1:B:264:GLU:HG3	2.03	0.41
1:A:335:LYS:HA	1:A:497:ILE:CD1	2.51	0.40
1:B:457:ASN:HB2	2:B:600:HEM:HAA1	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	472/495 (95%)	461 (98%)	11 (2%)	0	100	100
1	B	468/495 (94%)	459 (98%)	8 (2%)	1 (0%)	56	35
All	All	940/990 (95%)	920 (98%)	19 (2%)	1 (0%)	59	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	257	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	409/426 (96%)	406 (99%)	3 (1%)	91	83
1	B	405/426 (95%)	399 (98%)	6 (2%)	76	59
All	All	814/852 (96%)	805 (99%)	9 (1%)	84	70

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	201	LEU
1	A	491	ARG
1	A	515	LYS
1	B	52	LEU
1	B	109	GLU
1	B	179	ARG
1	B	201	LEU
1	B	257	ILE
1	B	463	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	100	ASN
1	A	180	ASN
1	A	332	ASN
1	A	359	ASN
1	A	452	HIS
1	B	60	ASN
1	B	100	ASN
1	B	164	HIS
1	B	269	HIS
1	B	312	HIS
1	B	313	ASN
1	B	332	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	600	1	49,50,50	2.51	15 (30%)	46,82,82	2.41	11 (23%)
3	GOL	B	1	-	5,5,5	0.27	0	5,5,5	0.37	0
2	HEM	B	600	1	49,50,50	3.01	17 (34%)	46,82,82	2.29	9 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	600	1	-	0/14/114/114	0/0/8/8
3	GOL	B	1	-	-	0/4/4/4	0/0/0/0
2	HEM	B	600	1	-	0/14/114/114	0/0/8/8

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	600	HEM	C2B-C1B	13.13	1.47	1.44
2	A	600	HEM	C2B-C1B	7.66	1.46	1.44
2	A	600	HEM	C3C-C2C	-6.01	1.33	1.43
2	B	600	HEM	C3C-C2C	-5.64	1.33	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	HEM	C3B-C2B	-5.29	1.34	1.43
2	B	600	HEM	C3B-CAB	5.03	1.56	1.40
2	A	600	HEM	C3D-C4D	4.99	1.45	1.44
2	B	600	HEM	C2D-C1D	4.90	1.45	1.44
2	B	600	HEM	C3B-C2B	-4.82	1.35	1.43
2	B	600	HEM	C3C-CAC	4.78	1.55	1.40
2	A	600	HEM	C3C-CAC	4.78	1.55	1.40
2	B	600	HEM	C4A-C3A	4.70	1.46	1.40
2	A	600	HEM	C3B-CAB	4.50	1.54	1.40
2	A	600	HEM	C4A-C3A	4.17	1.45	1.40
2	B	600	HEM	C3D-C2D	4.11	1.50	1.43
2	B	600	HEM	FE-NA	3.81	2.08	1.92
2	A	600	HEM	C3D-C2D	3.73	1.50	1.43
2	B	600	HEM	FE-ND	3.26	2.09	1.97
2	B	600	HEM	C3D-C4D	-3.16	1.43	1.44
2	A	600	HEM	FE-NA	3.16	2.06	1.92
2	A	600	HEM	FE-NB	3.01	2.08	1.97
2	B	600	HEM	FE-NC	2.98	2.09	1.97
2	A	600	HEM	FE-ND	2.89	2.08	1.97
2	A	600	HEM	FE-NC	2.72	2.08	1.97
2	A	600	HEM	CMD-C2D	2.59	1.55	1.47
2	B	600	HEM	CMB-C2B	2.56	1.55	1.47
2	B	600	HEM	FE-NB	2.49	2.06	1.97
2	A	600	HEM	CMB-C2B	2.43	1.55	1.47
2	B	600	HEM	CHB-C1B	2.35	1.39	1.35
2	B	600	HEM	CMC-C2C	2.34	1.54	1.47
2	A	600	HEM	CMC-C2C	2.31	1.54	1.47
2	B	600	HEM	CMD-C2D	2.21	1.54	1.47

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	HEM	C3B-C4B-NB	-9.51	107.20	114.00
2	B	600	HEM	C3B-C4B-NB	-7.83	108.40	114.00
2	A	600	HEM	C4D-ND-C1D	6.68	111.99	105.16
2	B	600	HEM	C4D-ND-C1D	6.22	111.53	105.16
2	B	600	HEM	CHD-C1D-ND	5.78	129.39	124.58
2	B	600	HEM	CBD-CAD-C3D	-5.07	103.30	114.37
2	A	600	HEM	C1B-NB-C4B	5.02	110.30	105.16
2	A	600	HEM	CBD-CAD-C3D	-4.40	104.78	114.37
2	A	600	HEM	CHC-C4B-NB	4.03	127.94	124.58
2	A	600	HEM	CHD-C1D-ND	3.93	127.85	124.58

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	600	HEM	C1B-NB-C4B	3.91	109.16	105.16
2	A	600	HEM	C2D-C1D-ND	-3.83	108.41	112.93
2	B	600	HEM	C2D-C1D-ND	-3.75	108.50	112.93
2	B	600	HEM	CHD-C4C-NC	-3.52	121.67	124.73
2	A	600	HEM	CHC-C1C-NC	-2.93	122.19	124.73
2	A	600	HEM	CMA-C3A-C4A	-2.72	124.43	128.62
2	A	600	HEM	CHA-C4D-ND	2.42	127.63	124.31
2	B	600	HEM	C1A-CHA-C4D	-2.17	124.62	127.47
2	B	600	HEM	C4A-C3A-C2A	2.15	108.49	107.00
2	A	600	HEM	CAD-C3D-C4D	2.00	128.13	124.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	465/495 (93%)	0.10	18 (3%) 37 44	17, 22, 34, 59	0
1	B	465/495 (93%)	0.14	19 (4%) 35 43	16, 24, 35, 54	0
All	All	930/990 (93%)	0.12	37 (3%) 36 44	16, 23, 35, 59	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	504	LEU	10.4
1	B	503	PRO	9.0
1	A	504	LEU	9.0
1	A	500	GLY	7.8
1	A	501	THR	7.1
1	B	501	THR	7.0
1	A	502	SER	6.9
1	B	502	SER	6.3
1	A	503	PRO	6.1
1	A	259	LEU	5.7
1	A	505	GLY	5.7
1	B	506	SER	5.4
1	A	65	ILE	5.1
1	B	258	GLY	4.9
1	A	257[A]	ILE	4.9
1	B	505	GLY	4.4
1	A	507	SER	3.9
1	A	63	LEU	3.7
1	A	499	VAL	3.6
1	B	259	LEU	3.6
1	B	500	GLY	3.3
1	B	448	LYS	2.9
1	A	506	SER	2.9
1	B	431	ASP	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	105	ALA	2.7
1	B	257	ILE	2.6
1	B	263	ILE	2.6
1	A	262	VAL	2.5
1	B	516	ALA	2.5
1	A	64	PRO	2.5
1	A	263	ILE	2.4
1	B	444	GLU	2.3
1	A	258	GLY	2.2
1	B	445	GLU	2.2
1	A	444	GLU	2.1
1	B	405	ASP	2.0
1	B	343	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	1	6/6	0.20	7.02	23,28,29,31	0
2	HEM	B	600	43/43	0.13	0.58	14,16,19,28	0
2	HEM	A	600	43/43	0.13	0.54	14,16,19,26	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.