



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 02:59 AM GMT

PDB ID : 2RD5
Title : Structural basis for the regulation of N-acetylglutamate kinase by PII in *Arabidopsis thaliana*
Authors : Mizuno, Y.; Moorhead, G.B.G.; Ng, K.K.S.
Deposited on : 2007-09-21
Resolution : 2.51 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	298	
1	B	298	
2	C	135	
2	D	135	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	MG	B	2002	-	X
3	MG	D	1001	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6328 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylglutamate kinase-like protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	0	0	0
			2074	1309	360	395	10			
1	B	283	Total	C	N	O	S	0	0	0
			2087	1317	362	398	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	INITIATING METHIONINE	UNP Q9SCL7
B	0	MET	-	INITIATING METHIONINE	UNP Q9SCL7

- Molecule 2 is a protein called PII protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	125	Total	C	N	O	S	0	0	0
			963	605	171	185	2			
2	D	126	Total	C	N	O	S	0	0	0
			971	610	172	186	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MET	-	INITIATING METHIONINE	UNP Q9ZST4
D	0	MET	-	INITIATING METHIONINE	UNP Q9ZST4

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ARGININE (three-letter code: ARG) (formula: $C_6H_{15}N_4O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			12	6	4	2		
4	B	1	Total	C	N	O	0	0
			12	6	4	2		

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
5	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 6 is N-ACETYL-L-GLUTAMATE (three-letter code: NLG) (formula: $C_7H_{11}NO_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			13	7	1	5		
6	B	1	Total	C	N	O	0	0
			13	7	1	5		

- Molecule 7 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
7	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	24	Total	O	0	0
			24	24		
8	B	25	Total	O	0	0
			25	25		
8	C	4	Total	O	0	0
			4	4		
8	D	11	Total	O	0	0
			11	11		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	171.13Å 171.13Å 171.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.51 32.93 – 2.51	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.00-2.51) 99.9 (32.93-2.51)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.23 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.201 , 0.229 0.213 , 0.236	Depositor DCC
R_{free} test set	2907 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	49.6	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 59.8	EDS
Estimated twinning fraction	0.031 for l,-k,h	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 57272 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6328	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP, ADP, NLG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2099	0.55	1/2841 (0.0%)
1	B	0.38	0/2113	0.57	0/2861
2	C	0.41	0/975	0.55	0/1307
2	D	0.39	0/983	0.57	0/1317
All	All	0.39	0/6170	0.56	1/8326 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	230	LEU	CA-CB-CG	5.25	127.38	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2074	0	2168	20	0
1	B	2087	0	2180	14	0
2	C	963	0	985	6	1
2	D	971	0	994	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	12	0	12	2	0
4	B	12	0	12	0	0
5	A	27	0	12	0	0
5	B	27	0	12	0	0
6	A	13	0	8	0	0
6	B	13	0	8	0	0
7	C	31	0	12	4	1
7	D	31	0	12	5	0
8	A	24	0	0	0	0
8	B	25	0	0	0	0
8	C	4	0	0	1	0
8	D	11	0	0	2	0
All	All	6328	0	6415	52	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (52) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:C:7:ILE:HG23	2:C:8:PRO:HD3	1.40	1.03
2:D:48:GLY:HA3	7:D:1000:ATP:O1A	1.66	0.94
1:A:74:HIS:HE1	1:A:117:ASN:HD22	1.16	0.92
2:C:48:GLY:HA3	7:C:1000:ATP:O1A	1.70	0.91
1:A:38:ILE:HD12	1:A:68:LEU:HD23	1.55	0.87
1:B:74:HIS:HE1	1:B:117:ASN:HD22	1.22	0.81
1:B:30:ILE:HG23	1:B:66:VAL:HG13	1.70	0.72
1:A:30:ILE:HG23	1:A:66:VAL:HG13	1.73	0.70
1:B:106:GLU:O	1:B:110:MET:HG2	1.93	0.67
1:B:222:GLU:HB2	1:B:231:ILE:HD11	1.77	0.66
2:D:99:GLY:HA2	7:D:1000:ATP:O2A	1.96	0.65
2:C:99:GLY:HA2	7:C:1000:ATP:O2A	1.98	0.63
1:A:285:ILE:HG22	1:A:286:MET:CE	2.30	0.61
1:A:74:HIS:CE1	1:A:117:ASN:HD22	2.08	0.59
2:C:7:ILE:CG2	2:C:8:PRO:HD3	2.24	0.57
2:D:77:VAL:HB	2:D:81:GLN:HG3	1.88	0.55
1:A:38:ILE:HG12	1:A:210:LYS:HB2	1.90	0.53
1:B:59:ASP:OD1	1:B:279:HIS:HD2	1.91	0.53
2:D:49:ALA:CB	2:D:64:GLU:HA	2.38	0.53
1:B:222:GLU:HB3	1:B:229:SER:HB2	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:285:ILE:HG22	1:A:286:MET:HE3	1.91	0.53
1:A:232:LYS:O	4:A:1000:ARG:NH2	2.44	0.51
2:D:102:LYS:HG3	7:D:1000:ATP:C8	2.46	0.51
1:A:285:ILE:HG22	1:A:286:MET:HE2	1.93	0.51
1:A:42:TYR:HB2	1:A:72:LEU:HD21	1.92	0.50
1:A:222:GLU:HB3	1:A:229:SER:HB2	1.93	0.49
1:B:253:ILE:HB	1:B:254:PRO:HD3	1.94	0.49
1:A:62:LEU:HD23	1:A:282:LEU:HD21	1.96	0.48
1:A:59:ASP:OD1	1:A:279:HIS:HD2	1.97	0.48
1:A:45:ALA:HA	1:A:48:THR:HG22	1.97	0.47
1:B:214:LEU:HD22	1:B:273:ILE:HD11	1.96	0.47
2:C:35:GLY:O	2:C:129:ASP:HB2	2.14	0.47
1:B:53:LYS:O	1:B:57:VAL:HG13	2.16	0.45
1:A:62:LEU:HD22	1:A:282:LEU:HD11	1.98	0.45
1:B:38:ILE:HD12	1:B:210:LYS:HB2	1.97	0.45
1:B:220:ILE:HD12	1:B:293:THR:HG21	1.99	0.45
1:A:240:LYS:HE3	8:D:1004:HOH:O	2.17	0.44
2:D:118:GLU:O	2:D:123:ALA:HB2	2.18	0.44
2:D:46:GLY:HA3	7:D:1000:ATP:O2'	2.18	0.44
2:C:102:LYS:HD2	7:C:1000:ATP:N7	2.33	0.43
7:D:1000:ATP:H5'2	8:D:1012:HOH:O	2.17	0.43
7:C:1000:ATP:H5'2	8:C:1005:HOH:O	2.19	0.43
1:B:74:HIS:CE1	1:B:117:ASN:HD22	2.15	0.43
1:A:74:HIS:HD2	1:A:75:GLY:O	2.01	0.43
1:B:59:ASP:OD2	1:B:279:HIS:HA	2.19	0.42
1:A:234:ILE:O	1:A:295:ILE:HA	2.20	0.42
2:D:49:ALA:HB2	2:D:64:GLU:HA	2.02	0.41
2:D:20:ARG:HA	2:D:21:PRO:HD3	1.94	0.41
1:A:284:GLU:O	4:A:1000:ARG:HB3	2.21	0.41
2:D:41:VAL:HA	2:D:72:LYS:O	2.21	0.40
1:B:284:GLU:HA	1:B:290:GLY:HA2	2.03	0.40
1:A:226:ASP:HA	1:A:227:PRO:HD3	1.77	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:C:115:ARG:NH1	7:C:1000:ATP:O1G[9_555]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/298 (94%)	274 (98%)	4 (1%)	1 (0%)	43	66
1	B	281/298 (94%)	275 (98%)	5 (2%)	1 (0%)	43	66
2	C	123/135 (91%)	117 (95%)	5 (4%)	1 (1%)	27	46
2	D	124/135 (92%)	118 (95%)	4 (3%)	2 (2%)	14	23
All	All	807/866 (93%)	784 (97%)	18 (2%)	5 (1%)	33	55

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	51	GLY
1	A	76	GLY
1	B	76	GLY
2	D	64	GLU
2	D	128	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/239 (94%)	214 (95%)	11 (5%)	35	59
1	B	227/239 (95%)	217 (96%)	10 (4%)	39	64
2	C	104/114 (91%)	103 (99%)	1 (1%)	85	97
2	D	105/114 (92%)	103 (98%)	2 (2%)	69	90
All	All	661/706 (94%)	637 (96%)	24 (4%)	47	73

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	TYR
1	A	34	ARG
1	A	58	SER
1	A	66	VAL
1	A	88	ASN
1	A	147	VAL
1	A	161	ARG
1	A	168	ARG
1	A	230	LEU
1	A	263	LEU
1	A	282	LEU
1	B	23	LEU
1	B	27	LEU
1	B	49	SER
1	B	66	VAL
1	B	72	LEU
1	B	123	LEU
1	B	145	ARG
1	B	168	ARG
1	B	230	LEU
1	B	263	LEU
2	C	31	LEU
2	D	32	LEU
2	D	80	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	HIS
1	A	119	ASN
1	A	278	GLN
1	A	279	HIS
1	B	74	HIS
1	B	152	GLN
1	B	279	HIS
2	C	81	GLN
2	C	87	ASN
2	D	87	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 3 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	ARG	A	1000	-	11,11,11	1.42	1 (9%)	13,13,13	0.83	0
5	ADP	A	2000	-	29,29,29	1.05	2 (6%)	45,45,45	1.80	8 (17%)
6	NLG	A	2001	-	12,12,12	3.42	3 (25%)	15,15,15	1.54	2 (13%)
4	ARG	B	1000	-	11,11,11	0.45	0	13,13,13	4.24	2 (15%)
5	ADP	B	2000	3	29,29,29	1.14	2 (6%)	45,45,45	1.84	8 (17%)
6	NLG	B	2001	-	12,12,12	3.36	3 (25%)	15,15,15	1.44	2 (13%)
7	ATP	C	1000	3	33,33,33	1.15	3 (9%)	52,52,52	1.82	10 (19%)
7	ATP	D	1000	3	33,33,33	1.08	2 (6%)	52,52,52	1.74	9 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ARG	A	1000	-	-	0/11/11/11	0/0/0/0
5	ADP	A	2000	-	-	0/16/32/32	0/1/3/3
6	NLG	A	2001	-	-	0/13/13/13	0/0/0/0
4	ARG	B	1000	-	-	0/11/11/11	0/0/0/0
5	ADP	B	2000	3	-	0/16/32/32	0/1/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NLG	B	2001	-	-	0/13/13/13	0/0/0/0
7	ATP	C	1000	3	-	0/22/38/38	0/1/3/3
7	ATP	D	1000	3	-	0/22/38/38	0/1/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	2001	NLG	C8-C7	-8.71	1.32	1.50
6	B	2001	NLG	C8-C7	-8.38	1.32	1.50
6	B	2001	NLG	CA-C	-6.29	1.35	1.52
6	A	2001	NLG	CA-C	-6.13	1.36	1.52
6	A	2001	NLG	CA-N2	-4.60	1.35	1.45
4	A	1000	ARG	OXT-C	-4.60	1.14	1.30
6	B	2001	NLG	CA-N2	-4.43	1.35	1.45
5	A	2000	ADP	C5-C4	3.36	1.48	1.40
7	C	1000	ATP	C5-C4	3.26	1.47	1.40
5	B	2000	ADP	C5-C4	3.21	1.47	1.40
7	D	1000	ATP	C5-C4	3.09	1.47	1.40
7	C	1000	ATP	C4-N9	-2.91	1.33	1.37
7	C	1000	ATP	O4'-C1'	2.65	1.45	1.41
7	D	1000	ATP	C4-N9	-2.55	1.34	1.37
5	B	2000	ADP	C4-N9	-2.53	1.34	1.37
5	A	2000	ADP	C4-N9	-2.03	1.34	1.37

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1000	ARG	OXT-C-O	-11.05	99.08	124.07
4	B	1000	ARG	OXT-C-CA	10.36	140.12	116.88
5	B	2000	ADP	N3-C2-N1	-6.69	123.12	128.71
5	A	2000	ADP	N3-C2-N1	-6.18	123.54	128.71
7	C	1000	ATP	N3-C2-N1	-6.06	123.64	128.71
7	D	1000	ATP	N3-C2-N1	-5.93	123.75	128.71
5	B	2000	ADP	N3-C4-N9	5.63	135.60	125.43
7	C	1000	ATP	N3-C4-N9	5.60	135.55	125.43
7	D	1000	ATP	N3-C4-N9	5.57	135.49	125.43
5	A	2000	ADP	N3-C4-N9	5.54	135.43	125.43
6	A	2001	NLG	CB-CA-C	3.72	118.72	110.71
7	C	1000	ATP	PA-O3A-PB	-3.65	120.97	131.68
6	A	2001	NLG	C-CA-N2	3.45	119.42	110.53
5	A	2000	ADP	C5-C4-N3	-3.41	118.28	125.70
5	B	2000	ADP	C5-C4-N3	-3.32	118.48	125.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	1000	ATP	C5-C4-N3	-3.28	118.56	125.70
6	B	2001	NLG	CB-CA-C	3.23	117.66	110.71
7	C	1000	ATP	O5'-C5'-C4'	3.22	120.76	108.94
7	C	1000	ATP	C5-C4-N3	-3.20	118.74	125.70
5	A	2000	ADP	C4-C5-N7	-3.18	106.80	109.52
7	D	1000	ATP	PA-O3A-PB	-3.10	122.60	131.68
7	D	1000	ATP	C4-C5-N7	-3.10	106.87	109.52
5	A	2000	ADP	C3'-C2'-C1'	3.05	105.67	100.91
7	C	1000	ATP	C4-C5-N7	-2.91	107.03	109.52
5	B	2000	ADP	C4-C5-N7	-2.89	107.04	109.52
5	A	2000	ADP	PA-O3A-PB	-2.70	123.77	131.68
7	D	1000	ATP	PB-O3B-PG	-2.67	123.86	131.68
5	B	2000	ADP	C2-N3-C4	2.56	121.29	114.01
5	A	2000	ADP	O4'-C1'-N9	2.53	110.79	108.44
5	A	2000	ADP	C2-N3-C4	2.52	121.18	114.01
7	D	1000	ATP	C2-N3-C4	2.44	120.94	114.01
7	C	1000	ATP	O4'-C4'-C5'	2.43	118.03	109.36
7	D	1000	ATP	O5'-C5'-C4'	2.41	117.77	108.94
7	C	1000	ATP	C8-N9-C4	2.40	108.73	106.90
7	C	1000	ATP	C2-N3-C4	2.40	120.83	114.01
5	B	2000	ADP	O4'-C1'-N9	2.37	110.64	108.44
5	B	2000	ADP	C3'-C2'-C1'	2.36	104.59	100.91
5	B	2000	ADP	PA-O3A-PB	-2.33	124.84	131.68
6	B	2001	NLG	C-CA-N2	2.34	116.55	110.53
7	D	1000	ATP	O4'-C4'-C5'	2.21	117.26	109.36
7	C	1000	ATP	PB-O3B-PG	-2.04	125.70	131.68

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/298 (94%)	-0.11	10 (3%) 41 42	52, 64, 93, 110	0
1	B	283/298 (94%)	-0.08	8 (2%) 50 53	54, 63, 84, 115	0
2	C	125/135 (92%)	0.02	7 (5%) 24 24	55, 62, 80, 104	0
2	D	126/135 (93%)	0.09	8 (6%) 19 19	56, 65, 84, 124	0
All	All	815/866 (94%)	-0.05	33 (4%) 35 37	52, 64, 90, 124	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	130	MET	8.0
2	C	128	GLY	6.7
1	A	225	GLU	6.1
2	D	7	ILE	5.8
2	C	7	ILE	5.6
1	B	18	TYR	5.1
2	D	128	GLY	4.9
2	C	129	ASP	4.5
1	A	18	TYR	4.0
2	D	129	ASP	4.0
2	C	52	GLY	3.7
1	B	15	SER	3.6
1	B	16	PRO	3.3
1	A	247	LYS	3.0
1	B	151	ALA	2.9
2	D	14	LYS	2.8
1	A	244	GLU	2.6
2	D	106	LEU	2.6
1	A	241	LYS	2.5
1	B	148	PRO	2.4
2	C	127	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	243	ILE	2.3
1	B	73	VAL	2.3
1	B	150	SER	2.3
2	D	5	ASP	2.2
1	A	251	GLY	2.2
1	B	179	ILE	2.2
1	A	224	LYS	2.1
1	A	223	ASN	2.1
2	C	5	ASP	2.1
2	D	52	GLY	2.1
1	A	297	GLY	2.0
2	C	16	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MG	D	1001	1/1	0.19	3.60	64,64,64,64	0
3	MG	B	2002	1/1	0.35	3.00	65,65,65,65	0
5	ADP	A	2000	27/27	0.34	1.70	71,75,95,98	27
4	ARG	B	1000	12/12	0.18	0.74	68,73,74,76	0
3	MG	C	1001	1/1	0.13	0.50	60,60,60,60	0
4	ARG	A	1000	12/12	0.16	0.50	73,77,80,85	0
6	NLG	A	2001	13/13	0.12	-0.04	53,58,64,71	0
7	ATP	C	1000	31/31	0.12	-0.40	43,50,64,73	0
6	NLG	B	2001	13/13	0.10	-0.62	41,51,55,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	ATP	D	1000	31/31	0.11	-0.63	55,60,66,78	0
5	ADP	B	2000	27/27	0.09	-0.86	42,51,55,63	0

6.5 Other polymers ⓘ

There are no such residues in this entry.