



wwPDB X-ray Structure Validation Summary Report i

Mar 1, 2014 – 01:10 AM GMT

PDB ID : 3RI4
Title : Ets1 cooperative binding to widely separated sites on promoter DNA
Authors : Babayeva, N.D.; Mino, K.; Tahirov, T.G.
Deposited on : 2011-04-12
Resolution : 3.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	163	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
1	D	163	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
2	B	16	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
2	E	16	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
3	C	16	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
3	F	16	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3567 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein C-ets-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	136	Total	C	N	O	S	0	0	0
			1117	720	193	200	4			
1	D	136	Total	C	N	O	S	0	0	0
			1117	720	193	200	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	279	MET	-	INITIATING METHIONINE	UNP P14921
A	288	TYR	SER	CONFLICT	UNP P14921
D	279	MET	-	INITIATING METHIONINE	UNP P14921
D	288	TYR	SER	CONFLICT	UNP P14921

- Molecule 2 is a DNA chain called TCR alpha promoter DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	16	Total	C	N	O	P	0	0	0
			321	154	59	93	15			
2	E	16	Total	C	N	O	P	0	0	0
			321	154	59	93	15			

- Molecule 3 is a DNA chain called TCR alpha promoter DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	16	Total	C	N	O	P	0	0	0
			329	157	62	95	15			
3	F	16	Total	C	N	O	P	0	0	0
			329	157	62	95	15			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total 6	O 6	0	0
4	B	6	Total 6	O 6	0	0
4	C	3	Total 3	O 3	0	0
4	D	8	Total 8	O 8	0	0
4	E	6	Total 6	O 6	0	0
4	F	4	Total 4	O 4	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein C-ets-1

Chain A:




- Molecule 1: Protein C-ets-1

Chain D:




- Molecule 2: TCR alpha promoter DNA

Chain B:



- Molecule 2: TCR alpha promoter DNA

Chain E:



- Molecule 3: TCR alpha promoter DNA

Chain C:

C101	A102	G103	A104	G105	G106	A107	T108	G109	T110	G111	G112	C113	T114	T115	C116
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

- Molecule 3: TCR alpha promoter DNA

Chain F: 

C301	A302	G303	A304	G305	G306	A307	T308	G309	T310	G311	G312	C313	T314	T315	C316
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.18Å 98.12Å 53.58Å 90.00° 109.67° 90.00°	Depositor
Resolution (Å)	29.84 – 3.00 29.84 – 2.99	Depositor EDS
% Data completeness (in resolution range)	94.5 (29.84-3.00) 93.2 (29.84-2.99)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	23.37 (at 3.00Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.230 , 0.284 0.243 , 0.295	Depositor DCC
R_{free} test set	493 reflections (5.63%)	DCC
Wilson B-factor (Å ²)	49.9	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 32.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtriage
Outliers	0 of 9359 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3567	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.44	0/1147	0.64	0/1548
1	D	0.48	0/1147	0.67	0/1548
2	B	0.57	0/359	0.83	0/551
2	E	0.57	0/359	0.78	0/551
3	C	0.62	0/369	0.83	1/569 (0.2%)
3	F	0.59	0/369	0.84	0/569
All	All	0.51	0/3750	0.73	1/5336 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	E	0	1
3	C	0	1
3	F	0	1
All	All	0	4

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	104	DA	O4'-C1'-N9	-5.07	104.45	108.00

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	1	DG	Sidechain
3	C	103	DG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	E	201	DG	Sidechain
3	F	303	DG	Sidechain

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1117	0	1112	104	0
1	D	1117	0	1112	103	0
2	B	321	0	181	31	0
2	E	321	0	181	27	0
3	C	329	0	182	28	0
3	F	329	0	182	30	0
4	A	6	0	0	3	0
4	B	6	0	0	5	0
4	C	3	0	0	1	0
4	D	8	0	0	2	0
4	E	6	0	0	0	0
4	F	4	0	0	1	0
All	All	3567	0	2950	300	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 46.

The worst 5 of 300 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:F:312:DG:H1'	3:F:313:DC:H5'	1.30	1.08
3:F:304:DA:H2''	3:F:305:DG:H5''	1.07	1.03
1:A:436:LYS:HB3	1:A:437:PRO:HD3	1.43	0.96
1:D:436:LYS:HB3	1:D:437:PRO:HD3	1.44	0.96
1:D:319:PRO:HB3	1:D:347:ASP:HB2	1.45	0.96

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	134/163 (82%)	124 (92%)	8 (6%)	2 (2%)	15	58
1	D	134/163 (82%)	124 (92%)	9 (7%)	1 (1%)	30	78
All	All	268/326 (82%)	248 (92%)	17 (6%)	3 (1%)	21	67

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	400	ASN
1	D	436	LYS
1	A	436	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/142 (83%)	113 (96%)	5 (4%)	40	83
1	D	118/142 (83%)	113 (96%)	5 (4%)	40	83
All	All	236/284 (83%)	226 (96%)	10 (4%)	40	83

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	422	LEU
1	D	309	ARG
1	D	380	ASN
1	A	421	LEU
1	D	313	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	351	GLN
1	A	419	GLN
1	D	336	GLN
1	D	351	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	136/163 (83%)	-0.13	0 100 100	9, 57, 87, 94	0
1	D	136/163 (83%)	-0.14	0 100 100	12, 51, 82, 89	0
2	B	16/16 (100%)	-0.50	0 100 100	15, 45, 61, 64	0
2	E	16/16 (100%)	-0.56	0 100 100	13, 45, 62, 77	0
3	C	16/16 (100%)	-0.60	0 100 100	19, 36, 53, 57	0
3	F	16/16 (100%)	-0.58	0 100 100	19, 38, 59, 65	0
All	All	336/390 (86%)	-0.22	0 100 100	9, 53, 85, 94	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.