



wwPDB X-ray Structure Validation Summary Report i

Feb 26, 2014 – 04:34 PM GMT

PDB ID : 1RKX
Title : Crystal Structure at 1.8 Angstrom of CDP-D-glucose 4,6-dehydratase from Yersinia pseudotuberculosis
Authors : Vogan, E.M.; Bellamacina, C.; He, X.; Liu, H.W.; Ringe, D.; Petsko, G.A.
Deposited on : 2003-11-23
Resolution : 1.80 Å (reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3513 (1.80-1.80)
Clashscore	79885	4461 (1.80-1.80)
Ramachandran outliers	78287	4404 (1.80-1.80)
Sidechain outliers	78261	4403 (1.80-1.80)
RSRZ outliers	66119	3515 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	357	
1	B	357	
1	C	357	
1	D	357	

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 12159 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CDP-glucose-4,6-dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	349	Total	C	N	O	S	0	0	0
			2785	1778	477	518	12			
1	B	346	Total	C	N	O	S	0	0	0
			2763	1764	474	513	12			
1	C	351	Total	C	N	O	S	0	0	0
			2802	1789	481	520	12			
1	D	347	Total	C	N	O	S	0	0	0
			2771	1770	475	514	12			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	230	Total	O	0	0
			230	230		
3	B	205	Total	O	0	0
			205	205		
3	C	214	Total	O	0	0
			214	214		
3	D	213	Total	O	0	0
			213	213		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CDP-glucose-4,6-dehydratase

Chain A:




- Molecule 1: CDP-glucose-4,6-dehydratase

Chain B:




- Molecule 1: CDP-glucose-4,6-dehydratase

Chain C:



- Molecule 1: CDP-glucose-4,6-dehydratase

Chain D:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	99.90Å 115.87Å 126.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.25 – 1.80 36.25 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.5 (36.25-1.80) 97.6 (36.25-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 1.81Å)	Xtriage
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.229 , 0.266 0.229 , 0.266	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	19.9	Xtriage
Anisotropy	0.427	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 36.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	4 of 133257 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12159	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.50 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 9.8343e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2864	0.57	1/3897 (0.0%)
1	B	0.32	0/2842	0.57	1/3867 (0.0%)
1	C	0.33	0/2883	0.57	1/3923 (0.0%)
1	D	0.33	0/2850	0.58	1/3878 (0.0%)
All	All	0.33	0/11439	0.57	4/15565 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	90	GLN	N-CA-C	-5.54	96.05	111.00
1	B	90	GLN	N-CA-C	-5.52	96.10	111.00
1	A	90	GLN	N-CA-C	-5.48	96.20	111.00
1	D	90	GLN	N-CA-C	-5.39	96.45	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2785	0	2673	85	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2763	0	2651	74	0
1	C	2802	0	2688	83	0
1	D	2771	0	2662	76	0
2	A	44	0	26	0	0
2	B	44	0	26	0	0
2	C	44	0	26	0	0
2	D	44	0	26	0	0
3	A	230	0	0	7	0
3	B	205	0	0	6	0
3	C	214	0	0	4	0
3	D	213	0	0	4	0
All	All	12159	0	10778	305	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 14.

The worst 5 of 305 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:3:ASN:H	1:A:247:GLN:HE22	1.15	0.93
1:A:184:HIS:HD2	1:A:186:THR:H	1.19	0.89
1:D:92:LEU:HD13	1:D:94:ARG:HH21	1.39	0.87
1:A:184:HIS:CD2	1:A:186:THR:H	1.93	0.87
1:D:94:ARG:HH12	1:D:205:ASP:CG	1.81	0.85

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/357 (97%)	339 (98%)	6 (2%)	0	100	100
1	B	342/357 (96%)	335 (98%)	7 (2%)	0	100	100
1	C	347/357 (97%)	341 (98%)	6 (2%)	0	100	100
1	D	343/357 (96%)	332 (97%)	10 (3%)	1 (0%)	50	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1377/1428 (96%)	1347 (98%)	29 (2%)	1 (0%)	59 41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	205	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	293/299 (98%)	285 (97%)	8 (3%)	57 39
1	B	291/299 (97%)	286 (98%)	5 (2%)	73 61
1	C	295/299 (99%)	290 (98%)	5 (2%)	73 61
1	D	292/299 (98%)	283 (97%)	9 (3%)	52 33
All	All	1171/1196 (98%)	1144 (98%)	27 (2%)	63 46

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	274	ASN
1	C	141	TRP
1	D	274	ASN
1	C	90	GLN
1	A	180	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 56 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	333	HIS
1	C	180	ASN
1	D	278	GLN
1	B	335	ASN
1	C	4	ASN

5.3.3 RNA [i](#)

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	A	360	-	48,48,48	2.06	8 (16%)	73,73,73	1.51	10 (13%)
2	NAD	B	360	-	48,48,48	2.08	9 (18%)	73,73,73	1.52	11 (15%)
2	NAD	C	360	-	48,48,48	2.06	6 (12%)	73,73,73	1.55	9 (12%)
2	NAD	D	360	-	48,48,48	2.03	9 (18%)	73,73,73	1.54	9 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	360	-	-	0/30/62/62	0/3/5/5
2	NAD	B	360	-	-	0/30/62/62	0/3/5/5
2	NAD	C	360	-	-	0/30/62/62	0/3/5/5
2	NAD	D	360	-	-	0/30/62/62	0/3/5/5

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	360	NAD	C2N-C3N	7.92	1.50	1.38
2	D	360	NAD	C2N-C3N	7.87	1.50	1.38
2	B	360	NAD	C2N-C3N	7.69	1.49	1.38
2	A	360	NAD	C2N-C3N	7.59	1.49	1.38
2	B	360	NAD	C4N-C3N	7.47	1.52	1.39

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	360	NAD	C5N-C4N-C3N	-5.69	112.93	120.32
2	A	360	NAD	C5N-C4N-C3N	-5.68	112.95	120.32
2	C	360	NAD	C5N-C4N-C3N	-5.62	113.03	120.32
2	D	360	NAD	C5N-C4N-C3N	-5.57	113.09	120.32
2	C	360	NAD	C3N-C7N-N7N	5.07	123.53	117.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	349/357 (97%)	0.11	5 (1%) 72 68	11, 20, 47, 64	0
1	B	346/357 (96%)	0.36	20 (5%) 22 16	12, 23, 60, 81	0
1	C	351/357 (98%)	0.40	18 (5%) 27 21	11, 25, 52, 82	0
1	D	347/357 (97%)	0.29	28 (8%) 12 8	12, 22, 54, 82	0
All	All	1393/1428 (97%)	0.29	71 (5%) 27 21	11, 22, 54, 82	0

The worst 5 of 71 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	222	ILE	8.3
1	B	224	ARG	7.8
1	B	223	ILE	7.7
1	B	289	TRP	6.1
1	A	300	ALA	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAD	D	360	44/44	0.12	-0.08	13,17,19,19	0
2	NAD	C	360	44/44	0.12	-0.33	14,19,21,24	0
2	NAD	A	360	44/44	0.11	-0.35	11,16,19,23	0
2	NAD	B	360	44/44	0.11	-0.55	15,18,21,23	0

6.5 Other polymers

There are no such residues in this entry.