



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 12:31 PM GMT

PDB ID : 1RM1
Title : Structure of a Yeast TFIIA/TBP/TATA-box DNA Complex
Authors : Jin, X.; Gewirth, D.T.; Geiger, J.H.
Deposited on : 2003-11-26
Resolution : 2.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance






The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	D	18	
2	E	18	
3	A	240	
4	B	122	
5	C	286	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4200 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*AP*TP*CP*GP*AP*TP*CP*GP*AP*TP*AP*T
P*AP*AP*AP*AP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	18	Total	C	N	O	P	0	0	0
			368	177	72	102	17			

- Molecule 2 is a DNA chain called 5'-D(P*CP*GP*TP*TP*TP*TP*AP*TP*AP*TP*CP*G
P*AP*TP*CP*GP*AP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	18	Total	C	N	O	P	0	0	0
			367	177	60	112	18			

- Molecule 3 is a protein called TATA-box binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	180	Total	C	N	O	S	0	0	0
			1416	921	242	247	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INITIATING METHIONINE	UNP P13393

- Molecule 4 is a protein called Transcription initiation factor IIA small chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	101	Total	C	N	O	S	0	0	0
			792	500	132	156	4			

- Molecule 5 is a protein called Transcription initiation factor IIA large chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	C	116	Total 956	C 599	N 159	O 195	S 3	0	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	119	Total 119	O 119	0	0
6	B	62	Total 62	O 62	0	0
6	C	72	Total 72	O 72	0	0
6	D	24	Total 24	O 24	0	0
6	E	24	Total 24	O 24	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-D(*AP*TP*CP*GP*AP*TP*CP*GP*AP*TP*AP*TP*AP*AP*AP*CP*G)-3'

Chain D: 




- Molecule 2: 5'-D(P*CP*GP*TP*TP*TP*TP*AP*TP*AP*TP*CP*GP*AP*TP*CP*GP*AP*T)-3'

Chain E: 




- Molecule 3: TATA-box binding protein

Chain A: 



- Molecule 4: Transcription initiation factor IIA small chain

Chain B: 



- Molecule 5: Transcription initiation factor IIA large chain

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	173.25Å 173.25Å 164.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.50 28.09 – 2.51	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.50) 90.1 (28.09-2.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.09 (at 2.51Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.186 , 0.252 0.191 , 0.250	Depositor DCC
R_{free} test set	2731 reflections (11.23%)	DCC
Wilson B-factor (Å ²)	40.1	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 29228 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4200	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.50	0/414	0.81	0/637
2	E	0.46	0/409	0.76	0/629
3	A	0.39	0/1443	0.61	0/1942
4	B	0.40	0/800	0.62	0/1080
5	C	0.38	0/970	0.56	0/1310
All	All	0.41	0/4036	0.65	0/5598

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	10	DT	Sidechain
1	D	15	DA	Sidechain

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the

Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	368	0	204	28	0
2	E	367	0	207	17	0
3	A	1416	0	1493	52	0
4	B	792	0	806	42	0
5	C	956	0	916	40	0
6	A	119	0	0	5	0
6	B	62	0	0	4	0
6	C	72	0	0	10	0
6	D	24	0	0	3	0
6	E	24	0	0	2	0
All	All	4200	0	3626	163	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 22.

The worst 5 of 163 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:9:DA:H2''	1:D:10:DT:H5'	1.36	1.03
4:B:11:ARG:HD2	6:B:138:HOH:O	1.62	0.99
4:B:87:VAL:HG12	4:B:88:GLU:H	1.32	0.91
4:B:86:THR:HG22	4:B:105:VAL:HG22	1.54	0.90
3:A:145:LYS:HA	3:A:145:LYS:HE3	1.55	0.89

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	178/240 (74%)	170 (96%)	5 (3%)	3 (2%)	14	22
4	B	97/122 (80%)	93 (96%)	4 (4%)	0	100	100
5	C	110/286 (38%)	103 (94%)	7 (6%)	0	100	100
All	All	385/648 (59%)	366 (95%)	16 (4%)	3 (1%)	27	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	154	ASP
3	A	78	CYS
3	A	110	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	152/205 (74%)	143 (94%)	9 (6%)	28	48
4	B	91/108 (84%)	83 (91%)	8 (9%)	14	26
5	C	107/260 (41%)	104 (97%)	3 (3%)	56	82
All	All	350/573 (61%)	330 (94%)	20 (6%)	29	50

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	B	19	LEU
4	B	23	LEU
4	B	117	ASN
3	A	233	VAL
3	A	234	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
4	B	84	GLN
4	B	117	ASN
5	C	59	GLN
4	B	57	GLN
5	C	43	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	18/18 (100%)	0.80	6 (33%) 1 1	24, 63, 90, 90	0
2	E	18/18 (100%)	1.27	7 (38%) 1 0	21, 69, 90, 90	0
3	A	180/240 (75%)	-0.38	0 100 100	15, 33, 67, 90	0
4	B	101/122 (82%)	-0.23	1 (0%) 79 81	18, 32, 68, 90	0
5	C	116/286 (40%)	-0.06	11 (9%) 8 8	19, 35, 89, 90	0
All	All	433/684 (63%)	-0.14	25 (5%) 22 23	15, 34, 88, 90	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	C	232	SER	4.2
5	C	236	GLU	3.9
2	E	34	DG	3.9
1	D	3	DC	3.9
5	C	229	TYR	3.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.