



# wwPDB X-ray Structure Validation Summary Report

Nov 26, 2014 – 12:42 PM EST

PDB ID : 4RS6  
Title : Crystal structure of the C domain of Polo like Kinase II in Homo Sapiens  
Authors : Shan, H.; Quan, J.; Wang, T.  
Deposited on : 2014-11-07  
Resolution : 2.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1439  
EDS : stable24195  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.1.3  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable24195

# 1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1718 (2.60-2.60)
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)
RSRZ outliers	66119	1718 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	242	
1	B	242	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3462 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-proteinkinase PLK2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	213	Total	C	N	O	S	0	0	0
			1724	1100	283	327	14			
1	B	213	Total	C	N	O	S	0	0	0
			1724	1100	283	327	14			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	444	GLY	-	EXPRESSION TAG	UNP Q9NYY3
A	445	PRO	-	EXPRESSION TAG	UNP Q9NYY3
A	446	LEU	-	EXPRESSION TAG	UNP Q9NYY3
A	447	GLY	-	EXPRESSION TAG	UNP Q9NYY3
A	448	SER	-	EXPRESSION TAG	UNP Q9NYY3
A	449	PRO	-	EXPRESSION TAG	UNP Q9NYY3
A	450	GLU	-	EXPRESSION TAG	UNP Q9NYY3
B	444	GLY	-	EXPRESSION TAG	UNP Q9NYY3
B	445	PRO	-	EXPRESSION TAG	UNP Q9NYY3
B	446	LEU	-	EXPRESSION TAG	UNP Q9NYY3
B	447	GLY	-	EXPRESSION TAG	UNP Q9NYY3
B	448	SER	-	EXPRESSION TAG	UNP Q9NYY3
B	449	PRO	-	EXPRESSION TAG	UNP Q9NYY3
B	450	GLU	-	EXPRESSION TAG	UNP Q9NYY3

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	11	Total	O	0	0
			11	11		
2	B	3	Total	O	0	0
			3	3		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-proteinkinase PLK2

Chain A: 



- Molecule 1: Serine/threonine-proteinkinase PLK2

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	153.07Å 153.07Å 153.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.83 – 2.60 24.83 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (24.83-2.60) 100.0 (24.83-2.60)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	0.15	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.31 (at 2.60Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.186 , 0.247 0.210 , 0.252	Depositor DCC
$R_{free}$ test set	947 reflections (5.13%)	DCC
Wilson B-factor (Å <sup>2</sup> )	62.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 42.7	EDS
Estimated twinning fraction	0.027 for -l,-k,-h	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 18474 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3462	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.80	0/1764	0.88	2/2391 (0.1%)
1	B	0.97	3/1764 (0.2%)	0.97	6/2391 (0.3%)
All	All	0.89	3/3528 (0.1%)	0.92	8/4782 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	586	GLY	CA-C	-18.02	1.23	1.51
1	B	585	ASP	CA-CB	-10.01	1.31	1.53
1	B	585	ASP	N-CA	-8.57	1.29	1.46

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	585	ASP	N-CA-CB	9.74	128.14	110.60
1	B	586	GLY	N-CA-C	9.03	135.67	113.10
1	B	588	ASP	N-CA-C	8.23	133.22	111.00
1	B	480	ARG	NE-CZ-NH1	7.41	124.00	120.30
1	A	597	ARG	N-CA-C	-6.80	92.64	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	586	GLY	Mainchain

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1724	0	1679	37	0
1	B	1724	0	1679	36	0
2	A	11	0	0	0	0
2	B	3	0	0	0	0
All	All	3462	0	3358	73	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 11.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:592:VAL:CG2	1:A:595:ILE:HD11	1.54	1.35
1:A:592:VAL:CG2	1:A:595:ILE:CD1	2.24	1.16
1:A:592:VAL:HG23	1:A:595:ILE:CD1	1.82	1.06
1:B:550:LEU:H	1:B:550:LEU:HD13	1.24	1.00
1:A:596:ARG:HH11	1:A:596:ARG:HB3	1.27	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/242 (87%)	196 (93%)	15 (7%)	0	100	100
1	B	211/242 (87%)	196 (93%)	15 (7%)	0	100	100
All	All	422/484 (87%)	392 (93%)	30 (7%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/218 (89%)	173 (89%)	21 (11%)	9	17
1	B	194/218 (89%)	175 (90%)	19 (10%)	12	21
All	All	388/436 (89%)	348 (90%)	40 (10%)	10	19

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	644	THR
1	B	498	SER
1	B	643	LEU
1	A	681	LEU
1	B	500	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	523	HIS
1	A	545	HIS
1	B	577	HIS
1	B	671	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	213/242 (88%)	-0.10	9 (4%) 35 31	24, 63, 96, 136	0
1	B	213/242 (88%)	0.14	16 (7%) 14 11	45, 70, 122, 152	0
All	All	426/484 (88%)	0.02	25 (5%) 22 18	24, 67, 112, 152	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	595	ILE	5.6
1	A	596	ARG	5.6
1	A	597	ARG	5.0
1	B	491	CYS	4.9
1	A	593	THR	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

### 6.5 Other polymers ⓘ

There are no such residues in this entry.