



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 04:24 AM GMT

PDB ID : 3RV3
Title : Crystal structure of E.coli biotin carboxylase in complex with two ADP and one Mg ion
Authors : Chou, C.Y.; Tong, L.
Deposited on : 2011-05-05
Resolution : 1.91 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4387 (1.94-1.90)
Clashscore	79885	5258 (1.94-1.90)
Ramachandran outliers	78287	5193 (1.94-1.90)
Sidechain outliers	78261	5194 (1.94-1.90)
RSRZ outliers	66119	4389 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	452	
1	B	452	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 7740 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Biotin carboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	0	0	0
			3441	2166	616	637	22			
1	B	447	Total	C	N	O	S	0	0	0
			3442	2168	615	637	22			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1C	GLY	-	EXPRESSION TAG	UNP P24182
A	1B	SER	-	EXPRESSION TAG	UNP P24182
A	1A	HIS	-	EXPRESSION TAG	UNP P24182
B	1C	GLY	-	EXPRESSION TAG	UNP P24182
B	1B	SER	-	EXPRESSION TAG	UNP P24182
B	1A	HIS	-	EXPRESSION TAG	UNP P24182

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	367	Total	O	0	0
			367	367		
4	B	380	Total	O	0	0
			380	380		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Biotin carboxylase



- Molecule 1: Biotin carboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	171.36Å 58.01Å 85.24Å 90.00° 94.64° 90.00°	Depositor
Resolution (Å)	30.00 – 1.91 29.43 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.4 (30.00-1.91) 99.4 (29.43-1.91)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.81 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.208 , 0.263 0.208 , 0.263	Depositor DCC
R_{free} test set	3281 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	18.2	Xtriage
Anisotropy	0.525	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 33.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	1 of 64727 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7740	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/3505	0.62	1/4730 (0.0%)
1	B	0.51	0/3505	0.61	0/4730
All	All	0.52	0/7010	0.61	1/9460 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	154	TYR	N-CA-C	5.75	126.54	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	154	TYR	Peptide

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit,

and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3441	0	3460	46	1
1	B	3442	0	3467	46	1
2	A	54	0	24	1	0
2	B	54	0	24	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	367	0	0	17	0
4	B	380	0	0	15	0
All	All	7740	0	6975	94	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (94) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:288:GLU:HG2	4:A:684:HOH:O	1.38	1.21
1:A:157:ILE:HD13	4:A:668:HOH:O	1.41	1.18
1:B:397:VAL:HB	4:B:820:HOH:O	1.46	1.14
1:B:407:GLN:HG2	4:B:466:HOH:O	1.66	0.94
1:B:190:LYS:HA	1:B:194:SER:O	1.68	0.93
1:A:18:LEU:HD23	4:A:743:HOH:O	1.77	0.84
1:A:342:GLU:HG3	4:A:752:HOH:O	1.76	0.84
1:B:158:ILE:CG2	1:B:198:VAL:HG11	2.09	0.82
1:A:108:GLU:HG2	4:A:689:HOH:O	1.80	0.80
1:B:234:ARG:NH1	1:B:234:ARG:HB3	1.96	0.79
1:A:156:VAL:HB	4:A:710:HOH:O	1.82	0.79
1:A:200:MET:HG2	4:A:710:HOH:O	1.82	0.79
1:A:19:ARG:HG3	4:A:743:HOH:O	1.84	0.77
1:B:181:SER:O	1:B:185:THR:HG23	1.87	0.74
1:B:234:ARG:NH2	4:B:591:HOH:O	2.18	0.72
2:B:1003:ADP:O2B	4:B:771:HOH:O	2.07	0.72
1:B:234:ARG:HH11	1:B:234:ARG:HB3	1.55	0.70
2:A:1002:ADP:O2A	4:A:684:HOH:O	2.11	0.68
1:A:200:MET:CG	4:A:710:HOH:O	2.41	0.67
1:A:116:LYS:O	1:A:120:ILE:HG12	1.94	0.67
1:A:22:LYS:HE2	4:B:619:HOH:O	1.95	0.65
1:A:18:LEU:CD2	1:A:22:LYS:HE3	2.26	0.64
1:A:27:LYS:NZ	4:A:734:HOH:O	2.30	0.63
1:A:363:PHE:HD1	4:A:596:HOH:O	1.81	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:205:GLU:O	1:B:438:HIS:HD2	1.83	0.61
1:A:370:HIS:CD2	1:A:370:HIS:H	2.18	0.61
1:B:147:ALA:O	1:B:151:ARG:HG3	2.00	0.61
1:A:23:GLU:OE1	1:A:310:LYS:HE3	1.99	0.61
1:B:158:ILE:HG23	1:B:198:VAL:HG11	1.83	0.61
1:A:18:LEU:CD2	4:A:743:HOH:O	2.40	0.60
1:B:252:LEU:HD12	4:B:565:HOH:O	2.01	0.60
1:B:255:TYR:O	1:B:259:ARG:HG2	2.04	0.58
1:B:403:LYS:O	1:B:407:GLN:HG3	2.03	0.58
1:B:158:ILE:CG2	1:B:198:VAL:CG1	2.79	0.58
1:A:205:GLU:O	1:A:438:HIS:CD2	2.56	0.58
1:A:155:PRO:HD2	1:A:202:LYS:HE3	1.87	0.57
1:A:234:ARG:HB3	1:A:234:ARG:HH11	1.70	0.57
1:B:158:ILE:HG23	1:B:198:VAL:CG1	2.34	0.57
1:B:254:ARG:HG3	4:B:648:HOH:O	2.05	0.56
1:A:228:ARG:HH21	1:A:294:GLN:NE2	2.03	0.56
1:A:180:GLN:O	1:A:184:MET:HG3	2.06	0.56
1:A:93:GLU:OE1	1:A:111:ARG:NH1	2.39	0.55
1:A:18:LEU:HD21	1:A:22:LYS:HE3	1.88	0.54
1:B:394:ASN:N	4:B:820:HOH:O	2.17	0.53
1:B:158:ILE:HG22	1:B:198:VAL:HG11	1.88	0.53
1:A:346:THR:OG1	1:A:348:LEU:HG	2.10	0.52
1:A:18:LEU:HD23	1:A:22:LYS:HE3	1.91	0.51
1:B:234:ARG:CB	1:B:234:ARG:HH11	2.22	0.51
1:B:228:ARG:HH21	1:B:294:GLN:NE2	2.09	0.51
1:A:234:ARG:HB3	1:A:234:ARG:NH1	2.26	0.51
1:A:205:GLU:O	1:A:438:HIS:HD2	1.93	0.50
1:B:251:GLU:HG3	4:B:468:HOH:O	2.12	0.50
1:A:141:ASP:HB2	4:A:495:HOH:O	2.13	0.49
1:A:23:GLU:OE1	1:A:310:LYS:CE	2.60	0.49
1:B:295:VAL:HG23	4:B:771:HOH:O	2.13	0.48
1:B:57:VAL:HG13	1:B:58:LYS:HG3	1.95	0.48
1:B:250:PRO:O	1:B:254:ARG:HG2	2.14	0.48
1:A:169:MET:HE3	4:A:668:HOH:O	2.13	0.47
1:B:307:ASP:O	1:B:311:GLU:HG2	2.14	0.47
1:B:344:PRO:HG3	1:B:416:THR:O	2.15	0.47
1:B:228:ARG:HH21	1:B:294:GLN:HE21	1.62	0.47
1:B:347:PHE:HE1	1:B:420:LEU:HD22	1.79	0.47
1:B:113:MET:CE	1:B:267:ILE:HG21	2.44	0.47
1:B:206:ASN:N	1:B:207:PRO:HD3	2.30	0.46
1:B:292:ARG:NH1	4:B:771:HOH:O	2.48	0.46
1:B:258:GLU:HG2	4:B:488:HOH:O	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:263:ALA:O	1:A:267:ILE:HG12	2.16	0.45
1:A:157:ILE:HD12	1:A:171:VAL:HG22	1.98	0.45
1:A:82:TYR:CZ	1:A:295:VAL:HG22	2.52	0.45
1:B:206:ASN:N	1:B:207:PRO:CD	2.80	0.45
1:A:93:GLU:HG3	1:A:107:ALA:CB	2.46	0.45
1:B:249:THR:OG1	1:B:251:GLU:HG2	2.16	0.44
1:A:131:VAL:CG1	1:A:204:LEU:HD11	2.47	0.44
1:B:253:ARG:NH1	4:B:813:HOH:O	2.51	0.44
1:B:259:ARG:NE	1:B:259:ARG:HA	2.33	0.44
1:B:82:TYR:CZ	1:B:295:VAL:HG22	2.52	0.44
1:A:442:LYS:HG3	4:A:548:HOH:O	2.18	0.44
1:A:296:GLU:O	1:A:299:VAL:HG22	2.18	0.43
1:B:2:LEU:HA	1:B:76:VAL:HG21	1.99	0.43
1:B:113:MET:HE2	1:B:267:ILE:HG21	1.98	0.43
1:B:205:GLU:HG3	4:B:531:HOH:O	2.18	0.42
1:B:228:ARG:NH2	1:B:294:GLN:HE21	2.17	0.42
1:A:105:PRO:HG2	1:A:291:THR:HB	2.01	0.42
1:B:13:ILE:HB	1:B:82:TYR:CE2	2.55	0.42
1:A:228:ARG:NH2	1:A:294:GLN:HE21	2.17	0.42
1:A:131:VAL:HG13	1:A:204:LEU:HD11	2.01	0.42
1:A:13:ILE:HB	1:A:82:TYR:CE2	2.55	0.42
1:A:157:ILE:HG13	1:A:170:ARG:O	2.19	0.41
1:A:440:LEU:O	1:A:444:LEU:HG	2.20	0.41
1:A:417:ASN:HB3	4:A:752:HOH:O	2.20	0.41
1:B:353:LYS:HA	1:B:376:THR:HA	2.03	0.41
1:B:234:ARG:HG2	1:B:235:ARG:HD2	2.02	0.41
1:A:27:LYS:HB2	1:A:27:LYS:HE3	1.74	0.41
1:B:205:GLU:N	4:B:756:HOH:O	2.53	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:98:SER:CB	1:B:251:GLU:OE2[4.546]	1.95	0.25

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/452 (98%)	434 (98%)	10 (2%)	1 (0%)	56	44
1	B	445/452 (98%)	429 (96%)	15 (3%)	1 (0%)	56	44
All	All	890/904 (98%)	863 (97%)	25 (3%)	2 (0%)	56	44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	TYR
1	B	195	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/363 (99%)	357 (99%)	2 (1%)	92	92
1	B	359/363 (99%)	352 (98%)	7 (2%)	69	62
All	All	718/726 (99%)	709 (99%)	9 (1%)	80	77

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	154	TYR
1	A	363	PHE
1	B	5	ILE
1	B	278	LEU
1	B	345	ASN
1	B	428	GLU
1	B	432	HIS
1	B	442	LYS
1	B	444	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	290	ASN
1	A	294	GLN
1	A	370	HIS
1	A	407	GLN
1	A	426	ASN
1	A	432	HIS
1	A	438	HIS
1	B	206	ASN
1	B	236	HIS
1	B	237	GLN
1	B	294	GLN
1	B	340	ASN
1	B	404	ASN
1	B	431	GLN
1	B	438	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADP	A	1000	-	29,29,29	1.08	3 (10%)	45,45,45	1.70	6 (13%)
2	ADP	A	1002	3	29,29,29	1.16	2 (6%)	45,45,45	1.92	9 (20%)
2	ADP	B	1001	3	29,29,29	1.06	2 (6%)	45,45,45	1.84	10 (22%)
2	ADP	B	1003	-	29,29,29	1.06	2 (6%)	45,45,45	1.80	8 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	1000	-	-	0/16/32/32	0/1/3/3
2	ADP	A	1002	3	-	0/16/32/32	0/1/3/3
2	ADP	B	1001	3	-	0/16/32/32	0/1/3/3
2	ADP	B	1003	-	-	0/16/32/32	0/1/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1002	ADP	C4-N9	-3.26	1.33	1.37
2	A	1002	ADP	C5-C4	3.16	1.47	1.40
2	A	1000	ADP	C5-C4	2.98	1.47	1.40
2	B	1003	ADP	C5-C4	2.96	1.47	1.40
2	B	1001	ADP	C4-N9	-2.92	1.33	1.37
2	B	1003	ADP	C4-N9	-2.92	1.33	1.37
2	B	1001	ADP	C5-C4	2.85	1.46	1.40
2	A	1000	ADP	C4-N9	-2.82	1.33	1.37
2	A	1000	ADP	C2-N3	2.30	1.36	1.32

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1003	ADP	N3-C2-N1	-7.29	122.61	128.71
2	B	1001	ADP	N3-C2-N1	-7.10	122.77	128.71
2	A	1002	ADP	N3-C2-N1	-6.83	123.00	128.71
2	A	1000	ADP	N3-C2-N1	-6.45	123.32	128.71
2	A	1000	ADP	N3-C4-N9	5.35	135.10	125.43
2	B	1003	ADP	N3-C4-N9	5.02	134.49	125.43
2	B	1001	ADP	N3-C4-N9	4.82	134.14	125.43
2	A	1002	ADP	N3-C4-N9	4.68	133.88	125.43
2	A	1002	ADP	O4'-C1'-N9	4.19	112.34	108.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1002	ADP	C2'-C1'-N9	-3.71	103.75	113.27
2	A	1002	ADP	C8-N9-C4	3.57	109.62	106.90
2	B	1003	ADP	O4'-C1'-N9	3.31	111.52	108.44
2	A	1000	ADP	C4-C5-N7	-3.27	106.72	109.52
2	A	1000	ADP	C5-C4-N3	-3.13	118.89	125.70
2	B	1001	ADP	C4-C5-N7	-2.95	106.99	109.52
2	B	1001	ADP	C8-N9-C4	2.91	109.12	106.90
2	B	1001	ADP	O4'-C1'-N9	2.80	111.05	108.44
2	B	1003	ADP	C5-C4-N3	-2.78	119.64	125.70
2	B	1001	ADP	C5-C4-N3	-2.63	119.97	125.70
2	B	1003	ADP	C8-N9-C4	2.50	108.81	106.90
2	A	1002	ADP	C1'-N9-C4	-2.46	122.39	126.64
2	B	1003	ADP	C4-C5-N7	-2.41	107.46	109.52
2	A	1000	ADP	C2-N3-C4	2.40	120.83	114.01
2	B	1001	ADP	C2'-C1'-N9	-2.39	107.14	113.27
2	B	1003	ADP	C2-N3-C4	2.39	120.80	114.01
2	A	1002	ADP	C2-N1-C6	2.33	122.97	118.77
2	B	1001	ADP	C2-N3-C4	2.30	120.55	114.01
2	A	1002	ADP	C5-C4-N3	-2.29	120.72	125.70
2	B	1001	ADP	C1'-N9-C4	-2.24	122.77	126.64
2	A	1002	ADP	C4-C5-N7	-2.22	107.62	109.52
2	A	1000	ADP	C8-N9-C4	2.19	108.57	106.90
2	B	1001	ADP	C2-N1-C6	2.13	122.62	118.77
2	B	1003	ADP	C2-N1-C6	2.06	122.50	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	447/452 (98%)	0.47	18 (4%) 36 37	7, 16, 29, 35	1 (0%)
1	B	447/452 (98%)	0.69	35 (7%) 13 13	9, 18, 34, 43	1 (0%)
All	All	894/904 (98%)	0.58	53 (5%) 22 22	7, 17, 32, 43	2 (0%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	447	GLN	7.2
1	B	445	GLY	6.9
1	B	446	LEU	6.3
1	A	154	TYR	4.4
1	B	189	ALA	4.2
1	B	322	SER	3.8
1	B	193	PHE	3.7
1	B	141	ASP	3.5
1	B	194	SER	3.4
1	B	255	TYR	3.3
1	A	348	LEU	3.2
1	A	345	ASN	3.1
1	A	203	TYR	3.1
1	A	346	THR	3.0
1	B	127	GLY	2.9
1	B	432	HIS	2.9
1	A	141	ASP	2.9
1	B	173	ARG	2.9
1	B	206	ASN	2.9
1	A	192	ALA	2.9
1	B	150	LYS	2.8
1	B	1	MET	2.8
1	A	344	PRO	2.8
1	B	323	ILE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	139	GLY	2.8
1	A	206	ASN	2.7
1	B	205	GLU	2.7
1	B	143	ASP	2.7
1	B	128	VAL	2.7
1	B	160	ALA	2.7
1	A	204	LEU	2.5
1	A	148	ILE	2.5
1	B	138	LEU	2.5
1	B	152	ILE	2.4
1	B	140	ASP	2.3
1	B	191	ALA	2.3
1	B	259	ARG	2.3
1	B	142	MET	2.3
1	A	151	ARG	2.3
1	A	347	PHE	2.3
1	A	1(A)	HIS	2.2
1	B	349	PRO	2.2
1	A	191	ALA	2.2
1	B	254	ARG	2.2
1	A	152	ILE	2.2
1	A	322	SER	2.2
1	B	281	ASN	2.2
1	B	185	THR	2.2
1	B	38	ASP	2.1
1	B	345	ASN	2.1
1	B	187	ALA	2.1
1	A	363	PHE	2.0
1	B	329	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ADP	A	1000	27/27	0.12	-	12,16,18,19	0
3	MG	B	1005	1/1	0.07	-	18,18,18,18	0
2	ADP	B	1001	27/27	0.12	-	14,17,20,20	0
3	MG	A	1004	1/1	0.06	-	15,15,15,15	0
2	ADP	B	1003	27/27	0.13	-	12,17,19,20	0
2	ADP	A	1002	27/27	0.14	-	13,21,22,25	0

6.5 Other polymers

There are no such residues in this entry.