



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 04:11 PM GMT

PDB ID : 3RVY
Title : Crystal structure of the NavAb voltage-gated sodium channel (Ile217Cys, 2.7 Å)
Authors : Payandeh, J.; Scheuer, T.; Zheng, N.; Catterall, W.A.
Deposited on : 2011-05-06
Resolution : 2.70 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1557 (2.70-2.70)
Clashscore	79885	1939 (2.70-2.70)
Ramachandran outliers	78287	1905 (2.70-2.70)
Sidechain outliers	78261	1905 (2.70-2.70)
RSRZ outliers	66119	1559 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	285	
1	B	285	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	PX4	A	4015	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3717 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ion transport protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S	0	0	0
			1810	1233	270	294	13			
1	B	221	Total	C	N	O	S	0	0	0
			1810	1233	270	294	13			

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	983	MET	-	EXPRESSION TAG	UNP A8EVM5
A	984	ASP	-	EXPRESSION TAG	UNP A8EVM5
A	985	TYR	-	EXPRESSION TAG	UNP A8EVM5
A	986	LYS	-	EXPRESSION TAG	UNP A8EVM5
A	987	ASP	-	EXPRESSION TAG	UNP A8EVM5
A	988	ASP	-	EXPRESSION TAG	UNP A8EVM5
A	989	ASP	-	EXPRESSION TAG	UNP A8EVM5
A	990	ASP	-	EXPRESSION TAG	UNP A8EVM5
A	991	LYS	-	EXPRESSION TAG	UNP A8EVM5
A	992	GLY	-	EXPRESSION TAG	UNP A8EVM5
A	993	SER	-	EXPRESSION TAG	UNP A8EVM5
A	994	LEU	-	EXPRESSION TAG	UNP A8EVM5
A	995	VAL	-	EXPRESSION TAG	UNP A8EVM5
A	996	PRO	-	EXPRESSION TAG	UNP A8EVM5
A	997	ARG	-	EXPRESSION TAG	UNP A8EVM5
A	998	GLY	-	EXPRESSION TAG	UNP A8EVM5
A	999	SER	-	EXPRESSION TAG	UNP A8EVM5
A	1000	HIS	-	EXPRESSION TAG	UNP A8EVM5
A	1217	CYS	ILE	ENGINEERED MUTATION	UNP A8EVM5
B	1983	MET	-	EXPRESSION TAG	UNP A8EVM5
B	1984	ASP	-	EXPRESSION TAG	UNP A8EVM5
B	1985	TYR	-	EXPRESSION TAG	UNP A8EVM5
B	1986	LYS	-	EXPRESSION TAG	UNP A8EVM5
B	1987	ASP	-	EXPRESSION TAG	UNP A8EVM5
B	1988	ASP	-	EXPRESSION TAG	UNP A8EVM5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1989	ASP	-	EXPRESSION TAG	UNP A8EVM5
B	1990	ASP	-	EXPRESSION TAG	UNP A8EVM5
B	1991	LYS	-	EXPRESSION TAG	UNP A8EVM5
B	1992	GLY	-	EXPRESSION TAG	UNP A8EVM5
B	1993	SER	-	EXPRESSION TAG	UNP A8EVM5
B	1994	LEU	-	EXPRESSION TAG	UNP A8EVM5
B	1995	VAL	-	EXPRESSION TAG	UNP A8EVM5
B	1996	PRO	-	EXPRESSION TAG	UNP A8EVM5
B	1997	ARG	-	EXPRESSION TAG	UNP A8EVM5
B	1998	GLY	-	EXPRESSION TAG	UNP A8EVM5
B	1999	SER	-	EXPRESSION TAG	UNP A8EVM5
B	2000	HIS	-	EXPRESSION TAG	UNP A8EVM5
B	2217	CYS	ILE	ENGINEERED MUTATION	UNP A8EVM5

- Molecule 2 is 1,2-DIMYRISTOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PX4) (formula: C₃₆H₇₃NO₈P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O P 22 14 7 1	0	0
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total C 5 5	0	0
2	A	1	Total C 6 6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total C 5 5	0	0
2	B	1	Total C O P 10 3 6 1	0	0
2	B	1	Total C O P 21 13 7 1	0	0
2	B	1	Total C O P 10 3 6 1	0	0
2	B	1	Total C 6 6	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O 1 1	0	0
3	B	1	Total O 1 1	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ion transport protein

Chain A:



- Molecule 1: Ion transport protein

Chain B:



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	125.59Å 125.48Å 192.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.70 42.27 – 2.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.70) 86.8 (42.27-2.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 2.69Å)	Xtriage
Refinement program	CNS 1.1 & 1.3	Depositor
R, R_{free}	0.266 , 0.273 0.279 , 0.286	Depositor DCC
R_{free} test set	1816 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	66.2	Xtriage
Anisotropy	0.535	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 71.7	EDS
Estimated twinning fraction	0.479 for k,h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 36494 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3717	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PX4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/1861	0.61	1/2531 (0.0%)
1	B	0.47	0/1861	0.61	0/2531
All	All	0.47	0/3722	0.61	1/5062 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1221	MET	CG-SD-CE	-7.49	88.21	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1810	0	1883	124	0
1	B	1810	0	1883	126	0
2	A	48	0	48	0	0
2	B	47	0	40	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3717	0	3854	244	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 32.

The worst 5 of 244 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:1217:CYS:SG	1:B:2214:VAL:HG22	2.00	1.01
1:B:2059:GLU:OE2	1:B:2108:ARG:NH2	1.94	0.99
1:B:2096:GLU:CD	1:B:2099:ARG:HD2	1.83	0.98
1:A:1001:MET:HG3	1:A:1004:ARG:HD3	1.48	0.95
1:B:2096:GLU:HG3	1:B:2099:ARG:HB3	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/285 (77%)	194 (89%)	25 (11%)	0	100	100
1	B	219/285 (77%)	188 (86%)	30 (14%)	1 (0%)	38	70
All	All	438/570 (77%)	382 (87%)	55 (13%)	1 (0%)	56	86

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2002	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/264 (77%)	197 (97%)	6 (3%)	53	84
1	B	203/264 (77%)	192 (95%)	11 (5%)	31	61
All	All	406/528 (77%)	389 (96%)	17 (4%)	40	73

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2028	THR
1	B	2029	MET
1	B	2133	VAL
1	B	2005	ILE
1	B	2165	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	2172	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PX4	A	4001	-	19,21,45	1.43	2 (10%)	23,25,53	1.05	0
2	PX4	A	4003	-	4,4,45	1.78	2 (50%)	6,6,53	1.91	2 (33%)
2	PX4	A	4005	-	4,4,45	1.21	1 (25%)	3,3,53	0.39	0
2	PX4	A	4007	-	5,5,45	1.06	1 (20%)	4,4,53	0.29	0
2	PX4	A	4013	-	4,4,45	1.82	2 (50%)	6,6,53	1.83	2 (33%)
2	PX4	A	4015	-	4,4,45	1.98	1 (25%)	3,3,53	0.46	0
2	PX4	B	4002	-	9,9,45	1.49	1 (11%)	12,12,53	11.03	5 (41%)
2	PX4	B	4011	-	19,20,45	1.69	2 (10%)	23,24,53	6.92	2 (8%)
2	PX4	B	4012	-	9,9,45	1.57	1 (11%)	12,12,53	11.76	5 (41%)
2	PX4	B	4017	-	5,5,45	0.97	0	4,4,53	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PX4	A	4001	-	-	0/24/24/49	0/0/0/0
2	PX4	A	4003	-	-	0/0/0/49	0/0/0/0
2	PX4	A	4005	-	-	0/2/2/49	0/0/0/0
2	PX4	A	4007	-	-	0/3/3/49	0/0/0/0
2	PX4	A	4013	-	-	0/0/0/49	0/0/0/0
2	PX4	A	4015	-	-	0/2/2/49	0/0/0/0
2	PX4	B	4002	-	-	0/8/8/49	0/0/0/0
2	PX4	B	4011	-	-	0/22/22/49	0/0/0/0
2	PX4	B	4012	-	-	0/8/8/49	0/0/0/0
2	PX4	B	4017	-	-	0/3/3/49	0/0/0/0

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	4011	PX4	C17-C16	-5.16	1.53	1.55
2	B	4011	PX4	P1-O3	4.64	1.70	1.61
2	B	4012	PX4	P1-O3	4.14	1.64	1.50
2	A	4001	PX4	C17-C16	-4.14	1.53	1.55
2	A	4015	PX4	C32-C33	-3.91	1.53	1.55

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4012	PX4	O7-C7-C6	-30.30	108.22	119.54
2	B	4002	PX4	O7-C7-C8	-26.92	109.49	119.54
2	B	4012	PX4	O7-C7-C8	-26.84	109.52	119.54
2	B	4002	PX4	O7-C7-C6	-26.70	109.57	119.54
2	B	4011	PX4	O7-C7-C8	-24.22	110.49	119.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	221/285 (77%)	0.45	13 (5%) 22 23	43, 87, 162, 215	0
1	B	221/285 (77%)	0.45	11 (4%) 28 30	43, 85, 157, 235	0
All	All	442/570 (77%)	0.45	24 (5%) 24 27	43, 87, 160, 235	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2095	PHE	4.7
1	B	2094	GLY	4.6
1	B	2098	LEU	4.0
1	A	1095	PHE	4.0
1	A	1220	ALA	3.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	PX4	A	4015	5/46	0.28	2.53	92,92,92,92	0
2	PX4	A	4007	6/46	0.24	1.95	66,66,66,66	0
2	PX4	B	4011	21/46	0.22	1.69	45,96,96,96	0
2	PX4	A	4001	22/46	0.21	0.80	45,89,89,89	0
2	PX4	B	4002	10/46	0.21	0.22	124,124,124,124	0
2	PX4	B	4017	6/46	0.19	-0.04	64,64,64,64	0
2	PX4	A	4013	5/46	0.18	-0.30	132,132,132,132	0
2	PX4	A	4003	5/46	0.16	-0.63	127,127,127,127	0
2	PX4	B	4012	10/46	0.17	-0.70	117,117,117,117	0
2	PX4	A	4005	5/46	0.20	-0.95	91,91,91,91	0

6.5 Other polymers ⓘ

There are no such residues in this entry.