



# wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 05:16 AM GMT

PDB ID : 1RZZ  
Title : PHOTOSYNTHETIC REACTION CENTER DOUBLE MUTANT FROM RHODOBACTER SPHAEROIDES WITH ASP L213 REPLACED WITH ASN AND ARG M233 REPLACED WITH CYS IN THE CHARGE-NEUTRAL DQAQB STATE (TETRAGONAL FORM)  
Authors : Xu, Q.; Axelrod, H.L.; Abresch, E.C.; Paddock, M.L.; Okamura, M.Y.; Feher, G.  
Deposited on : 2003-12-29  
Resolution : 2.40 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	L	281	
1	R	281	
2	M	307	
2	S	307	
3	H	260	
3	T	260	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	BCL	R	2002	-	X
4	BCL	S	2001	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Geometry	Electron density
4	BCL	S	2003	-	X
4	BCL	S	2004	-	X
5	U10	L	1009	-	X
5	U10	R	2009	-	X
5	U10	S	2008	-	X
8	SPO	M	1010	-	X
8	SPO	S	2010	-	X
9	LDA	M	1011	-	X
9	LDA	M	1012	-	X
9	LDA	M	1013	-	X
9	LDA	M	1014	-	X
9	LDA	S	2011	-	X

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 14481 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reaction center protein L chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	281	Total	C	N	O	S	0	0	0
			2232	1507	356	361	8			
1	R	281	Total	C	N	O	S	0	0	0
			2232	1507	356	361	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	213	ASN	ASP	ENGINEERED	UNP P02954
R	213	ASN	ASP	ENGINEERED	UNP P02954

- Molecule 2 is a protein called Reaction center protein M chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	299	Total	C	N	O	S	0	0	0
			2385	1594	388	392	11			
2	S	299	Total	C	N	O	S	0	0	0
			2385	1594	388	392	11			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	233	CYS	ARG	ENGINEERED	UNP P02953
S	233	CYS	ARG	ENGINEERED	UNP P02953

- Molecule 3 is a protein called Reaction center protein H chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	246	Total	C	N	O	S	0	0	0
			1869	1196	320	343	10			
3	T	246	Total	C	N	O	S	0	0	0
			1869	1196	320	343	10			

- Molecule 4 is BACTERIOCHLOROPHYLL A (three-letter code: BCL) (formula:  $C_{55}H_{74}MgN_4O_6$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	L	1	Total	C	Mg	N	O	0	0
			51	40	1	4	6		
4	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
4	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
4	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
4	R	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
4	S	1	Total	C	Mg	N	O	0	0
			51	40	1	4	6		
4	S	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
4	S	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		

- Molecule 5 is UBIQUINONE-10 (three-letter code: U10) (formula:  $C_{59}H_{90}O_4$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	C	O	0	0
			44	40	4		
5	M	1	Total	C	O	0	0
			38	34	4		
5	R	1	Total	C	O	0	0
			18	14	4		
5	S	1	Total	C	O	0	0
			32	28	4		

- Molecule 6 is FE (II) ION (three-letter code: FE2) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	S	1	Total	Fe	0	0
			1	1		
6	M	1	Total	Fe	0	0
			1	1		

- Molecule 7 is BACTERIOPHEOPHYTIN A (three-letter code: BPH) (formula: C<sub>55</sub>H<sub>76</sub>N<sub>4</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	M	1	Total	C	N	O	0	0
			55	45	4	6		
7	M	1	Total	C	N	O	0	0
			65	55	4	6		
7	R	1	Total	C	N	O	0	0
			65	55	4	6		
7	S	1	Total	C	N	O	0	0
			55	45	4	6		

- Molecule 8 is SPHEROIDENE (three-letter code: SPO) (formula: C<sub>41</sub>H<sub>60</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	M	1	Total	C	O	0	0
			42	41	1		
8	S	1	Total	C	O	0	0
			42	41	1		

- Molecule 9 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula:  $C_{14}H_{31}NO$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	M	1	Total	C	N	O	0	0
			16	14	1	1		
9	M	1	Total	C	N	O	0	0
			16	14	1	1		
9	M	1	Total	C	N	O	0	0
			16	14	1	1		
9	M	1	Total	C	N	O	0	0
			16	14	1	1		
9	S	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	L	72	Total	O	0	0
			72	72		
10	M	107	Total	O	0	0
			107	107		

*Continued on next page...*




*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	H	109	Total 109	O 109	0	0
10	R	47	Total 47	O 47	0	0
10	S	75	Total 75	O 75	0	0
10	T	63	Total 63	O 63	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Reaction center protein L chain




- Molecule 1: Reaction center protein L chain



- Molecule 2: Reaction center protein M chain




- Molecule 2: Reaction center protein M chain








• Molecule 3: Reaction center protein H chain

Chain H:



• Molecule 3: Reaction center protein H chain

Chain T:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	139.58Å 139.58Å 274.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.76 – 2.40 39.76 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.2 (39.76-2.40) 98.3 (39.76-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.25 (at 2.39Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.216 , 0.238 0.212 , 0.234	Depositor DCC
$R_{free}$ test set	5283 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.9	Xtriage
Anisotropy	0.306	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 38.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	1 of 104640 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	14481	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BCL, LDA, BPH, FE2, SPO, U10

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.42	0/2320	0.57	0/3175
1	R	0.37	0/2320	0.55	0/3175
2	M	0.41	0/2477	0.56	0/3383
2	S	0.39	0/2477	0.53	0/3383
3	H	0.35	0/1917	0.60	0/2608
3	T	0.31	0/1917	0.56	0/2608
All	All	0.38	0/13428	0.56	0/18332

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	2232	0	2189	57	0
1	R	2232	0	2189	60	0
2	M	2385	0	2296	51	0
2	S	2385	0	2296	63	0
3	H	1869	0	1884	36	0
3	T	1869	0	1884	52	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	L	183	0	189	22	0
4	M	66	0	74	13	0
4	R	66	0	74	5	0
4	S	183	0	189	21	0
5	L	44	0	56	3	0
5	M	38	0	47	4	0
5	R	18	0	15	0	0
5	S	32	0	39	1	0
6	M	1	0	0	0	0
6	S	1	0	0	0	0
7	M	120	0	127	6	0
7	R	65	0	74	5	0
7	S	55	0	53	2	0
8	M	42	0	60	4	0
8	S	42	0	60	3	0
9	M	64	0	124	3	0
9	S	16	0	31	0	0
10	H	109	0	0	1	0
10	L	72	0	0	1	0
10	M	107	0	0	2	0
10	R	47	0	0	2	0
10	S	75	0	0	3	0
10	T	63	0	0	1	0
All	All	14481	0	13950	317	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 11.

The worst 5 of 317 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:R:241:VAL:HG21	7:R:2006:BPH:HAC1	1.40	1.03
1:L:241:VAL:HG21	7:M:1006:BPH:HAC2	1.38	1.01
2:M:109:LEU:HD12	2:M:114:LEU:HD13	1.56	0.86
2:S:157:TRP:HB2	4:S:2003:BCL:H62	1.58	0.84
4:S:2001:BCL:HBC1	4:S:2003:BCL:CAD	2.09	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	279/281 (99%)	269 (96%)	10 (4%)	0	100	100
1	R	279/281 (99%)	263 (94%)	14 (5%)	2 (1%)	30	43
2	M	297/307 (97%)	285 (96%)	12 (4%)	0	100	100
2	S	297/307 (97%)	287 (97%)	10 (3%)	0	100	100
3	H	244/260 (94%)	236 (97%)	8 (3%)	0	100	100
3	T	244/260 (94%)	235 (96%)	9 (4%)	0	100	100
All	All	1640/1696 (97%)	1575 (96%)	63 (4%)	2 (0%)	59	78

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	R	4	SER
1	R	31	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	220/220 (100%)	207 (94%)	13 (6%)	28	42
1	R	220/220 (100%)	208 (94%)	12 (6%)	30	46
2	M	235/240 (98%)	220 (94%)	15 (6%)	25	37
2	S	235/240 (98%)	225 (96%)	10 (4%)	40	59
3	H	199/208 (96%)	194 (98%)	5 (2%)	60	80
3	T	199/208 (96%)	196 (98%)	3 (2%)	76	90
All	All	1308/1336 (98%)	1250 (96%)	58 (4%)	39	58

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	M	216	PHE
3	H	231	ASP
2	S	216	PHE
2	M	232	GLU
3	H	123	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	R	87	GLN
1	R	159	ASN
3	T	98	HIS
3	H	206	ASN
3	T	128	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 25 ligands modelled in this entry, 2 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	BCL	L	1001	-	59,59,74	2.42	15 (25%)	77,97,115	2.05	21 (27%)
4	BCL	L	1002	-	74,74,74	2.01	13 (17%)	97,115,115	1.94	27 (27%)
4	BCL	L	1004	-	74,74,74	2.12	13 (17%)	97,115,115	2.00	27 (27%)
5	U10	L	1009	-	44,44,63	2.38	20 (45%)	55,56,79	3.15	16 (29%)
4	BCL	M	1003	-	74,74,74	2.01	12 (16%)	97,115,115	1.97	30 (30%)
7	BPH	M	1005	-	59,60,70	1.71	12 (20%)	80,89,101	2.09	26 (32%)
7	BPH	M	1006	-	70,70,70	1.71	13 (18%)	94,101,101	2.03	30 (31%)
5	U10	M	1008	-	38,38,63	2.04	13 (34%)	47,49,79	2.00	9 (19%)
8	SPO	M	1010	-	41,41,41	3.42	23 (56%)	50,50,50	3.94	17 (34%)
9	LDA	M	1011	-	15,15,15	3.64	2 (13%)	17,17,17	2.50	5 (29%)
9	LDA	M	1012	-	15,15,15	3.93	2 (13%)	17,17,17	2.45	5 (29%)
9	LDA	M	1013	-	15,15,15	3.86	2 (13%)	17,17,17	2.42	5 (29%)
9	LDA	M	1014	-	15,15,15	3.92	2 (13%)	17,17,17	2.49	5 (29%)
4	BCL	R	2002	-	74,74,74	2.10	13 (17%)	97,115,115	1.94	29 (29%)
7	BPH	R	2006	-	70,70,70	1.72	13 (18%)	94,101,101	2.04	27 (28%)
5	U10	R	2009	-	18,18,63	2.20	8 (44%)	23,25,79	2.25	6 (26%)
4	BCL	S	2001	-	59,59,74	2.33	15 (25%)	77,97,115	2.00	21 (27%)
4	BCL	S	2003	-	74,74,74	2.06	12 (16%)	97,115,115	2.04	34 (35%)
4	BCL	S	2004	-	74,74,74	2.15	15 (20%)	97,115,115	1.92	26 (26%)
7	BPH	S	2005	-	59,60,70	1.70	11 (18%)	80,89,101	2.18	28 (35%)
5	U10	S	2008	-	31,32,63	2.13	10 (32%)	39,41,79	1.99	9 (23%)
8	SPO	S	2010	-	41,41,41	3.31	24 (58%)	50,50,50	4.05	17 (34%)
9	LDA	S	2011	-	15,15,15	3.82	2 (13%)	17,17,17	2.48	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BCL	L	1001	-	-	0/23/119/137	0/0/9/9
4	BCL	L	1002	-	-	0/41/137/137	0/0/9/9
4	BCL	L	1004	-	-	0/41/137/137	0/0/9/9
5	U10	L	1009	-	-	1/41/65/87	0/1/1/1
4	BCL	M	1003	-	-	0/41/137/137	0/0/9/9
7	BPH	M	1005	-	-	0/37/93/105	0/0/6/6
7	BPH	M	1006	-	-	1/49/105/105	0/0/6/6

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	U10	M	1008	-	-	0/33/57/87	0/1/1/1
8	SPO	M	1010	-	-	0/47/47/47	0/0/0/0
9	LDA	M	1011	-	-	0/13/13/13	0/0/0/0
9	LDA	M	1012	-	-	0/13/13/13	0/0/0/0
9	LDA	M	1013	-	-	0/13/13/13	0/0/0/0
9	LDA	M	1014	-	-	0/13/13/13	0/0/0/0
4	BCL	R	2002	-	-	0/41/137/137	0/0/9/9
7	BPH	R	2006	-	-	0/49/105/105	0/0/6/6
5	U10	R	2009	-	-	0/9/33/87	0/1/1/1
4	BCL	S	2001	-	-	0/23/119/137	0/0/9/9
4	BCL	S	2003	-	-	0/41/137/137	0/0/9/9
4	BCL	S	2004	-	-	0/41/137/137	0/0/9/9
7	BPH	S	2005	-	-	0/37/93/105	0/0/6/6
5	U10	S	2008	-	-	0/25/50/87	0/1/1/1
8	SPO	S	2010	-	-	0/47/47/47	0/0/0/0
9	LDA	S	2011	-	-	0/13/13/13	0/0/0/0

The worst 5 of 265 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	M	1012	LDA	O1-N1	-14.55	1.25	1.39
9	M	1014	LDA	O1-N1	-14.46	1.25	1.39
9	M	1013	LDA	O1-N1	-14.27	1.26	1.39
9	S	2011	LDA	O1-N1	-14.11	1.26	1.39
9	M	1011	LDA	O1-N1	-13.40	1.26	1.39

The worst 5 of 425 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	S	2010	SPO	C3-C1-C4	-16.32	84.02	110.97
5	L	1009	U10	C32-C33-C34	16.13	162.60	127.80
8	M	1010	SPO	C3-C1-C4	-16.00	84.55	110.97
8	S	2010	SPO	C2-C1-C4	-15.09	86.04	110.97
8	M	1010	SPO	C2-C1-C4	-14.59	86.88	110.97

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	L	1009	U10	C34-C33-C32-C31
7	M	1006	BPH	C1-C2-C3-C4

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	281/281 (100%)	-0.12	4 (1%) 72 71	17, 32, 49, 58	0
1	R	281/281 (100%)	0.04	11 (3%) 37 35	23, 41, 58, 69	0
2	M	299/307 (97%)	-0.27	2 (0%) 84 84	17, 27, 41, 67	0
2	S	299/307 (97%)	-0.14	2 (0%) 84 84	24, 35, 48, 69	0
3	H	246/260 (94%)	-0.14	8 (3%) 44 42	22, 34, 54, 88	0
3	T	246/260 (94%)	0.16	18 (7%) 15 13	31, 45, 70, 80	0
All	All	1652/1696 (97%)	-0.08	45 (2%) 52 49	17, 36, 57, 88	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	252	VAL	5.4
3	H	255	MET	4.8
3	H	254	ALA	4.7
3	H	256	LEU	4.6
1	R	1	ALA	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	U10	R	2009	18/63	0.37	8.08	81,85,86,87	0
9	LDA	M	1011	16/16	0.25	5.08	49,57,74,74	0
5	U10	L	1009	44/63	0.32	4.31	66,72,79,79	0
9	LDA	M	1013	16/16	0.28	4.13	69,71,80,80	0
9	LDA	S	2011	16/16	0.34	3.82	74,79,85,85	0
8	SPO	M	1010	42/42	0.22	3.32	16,28,47,51	0
9	LDA	M	1014	16/16	0.31	3.29	50,55,59,60	0
8	SPO	S	2010	42/42	0.21	3.16	24,35,51,54	0
5	U10	S	2008	32/63	0.29	2.96	36,42,51,54	0
4	BCL	S	2001	51/66	0.18	2.91	28,31,41,43	0
4	BCL	S	2003	66/66	0.19	2.63	27,33,48,55	0
9	LDA	M	1012	16/16	0.26	2.42	60,62,65,65	0
4	BCL	R	2002	66/66	0.17	2.31	29,35,44,48	0
4	BCL	S	2004	66/66	0.20	2.09	26,34,58,60	0
7	BPH	S	2005	55/65	0.17	1.85	25,30,55,58	0
4	BCL	L	1001	51/66	0.21	1.83	17,24,51,55	0
7	BPH	M	1006	65/65	0.22	1.75	23,28,37,40	0
4	BCL	L	1002	66/66	0.18	1.65	20,26,29,33	0
5	U10	M	1008	38/63	0.25	1.60	23,30,53,53	0
4	BCL	L	1004	66/66	0.25	1.57	16,25,45,52	0
7	BPH	R	2006	65/65	0.17	1.48	31,38,45,46	0
4	BCL	M	1003	66/66	0.17	1.42	20,25,39,44	0
7	BPH	M	1005	55/65	0.16	0.78	17,23,43,46	0
6	FE2	S	2007	1/1	0.13	-1.29	30,30,30,30	0
6	FE2	M	1007	1/1	0.12	-2.95	20,20,20,20	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.