



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 01:19 PM GMT

PDB ID : 1S3B
Title : Crystal structure of MAOB in complex with N-methyl-N-propargyl-1(R)-aminindan
Authors : Binda, C.; Hubalek, F.; Li, M.; Herzig, Y.; Sterling, J.; Edmondson, D.E.; Mattevi, A.
Deposited on : 2004-01-13
Resolution : 1.65 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1404 (1.68-1.64)
Clashscore	79885	1001 (1.66-1.66)
Ramachandran outliers	78287	1581 (1.68-1.64)
Sidechain outliers	78261	1580 (1.68-1.64)
RSRZ outliers	66119	1404 (1.68-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	520	
1	B	520	

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 8864 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amine oxidase [flavin-containing] B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	499	Total	C	N	O	S	0	0	0
			3971	2538	681	728	24			
1	B	494	Total	C	N	O	S	0	0	0
			3940	2519	676	721	24			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: RMA, FAD) (formula: $C_{13}H_{15}N$, $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	2	Total	C	N	O	P	0	0
			67	40	10	15	2		
2	B	2	Total	C	N	O	P	0	0
			67	40	10	15	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	376	Total 376	O 376	0	0
3	B	443	Total 443	O 443	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Amine oxidase [flavin-containing] B



• Molecule 1: Amine oxidase [flavin-containing] B



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	130.76 Å 222.91 Å 86.17 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.65 14.96 – 1.65	Depositor EDS
% Data completeness (in resolution range)	98.1 (15.00-1.65) 98.1 (14.96-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 1.65 Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.203 , 0.223 0.207 , 0.226	Depositor DCC
R_{free} test set	3732 reflections (2.59%)	DCC
Wilson B-factor (Å ²)	14.6	Xtriage
Anisotropy	0.301	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 63.1	EDS
Estimated twinning fraction	0.012 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.015 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 147566 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8864	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RMA, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/4068	0.65	2/5522 (0.0%)
1	B	0.36	0/4037	0.65	7/5479 (0.1%)
All	All	0.37	0/8105	0.65	9/11001 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	445	ARG	NE-CZ-NH2	-6.55	117.02	120.30
1	A	315	MET	CG-SD-CE	5.82	109.51	100.20
1	B	315	MET	CG-SD-CE	5.82	109.51	100.20
1	A	445	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	B	227	ASP	CB-CG-OD2	5.38	123.15	118.30
1	B	445	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	B	310	ASP	CB-CG-OD2	5.13	122.92	118.30
1	B	6	ASP	CB-CG-OD2	5.11	122.90	118.30
1	B	330	ASP	CB-CG-OD2	5.08	122.88	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3971	0	3967	23	0
1	B	3940	0	3937	31	1
2	A	67	0	44	4	0
2	B	67	0	44	2	0
3	A	376	0	0	2	0
3	B	443	0	0	5	0
All	All	8864	0	7992	54	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (54) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:92:VAL:HB	1:B:318:ASP:OD2	1.79	0.83
1:A:28:LEU:HD11	1:A:454:MET:HE1	1.64	0.78
1:B:28:LEU:HD11	1:B:454:MET:HE1	1.67	0.75
1:B:117:ASN:HD22	1:B:120:ARG:HH21	1.37	0.72
1:A:117:ASN:HD22	1:A:120:ARG:HH21	1.38	0.71
1:B:92:VAL:CG2	1:B:318:ASP:OD2	2.42	0.68
1:B:414:LEU:HD12	3:B:811:HOH:O	1.95	0.67
1:B:251:ASN:H	1:B:251:ASN:HD22	1.44	0.66
1:B:92:VAL:CB	1:B:318:ASP:OD2	2.44	0.65
1:A:251:ASN:H	1:A:251:ASN:HD22	1.43	0.64
1:A:451:LEU:HA	1:A:454:MET:HE2	1.81	0.60
1:B:451:LEU:HA	1:B:454:MET:HE2	1.83	0.60
1:A:445:ARG:HD2	1:A:463:TRP:CH2	2.39	0.58
1:B:92:VAL:HG23	1:B:318:ASP:OD2	2.04	0.57
1:B:445:ARG:HD3	3:B:616:HOH:O	2.04	0.56
1:B:445:ARG:HD2	1:B:463:TRP:CH2	2.40	0.56
1:A:445:ARG:HD2	1:A:463:TRP:CZ2	2.42	0.54
1:A:445:ARG:HD3	3:A:623:HOH:O	2.11	0.50
1:B:445:ARG:HD2	1:B:463:TRP:CZ2	2.46	0.50
1:A:271:LYS:HE2	3:A:767:HOH:O	2.12	0.50
1:A:233:ARG:HG3	1:A:251:ASN:HD21	1.77	0.50
1:B:233:ARG:HG3	1:B:251:ASN:HD21	1.77	0.50
1:B:168:PHE:CE1	1:B:172:CYS:SG	3.05	0.50
1:A:321:GLU:CD	1:A:321:GLU:H	2.15	0.49
1:B:321:GLU:CD	1:B:321:GLU:H	2.17	0.48
1:A:58:GLY:HA2	2:A:600:FAD:C4X	2.44	0.48
1:A:398:TYR:CZ	2:A:601:RMA:H111	2.48	0.48
1:B:58:GLY:HA2	2:B:600:FAD:C4X	2.46	0.46
1:B:222:MET:HE3	3:B:859:HOH:O	2.15	0.46
1:B:398:TYR:CZ	2:B:601:RMA:H111	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:237:TYR:HB3	1:B:248:GLU:HB3	1.98	0.45
1:A:168:PHE:CE1	1:A:172:CYS:SG	3.09	0.45
1:B:308:LYS:HE2	1:B:308:LYS:HB2	1.88	0.45
1:B:366:GLU:O	1:B:370:LYS:HG3	2.17	0.44
1:A:237:TYR:HB3	1:A:248:GLU:HB3	1.99	0.44
1:A:265:PRO:HD2	1:A:268:LEU:HD12	2.00	0.44
1:B:111:THR:HG22	1:B:115:HIS:CD2	2.52	0.43
1:B:323:PRO:HD2	1:B:367:LEU:HD22	2.00	0.43
1:A:308:LYS:HB2	1:A:308:LYS:HE2	1.87	0.43
1:A:323:PRO:HD2	1:A:367:LEU:HD22	2.00	0.43
1:A:117:ASN:HD22	1:A:120:ARG:NH2	2.12	0.43
1:B:251:ASN:H	1:B:251:ASN:ND2	2.13	0.43
1:A:251:ASN:ND2	1:A:251:ASN:H	2.13	0.43
1:A:172:CYS:SG	2:A:601:RMA:H6	2.59	0.42
1:B:454:MET:HE3	1:B:456:LYS:HG3	2.02	0.42
1:B:489:VAL:N	1:B:490:PRO:HD2	2.35	0.42
1:B:108:ASN:HA	1:B:109:PRO:HD3	1.87	0.42
1:B:309:LYS:NZ	1:B:373:GLY:O	2.53	0.41
1:B:271:LYS:HE2	3:B:858:HOH:O	2.20	0.41
1:A:489:VAL:N	1:A:490:PRO:HD2	2.35	0.41
1:A:23:LEU:HB2	1:A:30:VAL:HG11	2.03	0.41
1:B:494:ARG:NH1	3:B:926:HOH:O	2.54	0.40
1:A:434:GLY:O	2:A:600:FAD:H1'2	2.21	0.40
1:B:23:LEU:HB2	1:B:30:VAL:HG11	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:25:ASP:O	1:B:49:GLN:NE2[4_565]	2.01	0.19

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/520 (96%)	486 (98%)	11 (2%)	0	100	100
1	B	492/520 (95%)	480 (98%)	12 (2%)	0	100	100
All	All	989/1040 (95%)	966 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/444 (96%)	414 (97%)	13 (3%)	53	21
1	B	424/444 (96%)	415 (98%)	9 (2%)	66	37
All	All	851/888 (96%)	829 (97%)	22 (3%)	59	27

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
1	A	93	LYS
1	A	106	VAL
1	A	155	LEU
1	A	251	ASN
1	A	280	MET
1	A	350	ARG
1	A	379	GLU
1	A	397	CYS
1	A	460	ASP
1	A	495	LEU
1	A	498	LEU
1	A	500	THR
1	B	49	GLN
1	B	155	LEU
1	B	242	ARG
1	B	251	ASN
1	B	280	MET
1	B	350	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	379	GLU
1	B	460	ASP
1	B	495	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
1	A	251	ASN
1	B	117	ASN
1	B	170	ASN
1	B	251	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	600	1,2	58,58,58	1.02	6 (10%)	85,89,89	1.85	13 (15%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	RMA	A	601	2	15,15,15	4.10	4 (26%)	20,20,20	10.12	6 (30%)
2	FAD	B	600	1,2	58,58,58	1.01	5 (8%)	85,89,89	1.85	13 (15%)
2	RMA	B	601	2	15,15,15	4.13	4 (26%)	20,20,20	10.03	7 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	600	1,2	-	0/34/50/50	0/1/6/6
2	RMA	A	601	2	-	0/6/16/16	0/0/2/2
2	FAD	B	600	1,2	-	0/34/50/50	0/1/6/6
2	RMA	B	601	2	-	0/6/16/16	0/0/2/2

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	RMA	C11-N10	-11.08	1.29	1.46
2	A	601	RMA	C11-N10	-11.04	1.29	1.46
2	B	601	RMA	C10-N10	-8.66	1.29	1.46
2	A	601	RMA	C10-N10	-8.62	1.29	1.46
2	B	601	RMA	C12-C13	5.47	1.30	1.17
2	A	601	RMA	C12-C13	5.38	1.30	1.17
2	B	601	RMA	C11-C12	4.80	1.54	1.47
2	A	601	RMA	C11-C12	4.66	1.54	1.47
2	A	600	FAD	C1'-C2'	2.94	1.54	1.51
2	A	600	FAD	C2A-N3A	2.78	1.37	1.32
2	B	600	FAD	C2A-N3A	2.68	1.37	1.32
2	B	600	FAD	C1'-N10	2.65	1.51	1.48
2	B	600	FAD	C1'-C2'	2.53	1.54	1.51
2	A	600	FAD	C5X-N5	2.41	1.39	1.35
2	A	600	FAD	C1'-N10	2.34	1.50	1.48
2	B	600	FAD	C2A-N1A	2.31	1.38	1.33
2	A	600	FAD	C2A-N1A	2.30	1.38	1.33
2	B	600	FAD	C5X-N5	2.12	1.38	1.35
2	A	600	FAD	C10-N1	2.03	1.39	1.35

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	RMA	C11-C12-C13	-43.93	125.19	177.80
2	B	601	RMA	C11-C12-C13	-43.46	125.76	177.80
2	A	600	FAD	N3A-C2A-N1A	-10.44	119.98	128.71
2	B	600	FAD	N3A-C2A-N1A	-10.39	120.02	128.71
2	B	601	RMA	C10-N10-C11	7.07	120.27	110.12
2	A	601	RMA	C10-N10-C11	6.86	119.96	110.12
2	A	600	FAD	C9A-N10-C10	-5.81	116.07	121.77
2	B	600	FAD	C9A-N10-C10	-5.58	116.30	121.77
2	A	600	FAD	C2-N1-C10	5.11	120.12	114.98
2	A	601	RMA	C12-C11-N10	5.07	121.69	111.77
2	B	600	FAD	C2-N1-C10	4.95	119.97	114.98
2	B	601	RMA	C12-C11-N10	4.92	121.39	111.77
2	A	601	RMA	C11-N10-C9	3.54	118.63	113.65
2	B	601	RMA	C11-N10-C9	3.51	118.57	113.65
2	B	600	FAD	C4X-C10-N10	-3.29	118.87	120.51
2	B	600	FAD	N3A-C4A-N9A	3.00	130.85	125.43
2	A	600	FAD	C4X-C10-N1	-2.89	119.84	122.73
2	B	600	FAD	C9A-C5X-N5	-2.85	118.00	122.37
2	A	600	FAD	N3A-C4A-N9A	2.80	130.49	125.43
2	A	600	FAD	C5'-C4'-C3'	-2.60	107.16	112.06
2	A	601	RMA	C2-C1-C9	2.59	108.58	106.13
2	A	600	FAD	C4-N3-C2	-2.55	120.15	125.39
2	B	601	RMA	C2-C1-C9	2.49	108.49	106.13
2	B	600	FAD	C5'-C4'-C3'	-2.44	107.46	112.06
2	A	600	FAD	C1B-N9A-C4A	-2.41	122.46	126.64
2	B	601	RMA	C6-C7-C8	-2.41	117.53	120.95
2	A	600	FAD	C4X-C10-N10	-2.40	119.31	120.51
2	B	600	FAD	C4X-C10-N1	-2.40	120.33	122.73
2	A	600	FAD	C2'-C1'-N10	2.34	115.55	112.45
2	B	600	FAD	C2'-C1'-N10	2.33	115.55	112.45
2	A	600	FAD	C9A-C5X-N5	-2.29	118.85	122.37
2	B	600	FAD	C4X-N5-C5X	2.28	119.25	116.69
2	A	601	RMA	C6-C7-C8	-2.26	117.75	120.95
2	A	600	FAD	C4A-C5A-N7A	-2.23	107.61	109.52
2	B	600	FAD	C4-N3-C2	-2.23	120.81	125.39
2	B	600	FAD	N7A-C8A-N9A	-2.20	108.12	114.36
2	A	600	FAD	N7A-C8A-N9A	-2.14	108.30	114.36
2	B	600	FAD	C1B-N9A-C4A	-2.12	122.97	126.64
2	B	601	RMA	C5-C4-C3	-2.10	117.59	120.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	499/520 (95%)	0.31	43 (8%) 11 9	28, 37, 55, 78	0
1	B	494/520 (95%)	0.35	32 (6%) 18 17	28, 37, 52, 72	0
All	All	993/1040 (95%)	0.33	75 (7%) 14 12	28, 37, 53, 78	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	500	THR	8.5
1	B	496	ILE	7.9
1	B	495	LEU	7.8
1	A	499	THR	7.5
1	A	498	LEU	5.9
1	B	494	ARG	5.8
1	A	252	HIS	5.2
1	A	107	TRP	5.1
1	B	103	PHE	4.9
1	B	107	TRP	4.7
1	A	494	ARG	4.6
1	B	243	GLU	4.5
1	A	497	GLY	4.5
1	A	103	PHE	4.5
1	A	501	ILE	4.4
1	A	3	ASN	4.2
1	A	479	THR	4.2
1	B	93	LYS	4.1
1	B	3	ASN	4.0
1	A	480	THR	3.7
1	B	242	ARG	3.7
1	A	460	ASP	3.7
1	B	92	VAL	3.7
1	A	481	PHE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	227	ASP	3.5
1	B	460	ASP	3.3
1	B	119	TRP	3.3
1	B	252	HIS	3.2
1	B	95	LYS	3.2
1	B	481	PHE	3.2
1	B	89	ILE	3.2
1	B	482	LEU	3.2
1	B	246	LEU	3.1
1	A	242	ARG	3.1
1	B	112	TYR	3.0
1	A	478	THR	2.9
1	A	89	ILE	2.8
1	A	354	ARG	2.8
1	B	254	MET	2.7
1	A	49	GLN	2.7
1	A	102	PRO	2.7
1	B	49	GLN	2.6
1	A	224	LEU	2.6
1	A	321	GLU	2.6
1	B	475	GLN	2.5
1	A	470	VAL	2.5
1	B	492	LEU	2.5
1	A	154	LYS	2.5
1	B	354	ARG	2.4
1	A	495	LEU	2.4
1	B	493	LEU	2.4
1	A	486	LEU	2.4
1	A	78	GLU	2.4
1	A	483	GLU	2.4
1	A	155	LEU	2.3
1	A	243	GLU	2.3
1	A	484	ARG	2.3
1	A	93	LYS	2.2
1	A	112	TYR	2.2
1	A	459	GLU	2.2
1	A	157	TRP	2.2
1	A	226	GLY	2.2
1	B	110	ILE	2.2
1	A	251	ASN	2.2
1	A	29	ASN	2.2
1	A	482	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	227	ASP	2.2
1	B	4	LYS	2.1
1	B	157	TRP	2.1
1	A	471	ASP	2.1
1	B	479	THR	2.1
1	A	412	ARG	2.1
1	B	303	GLU	2.0
1	A	302	LYS	2.0
1	B	78	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	RMA	A	601	14/14	0.11	1.93	37,43,44,44	0
2	RMA	B	601	14/14	0.10	1.01	38,42,43,44	0
2	FAD	B	600	53/53	0.08	-0.13	28,31,33,33	0
2	FAD	A	600	53/53	0.07	-0.40	29,31,33,33	0

6.5 Other polymers ⓘ

There are no such residues in this entry.