



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 12, 2014 – 10:45 AM EDT

PDB ID : 3S48  
Title : Human Alpha-Haemoglobin Complexed with the First NEAT Domain of IsdH from *Staphylococcus aureus*  
Authors : Kumar, K.K.; Jacques, D.A.; Caradoc-Davies, T.T.; Guss, J.M.; Gell, D.A.  
Deposited on : 2011-05-19  
Resolution : 3.05 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1439  
EDS : stable23489  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable23489

# 1 Overall quality at a glance





The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2079 (3.12-3.00)
Clashscore	79885	2629 (3.12-3.00)
Ramachandran outliers	78287	2536 (3.12-3.00)
Sidechain outliers	78261	2539 (3.12-3.00)
RSRZ outliers	66119	2081 (3.12-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	164	
1	B	164	
2	C	141	
2	D	141	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4486 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Iron-regulated surface determinant protein H.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	143	Total	C	N	O	0	0	0
			1171	748	188	235			
1	B	141	Total	C	N	O	0	0	0
			1145	733	186	226			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	66	GLY	-	EXPRESSION TAG	UNP Q6G8J7
A	67	SER	-	EXPRESSION TAG	UNP Q6G8J7
A	68	SER	-	EXPRESSION TAG	UNP Q6G8J7
A	69	HIS	-	EXPRESSION TAG	UNP Q6G8J7
A	70	HIS	-	EXPRESSION TAG	UNP Q6G8J7
A	71	HIS	-	EXPRESSION TAG	UNP Q6G8J7
A	72	HIS	-	EXPRESSION TAG	UNP Q6G8J7
A	73	HIS	-	EXPRESSION TAG	UNP Q6G8J7
A	74	HIS	-	EXPRESSION TAG	UNP Q6G8J7
A	75	SER	-	EXPRESSION TAG	UNP Q6G8J7
A	76	SER	-	EXPRESSION TAG	UNP Q6G8J7
A	77	GLY	-	EXPRESSION TAG	UNP Q6G8J7
A	78	LEU	-	EXPRESSION TAG	UNP Q6G8J7
A	79	VAL	-	EXPRESSION TAG	UNP Q6G8J7
A	80	PRO	-	EXPRESSION TAG	UNP Q6G8J7
A	81	ARG	-	EXPRESSION TAG	UNP Q6G8J7
A	82	GLY	-	EXPRESSION TAG	UNP Q6G8J7
A	83	SER	-	EXPRESSION TAG	UNP Q6G8J7
A	84	HIS	-	EXPRESSION TAG	UNP Q6G8J7
A	85	MET	-	EXPRESSION TAG	UNP Q6G8J7
B	66	GLY	-	EXPRESSION TAG	UNP Q6G8J7
B	67	SER	-	EXPRESSION TAG	UNP Q6G8J7
B	68	SER	-	EXPRESSION TAG	UNP Q6G8J7
B	69	HIS	-	EXPRESSION TAG	UNP Q6G8J7
B	70	HIS	-	EXPRESSION TAG	UNP Q6G8J7

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	71	HIS	-	EXPRESSION TAG	UNP Q6G8J7
B	72	HIS	-	EXPRESSION TAG	UNP Q6G8J7
B	73	HIS	-	EXPRESSION TAG	UNP Q6G8J7
B	74	HIS	-	EXPRESSION TAG	UNP Q6G8J7
B	75	SER	-	EXPRESSION TAG	UNP Q6G8J7
B	76	SER	-	EXPRESSION TAG	UNP Q6G8J7
B	77	GLY	-	EXPRESSION TAG	UNP Q6G8J7
B	78	LEU	-	EXPRESSION TAG	UNP Q6G8J7
B	79	VAL	-	EXPRESSION TAG	UNP Q6G8J7
B	80	PRO	-	EXPRESSION TAG	UNP Q6G8J7
B	81	ARG	-	EXPRESSION TAG	UNP Q6G8J7
B	82	GLY	-	EXPRESSION TAG	UNP Q6G8J7
B	83	SER	-	EXPRESSION TAG	UNP Q6G8J7
B	84	HIS	-	EXPRESSION TAG	UNP Q6G8J7
B	85	MET	-	EXPRESSION TAG	UNP Q6G8J7

- Molecule 2 is a protein called Hemoglobin subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	140	Total	C	N	O	S	0	0	0
			1050	674	183	190	3			
2	C	138	Total	C	N	O	S	0	0	0
			1032	661	181	187	3			

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total 1	O 1	0	0
4	C	1	Total 1	O 1	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Iron-regulated surface determinant protein H

Chain A: 



- Molecule 1: Iron-regulated surface determinant protein H

Chain B: 



- Molecule 2: Hemoglobin subunit alpha

Chain D: 



- Molecule 2: Hemoglobin subunit alpha

Chain C: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	115.61Å 115.61Å 142.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	89.74 – 3.05 36.56 – 3.05	Depositor EDS
% Data completeness (in resolution range)	99.5 (89.74-3.05) 99.5 (36.56-3.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.26 (at 3.06Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.240 , 0.268 0.237 , 0.263	Depositor DCC
$R_{free}$ test set	904 reflections (5.37%)	DCC
Wilson B-factor (Å <sup>2</sup> )	101.5	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 50.7	EDS
Estimated twinning fraction	0.029 for -h,k,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 17737 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4486	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	134.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.26	0/1200	0.40	0/1631
1	B	0.26	0/1174	0.37	0/1597
2	C	0.26	0/1059	0.34	0/1441
2	D	0.26	0/1078	0.35	0/1468
All	All	0.26	0/4511	0.37	0/6137

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1171	0	1121	15	0
1	B	1145	0	1093	5	0
2	C	1032	0	1028	9	0
2	D	1050	0	1047	14	0
3	C	43	0	30	0	0
3	D	43	0	30	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4486	0	4349	39	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (39) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:131:ILE:HD12	1:A:150:ILE:HD12	1.52	0.92
1:A:155:THR:HG21	2:D:8:THR:HG21	1.74	0.68
1:A:178:VAL:HG13	1:A:179:PRO:HA	1.79	0.65
1:A:131:ILE:HD12	1:A:150:ILE:CD1	2.27	0.63
1:A:131:ILE:CD1	1:A:150:ILE:HD12	2.25	0.63
2:C:69:ALA:HB1	2:C:76:MET:HE3	1.83	0.60
1:A:153:ALA:HB2	1:A:183:ALA:HB2	1.86	0.56
1:A:108:PRO:HG3	1:A:229:LEU:HD22	1.92	0.52
1:A:94:ILE:HG22	1:A:172:LEU:HD11	1.92	0.52
2:D:17:VAL:HG13	2:D:24:TYR:CE2	2.47	0.50
1:A:99:LEU:N	1:A:99:LEU:HD12	2.27	0.50
2:D:123:ALA:HB2	2:C:35:SER:HA	1.93	0.50
2:C:69:ALA:HB1	2:C:76:MET:CE	2.43	0.49
1:A:153:ALA:CB	1:A:183:ALA:HB2	2.43	0.48
2:D:36:PHE:CZ	2:D:100:LEU:HD22	2.47	0.48
1:A:148:LEU:HB3	1:A:150:ILE:HD11	1.95	0.47
1:B:178:VAL:HG13	1:B:179:PRO:HA	1.97	0.46
2:C:7:LYS:HG2	2:C:73:VAL:HG11	1.98	0.46
2:D:76:MET:N	2:D:77:PRO:HD2	2.31	0.46
1:A:155:THR:HG21	2:D:8:THR:CG2	2.41	0.46
2:D:129:LEU:HA	2:D:132:VAL:HG12	1.98	0.46
2:C:36:PHE:CE1	2:C:100:LEU:HD22	2.51	0.45
1:A:152:THR:CG2	2:D:11:LYS:HZ1	2.28	0.45
1:B:163:GLU:HB2	1:B:168:LEU:HD11	1.99	0.45
2:C:25:GLY:O	2:C:29:LEU:HD23	2.16	0.44
2:D:32:MET:SD	2:D:39:THR:HG21	2.58	0.44
1:A:195:GLU:C	1:A:196:LEU:HD12	2.38	0.43
2:C:97:ASN:N	2:C:97:ASN:HD22	2.17	0.43
1:A:116:LEU:HD13	1:A:127:HIS:HB3	2.00	0.43
2:D:129:LEU:O	2:D:132:VAL:HG12	2.19	0.43
2:D:6:ASP:O	2:D:10:VAL:HG23	2.19	0.42
1:B:217:LEU:HD22	1:B:219:PHE:CE2	2.55	0.42
1:B:144:ALA:HB2	1:B:225:ASN:ND2	2.35	0.41
1:B:173:VAL:HG11	1:B:186:ARG:HD2	2.02	0.41
2:D:1:VAL:HG22	2:D:127:LYS:HE2	2.02	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:32:MET:CE	2:D:39:THR:HG21	2.50	0.41
2:D:80:LEU:HD12	2:D:80:LEU:N	2.36	0.41
2:C:131:SER:O	2:C:135:VAL:HG23	2.21	0.41
2:C:76:MET:N	2:C:77:PRO:CD	2.83	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/164 (86%)	130 (92%)	10 (7%)	1 (1%)	30	77
1	B	139/164 (85%)	129 (93%)	10 (7%)	0	100	100
2	C	136/141 (96%)	130 (96%)	6 (4%)	0	100	100
2	D	138/141 (98%)	132 (96%)	6 (4%)	0	100	100
All	All	554/610 (91%)	521 (94%)	32 (6%)	1 (0%)	56	91

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	205	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/150 (87%)	131 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	126/150 (84%)	126 (100%)	0	100	100
2	C	108/113 (96%)	107 (99%)	1 (1%)	87	97
2	D	110/113 (97%)	110 (100%)	0	100	100
All	All	475/526 (90%)	474 (100%)	1 (0%)	96	99

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	97	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	D	9	ASN
1	B	120	ASN
1	B	225	ASN
2	C	97	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	HEM	C	500	2,4	42,50,50	3.74	13 (30%)	27,82,82	1.19	2 (7%)
3	HEM	D	500	2,4	42,50,50	3.74	13 (30%)	27,82,82	1.17	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	C	500	2,4	-	0/14/114/114	0/0/8/8
3	HEM	D	500	2,4	-	0/14/114/114	0/0/8/8

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	500	HEM	C3C-C2C	-14.79	1.33	1.45
3	D	500	HEM	C3C-C2C	-14.75	1.33	1.45
3	D	500	HEM	C3B-C2B	-12.16	1.33	1.45
3	C	500	HEM	C3B-C2B	-11.92	1.34	1.45
3	C	500	HEM	CMC-C2C	5.74	1.54	1.45
3	C	500	HEM	CMB-C2B	5.72	1.54	1.45
3	D	500	HEM	CMD-C2D	5.71	1.54	1.45
3	C	500	HEM	CMD-C2D	5.69	1.54	1.45
3	D	500	HEM	CMC-C2C	5.68	1.54	1.45
3	D	500	HEM	CMB-C2B	5.58	1.54	1.45
3	C	500	HEM	C3B-CAB	4.45	1.54	1.40
3	C	500	HEM	C3C-CAC	4.43	1.54	1.40
3	D	500	HEM	C3C-CAC	4.41	1.54	1.40
3	D	500	HEM	C3B-CAB	4.40	1.54	1.40
3	D	500	HEM	C3D-C2D	3.89	1.53	1.43
3	C	500	HEM	C3D-C2D	3.87	1.53	1.43
3	C	500	HEM	FE-ND	3.43	2.09	1.95
3	D	500	HEM	FE-ND	3.40	2.09	1.95
3	C	500	HEM	FE-NB	2.95	2.06	1.95
3	D	500	HEM	FE-NB	2.87	2.05	1.95
3	C	500	HEM	FE-NC	2.51	2.05	1.95
3	D	500	HEM	FE-NC	2.49	2.05	1.95
3	D	500	HEM	C4D-ND	2.49	1.40	1.33

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	500	HEM	C1D-ND	2.45	1.39	1.33
3	C	500	HEM	C4D-ND	2.44	1.39	1.33
3	C	500	HEM	C1D-ND	2.44	1.39	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	500	HEM	C3A-C4A-NA	-2.39	107.90	109.50
3	D	500	HEM	CBD-CAD-C3D	-2.24	109.66	114.51
3	C	500	HEM	CBD-CAD-C3D	-2.23	109.68	114.51
3	D	500	HEM	C3A-C4A-NA	-2.14	108.07	109.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	143/164 (87%)	-0.12	0 100 100	64, 78, 106, 116	0
1	B	141/164 (85%)	0.94	21 (14%) 3 1	189, 249, 298, 323	0
2	C	138/141 (97%)	-0.04	0 100 100	82, 114, 155, 173	0
2	D	140/141 (99%)	-0.07	0 100 100	67, 89, 134, 148	0
All	All	562/610 (92%)	0.18	21 (3%) 39 7	64, 102, 277, 323	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	137	VAL	4.6
1	B	138	TYR	4.3
1	B	87	ASP	4.1
1	B	194	GLN	3.4
1	B	195	GLU	3.2
1	B	197	LYS	3.1
1	B	136	ASP	3.0
1	B	216	LYS	3.0
1	B	106	ILE	3.0
1	B	221	LYS	2.7
1	B	161	VAL	2.5
1	B	116	LEU	2.5
1	B	105	ASP	2.4
1	B	109	ARG	2.4
1	B	157	LYS	2.4
1	B	162	TYR	2.3
1	B	193	THR	2.2
1	B	95	LYS	2.2
1	B	220	ALA	2.1
1	B	160	GLU	2.1
1	B	101	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	HEM	C	500	43/43	0.31	1.03	81,92,112,123	0
3	HEM	D	500	43/43	0.27	-0.03	69,82,99,102	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.