



Full wwPDB X-ray Structure Validation Report

Mar 1, 2014 – 01:28 AM GMT

PDB ID : 3SPT
Title : Crystal structure of GlmU from Mycobacterium tuberculosis in complex with ACETYL COENZYME A and URIDINE-DIPHOSPHATE-N-ACETYLGLUCOSAMINE
Authors : Jagtap, P.A.; Prakash, B.
Deposited on : 2011-07-03
Resolution : 2.33 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4049 (2.38-2.30)
Clashscore	79885	1094 (2.36-2.32)
Ramachandran outliers	78287	1080 (2.36-2.32)
Sidechain outliers	78261	1081 (2.36-2.32)
RSRZ outliers	66119	4050 (2.38-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	501	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3776 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional protein glmU.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	474	Total	C	N	O	S	0	0	0
			3471	2145	635	685	6			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	EXPRESSION TAG	UNP P96382
A	-4	HIS	-	EXPRESSION TAG	UNP P96382
A	-3	HIS	-	EXPRESSION TAG	UNP P96382
A	-2	HIS	-	EXPRESSION TAG	UNP P96382
A	-1	HIS	-	EXPRESSION TAG	UNP P96382
A	0	HIS	-	EXPRESSION TAG	UNP P96382

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Mg	0	0
			3	3		

- Molecule 3 is ACETYL COENZYME *A (three-letter code: ACO) (formula: C₂₃H₃₈N₇O₁₇P₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	51	23	7	17	3	0	0

- Molecule 4 is URIDINE-DIPHOSPHATE-N-ACETYLGLUCOSAMINE (three-letter code: UD1) (formula: $C_{17}H_{27}N_3O_{17}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	39	17	3	17	2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	212	Total 212	O 212	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional protein glmU

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	110.19Å 110.19Å 361.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.68 – 2.33 19.68 – 2.33	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.68-2.33) 99.8 (19.68-2.33)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 2.33Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.192 , 0.224 0.195 , 0.225	Depositor DCC
R_{free} test set	1001 reflections (2.82%)	DCC
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.220	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 17.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	5 of 36444 reflections (0.014%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3776	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, UD1, ACO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.12	6/3525 (0.2%)	1.07	16/4816 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	246	GLU	CG-CD	7.66	1.63	1.51
1	A	376	THR	CB-CG2	-7.59	1.27	1.52
1	A	342	TYR	CD1-CE1	5.26	1.47	1.39
1	A	95	CYS	CB-SG	5.15	1.91	1.82
1	A	84	ASP	CB-CG	5.09	1.62	1.51
1	A	322	VAL	CB-CG2	5.06	1.63	1.52

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	ARG	NE-CZ-NH2	-12.26	114.17	120.30
1	A	47	LEU	CB-CG-CD1	-8.91	95.85	111.00
1	A	413	ARG	NE-CZ-NH2	-8.77	115.91	120.30
1	A	76	ARG	NE-CZ-NH1	7.22	123.91	120.30
1	A	301	ARG	NE-CZ-NH1	5.85	123.23	120.30
1	A	241	ARG	NE-CZ-NH2	-5.85	117.38	120.30
1	A	84	ASP	CB-CG-OD2	5.76	123.48	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	21	ARG	CG-CD-NE	-5.74	99.75	111.80
1	A	199	LEU	CB-CG-CD1	5.55	120.43	111.00
1	A	467	GLY	N-CA-C	-5.51	99.31	113.10
1	A	413	ARG	CG-CD-NE	-5.33	100.62	111.80
1	A	21	ARG	NE-CZ-NH2	-5.27	117.66	120.30
1	A	218	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	A	51	ARG	NE-CZ-NH1	-5.14	117.73	120.30
1	A	59	ASP	CB-CG-OD1	-5.07	113.74	118.30
1	A	247	LEU	CB-CG-CD1	-5.05	102.41	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	466	PRO	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3471	0	3474	82	1
2	A	3	0	0	0	0
3	A	51	0	32	3	0
4	A	39	0	25	2	0
5	A	212	0	0	8	2
All	All	3776	0	3531	85	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

All (85) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:27:VAL:CG1	1:A:37:LEU:HB2	1.67	1.25
1:A:143:THR:HG22	1:A:228:HIS:CE1	1.81	1.15
1:A:27:VAL:HG13	1:A:37:LEU:HB2	1.08	1.07

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:143:THR:HG22	1:A:228:HIS:HE1	0.92	1.07
1:A:143:THR:CG2	1:A:228:HIS:HE1	1.70	1.02
1:A:143:THR:CG2	1:A:228:HIS:CE1	2.48	0.93
1:A:27:VAL:HG13	1:A:37:LEU:CB	2.00	0.90
1:A:83:GLN:HE21	1:A:91:HIS:HD2	1.22	0.87
1:A:83:GLN:HE21	1:A:91:HIS:CD2	1.94	0.85
1:A:305:GLY:HA3	1:A:322:VAL:HA	1.66	0.76
1:A:402:SER:OG	1:A:404:ARG:NH1	2.22	0.72
1:A:161:VAL:HG21	1:A:226:ALA:HB2	1.72	0.69
1:A:31:LEU:HD23	1:A:31:LEU:N	2.08	0.69
1:A:27:VAL:HG11	1:A:37:LEU:HB2	1.74	0.67
1:A:39:HIS:HD2	5:A:539:HOH:O	1.77	0.67
1:A:27:VAL:CG1	1:A:37:LEU:CB	2.60	0.67
1:A:374:HIS:O	1:A:376:THR:CG2	2.42	0.67
1:A:284:THR:HG21	1:A:299:GLY:O	1.95	0.67
1:A:160:GLU:CD	5:A:702:HOH:O	2.34	0.66
1:A:83:GLN:NE2	1:A:91:HIS:HD2	1.93	0.66
1:A:115:THR:HG22	1:A:118:LEU:HB2	1.78	0.65
1:A:374:HIS:O	1:A:376:THR:HG22	1.96	0.65
1:A:7:THR:HG22	1:A:107:ASN:HB2	1.78	0.64
1:A:399:ASP:OD1	1:A:400:GLY:O	2.16	0.63
1:A:375:LEU:O	1:A:392:SER:HA	1.99	0.61
1:A:418:THR:CG2	1:A:432:THR:HG21	2.32	0.59
1:A:46:LYS:HE3	5:A:700:HOH:O	2.02	0.58
1:A:432:THR:HG22	1:A:432:THR:O	2.02	0.58
1:A:31:LEU:HD23	1:A:31:LEU:H	1.67	0.58
1:A:418:THR:OG1	1:A:432:THR:CG2	2.52	0.58
1:A:117:LEU:HD11	1:A:251:LEU:HB2	1.87	0.57
1:A:415:GLY:N	1:A:432:THR:O	2.35	0.57
3:A:496:ACO:H31	5:A:592:HOH:O	2.04	0.56
1:A:418:THR:OG1	1:A:432:THR:HG21	2.06	0.55
1:A:161:VAL:HG21	1:A:226:ALA:CB	2.36	0.55
1:A:286:ILE:HD12	1:A:304:VAL:HG21	1.89	0.54
1:A:305:GLY:HA3	1:A:322:VAL:HG22	1.90	0.54
1:A:160:GLU:OE2	5:A:702:HOH:O	2.19	0.54
1:A:41:LEU:HD12	1:A:70:LEU:HD13	1.89	0.54
1:A:39:HIS:CD2	5:A:539:HOH:O	2.57	0.53
1:A:115:THR:O	1:A:115:THR:HG22	2.08	0.53
1:A:138:THR:HA	1:A:225:HIS:O	2.10	0.52
1:A:405:ARG:O	1:A:423:PRO:HA	2.10	0.52
1:A:281:GLY:O	1:A:284:THR:HG23	2.10	0.52
1:A:338:GLY:HA2	1:A:341:THR:OG1	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:10:LEU:HD13	1:A:96:GLY:HA3	1.93	0.51
1:A:418:THR:HG21	1:A:432:THR:HG21	1.92	0.51
1:A:305:GLY:CA	1:A:322:VAL:HG22	2.42	0.50
1:A:31:LEU:O	1:A:39:HIS:HE1	1.93	0.50
1:A:31:LEU:N	1:A:31:LEU:CD2	2.75	0.50
1:A:284:THR:CG2	1:A:299:GLY:O	2.59	0.49
1:A:339:PRO:O	1:A:357:ALA:HA	2.12	0.49
1:A:304:VAL:O	1:A:305:GLY:C	2.51	0.49
1:A:304:VAL:O	1:A:305:GLY:O	2.30	0.48
1:A:6:ASP:N	5:A:594:HOH:O	2.47	0.48
1:A:143:THR:HG23	1:A:228:HIS:CE1	2.45	0.48
1:A:374:HIS:O	1:A:376:THR:HG23	2.14	0.48
1:A:76:ARG:HD3	1:A:77:THR:O	2.13	0.48
1:A:34:ARG:HB2	1:A:39:HIS:CE1	2.49	0.47
1:A:27:VAL:HG11	1:A:37:LEU:HD22	1.97	0.47
3:A:496:ACO:O1A	3:A:496:ACO:O5A	2.31	0.47
1:A:391:ALA:H	3:A:496:ACO:CH3	2.28	0.47
1:A:198:ARG:HG3	1:A:198:ARG:HH11	1.79	0.47
1:A:432:THR:CG2	1:A:432:THR:O	2.63	0.46
1:A:164:ILE:HD12	1:A:214:ILE:HG13	1.96	0.46
1:A:182:ALA:HB1	1:A:235:VAL:HB	1.97	0.46
1:A:150:TYR:HB3	1:A:180:VAL:HB	1.98	0.45
1:A:311:THR:O	1:A:327:SER:HA	2.16	0.45
1:A:462:GLN:O	1:A:466:PRO:HB3	2.17	0.45
1:A:13:ALA:HB1	1:A:27:VAL:HG21	1.99	0.44
4:A:498:UD1:O5B	4:A:498:UD1:H6	2.17	0.44
1:A:415:GLY:H	1:A:432:THR:HG22	1.83	0.44
1:A:284:THR:HG22	1:A:302:CYS:H	1.83	0.43
1:A:398:TYR:CE1	1:A:400:GLY:HA2	2.53	0.43
1:A:392:SER:O	1:A:417:ASP:HA	2.18	0.43
1:A:62:ARG:O	1:A:66:LEU:CD1	2.66	0.43
1:A:26:LYS:HE3	1:A:114:ASP:HB3	1.99	0.43
1:A:286:ILE:CD1	1:A:304:VAL:HG21	2.47	0.43
1:A:415:GLY:HA3	1:A:432:THR:O	2.19	0.42
1:A:166:GLU:CD	4:A:498:UD1:H8'3	2.40	0.42
1:A:26:LYS:O	1:A:28:LEU:N	2.52	0.42
1:A:404:ARG:HD3	1:A:423:PRO:HD3	2.02	0.42
1:A:77:THR:HG23	5:A:611:HOH:O	2.19	0.41
1:A:164:ILE:CD1	1:A:214:ILE:HG13	2.49	0.41
1:A:162:MET:SD	1:A:218:ARG:HD3	2.61	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
5:A:534:HOH:O	5:A:534:HOH:O[3_555]	1.35	0.85
1:A:417:ASP:OD2	5:A:534:HOH:O[3_555]	1.87	0.33

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	472/501 (94%)	454 (96%)	15 (3%)	3 (1%)	33 38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	305	GLY
1	A	27	VAL
1	A	400	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	370/394 (94%)	346 (94%)	24 (6%)	24 27

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LEU
1	A	46	LYS
1	A	58	HIS
1	A	59	ASP
1	A	62	ARG
1	A	76	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	77	THR
1	A	82	LEU
1	A	87	LEU
1	A	94	LEU
1	A	143	THR
1	A	199	LEU
1	A	200	SER
1	A	206	GLN
1	A	218	ARG
1	A	219	SER
1	A	247	LEU
1	A	284	THR
1	A	291	GLN
1	A	376	THR
1	A	392	SER
1	A	405	ARG
1	A	424	VAL
1	A	432	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	HIS
1	A	91	HIS
1	A	202	ASN
1	A	228	HIS
1	A	291	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ACO	A	496	-	53,53,53	0.98	3 (5%)	79,79,79	4.80	47 (59%)
4	UD1	A	498	2	41,41,41	0.84	2 (4%)	58,62,62	1.30	4 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACO	A	496	-	-	2/51/67/67	0/1/3/3
4	UD1	A	498	2	-	0/25/63/63	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	496	ACO	C8A-N7A	3.59	1.41	1.34
4	A	498	UD1	C2-N1	2.82	1.41	1.38
3	A	496	ACO	C8A-N9A	2.80	1.40	1.36
3	A	496	ACO	CBP-CAP	-2.61	1.52	1.55
4	A	498	UD1	C6-C5	2.07	1.39	1.36

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	496	ACO	C3P-N4P-C5P	20.72	165.03	122.84
3	A	496	ACO	O8A-P3B-O7A	11.77	148.90	110.44
3	A	496	ACO	O2B-C2B-C3B	11.45	144.91	111.20
3	A	496	ACO	C3P-C2P-S1P	10.11	142.20	111.04
3	A	496	ACO	O3A-P2A-O6A	9.57	146.20	103.41
3	A	496	ACO	CDP-CBP-CAP	9.34	125.01	108.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	496	ACO	C7P-C6P-C5P	-7.99	98.64	112.25
3	A	496	ACO	N3A-C2A-N1A	-7.16	122.72	128.71
3	A	496	ACO	CCP-CBP-CAP	-6.87	98.65	108.70
3	A	496	ACO	CEP-CBP-CCP	-6.75	99.03	108.76
3	A	496	ACO	CEP-CBP-CDP	-6.58	99.71	109.08
3	A	496	ACO	C6P-C7P-N8P	6.24	125.31	111.87
3	A	496	ACO	CDP-CBP-CCP	6.01	117.42	108.76
3	A	496	ACO	C2B-C1B-N9A	5.73	127.96	113.27
3	A	496	ACO	O5P-C5P-N4P	5.38	133.61	122.94
3	A	496	ACO	O4B-C4B-C5B	5.29	128.25	109.36
4	A	498	UD1	N3-C2-N1	5.17	120.28	115.97
3	A	496	ACO	O2A-P1A-O5B	5.16	134.53	108.51
3	A	496	ACO	O6A-CCP-CBP	-4.82	102.51	110.57
3	A	496	ACO	O4B-C1B-C2B	4.77	114.08	106.77
3	A	496	ACO	P2A-O3A-P1A	-4.77	117.70	131.68
3	A	496	ACO	O9P-C9P-N8P	4.65	132.49	123.05
3	A	496	ACO	C6P-C5P-N4P	-4.58	108.18	116.50
3	A	496	ACO	C5B-C4B-C3B	-4.39	98.89	114.19
3	A	496	ACO	O4B-C1B-N9A	-4.36	104.39	108.44
3	A	496	ACO	N3A-C4A-N9A	4.04	132.73	125.43
4	A	498	UD1	PB-O3A-PA	-3.81	120.52	131.68
3	A	496	ACO	O3B-P3B-O7A	-3.66	96.55	106.79
3	A	496	ACO	C2B-C3B-C4B	3.66	110.29	103.16
3	A	496	ACO	O9A-P3B-O3B	-3.55	96.86	107.09
3	A	496	ACO	CBP-CAP-C9P	3.46	116.09	112.73
3	A	496	ACO	CEP-CBP-CAP	3.43	114.76	108.82
4	A	498	UD1	C3B-C2B-C1B	3.38	106.20	100.91
3	A	496	ACO	P1A-O5B-C5B	-3.30	98.32	122.03
3	A	496	ACO	O3A-P1A-O5B	3.13	117.40	103.41
4	A	498	UD1	C2-N1-C1B	2.93	120.05	118.21
3	A	496	ACO	O9A-P3B-O7A	-2.81	101.26	110.44
3	A	496	ACO	O5B-P1A-O1A	-2.78	98.49	109.37
3	A	496	ACO	CAP-C9P-N8P	-2.77	110.65	116.57
3	A	496	ACO	C7P-N8P-C9P	2.76	128.24	122.57
3	A	496	ACO	O2B-C2B-C1B	-2.73	102.96	111.23
3	A	496	ACO	C5A-C4A-N3A	-2.71	119.80	125.70
3	A	496	ACO	C8A-N9A-C4A	2.65	108.92	106.90
3	A	496	ACO	C4A-C5A-N7A	-2.55	107.33	109.52
3	A	496	ACO	O6A-P2A-O4A	-2.48	99.68	109.37
3	A	496	ACO	OAP-CAP-C9P	2.46	117.10	110.76
3	A	496	ACO	P2A-O6A-CCP	-2.36	108.36	121.66
3	A	496	ACO	N7A-C8A-N9A	-2.31	107.83	114.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	496	ACO	O3B-C3B-C2B	-2.20	102.91	111.54
3	A	496	ACO	O3A-P2A-O4A	-2.03	96.71	111.28
3	A	496	ACO	N6A-C6A-N1A	2.01	123.32	119.36

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	496	ACO	CH3-C-S1P-C2P
3	A	496	ACO	O-C-S1P-C2P

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	474/501 (94%)	-0.42	6 (1%) 74 83	15, 28, 49, 72	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	305	GLY	5.1
1	A	432	THR	3.6
1	A	478	SER	3.3
1	A	401	THR	2.3
1	A	132	ALA	2.1
1	A	304	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MG	A	500	1/1	0.10	1.36	23,23,23,23	1
4	UD1	A	498	39/39	0.13	1.11	20,20,20,20	0
3	ACO	A	496	51/51	0.10	0.03	21,33,48,57	0
2	MG	A	499	1/1	0.08	-0.07	2,2,2,2	1
2	MG	A	497	1/1	0.04	-2.50	34,34,34,34	0

6.5 Other polymers

There are no such residues in this entry.