



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 07:05 AM GMT

PDB ID : 3T2G
Title : Fructose-1,6-bisphosphatealdolase/phosphatase from Thermoproteus neutrophilus, Y229F variant with DHAP
Authors : Du, J.; Say, R.; Lue, W.; Fuchs, G.; Einsle, O.
Deposited on : 2011-07-22
Resolution : 3.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	407	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3032 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructose-1,6-bisphosphatealdolase/phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	383	Total	C	N	O	S	0	0	0
			3005	1936	515	542	12			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	229	PHE	TYR	ENGINEERED MUTATION	UNP B1YAL1
A	400	LEU	-	EXPRESSION TAG	UNP B1YAL1
A	401	GLU	-	EXPRESSION TAG	UNP B1YAL1
A	402	HIS	-	EXPRESSION TAG	UNP B1YAL1
A	403	HIS	-	EXPRESSION TAG	UNP B1YAL1
A	404	HIS	-	EXPRESSION TAG	UNP B1YAL1
A	405	HIS	-	EXPRESSION TAG	UNP B1YAL1
A	406	HIS	-	EXPRESSION TAG	UNP B1YAL1
A	407	HIS	-	EXPRESSION TAG	UNP B1YAL1

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is 1,3-DIHYDROXYACETONEPHOSPHATE (three-letter code: 13P) (formula: C₃H₇O₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			10	3	6	1		
3	A	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	O	0	0
			5	5		

4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	112.43Å 112.43Å 151.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.72 – 3.00 47.72 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.72-3.00) 99.9 (47.72-3.00)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.175 , 0.234 0.175 , 0.234	Depositor DCC
R_{free} test set	505 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	52.1	Xtriage
Anisotropy	0.214	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 17.3	EDS
Estimated twinning fraction	0.012 for -1/2*h+1/2*k-1/2*l, 1/2*h-1/2*k-1/2*l, -h-k 0.032 for -1/2*h-1/2*k+1/2*l, -1/2*h-1/2*k-1/2*l, h-k	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 10089 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3032	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 13P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/3083	0.75	1/4173 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	208	ARG	NE-CZ-NH2	-5.47	117.57	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3005	0	3003	78	0
2	A	2	0	0	0	0
3	A	20	0	10	5	0
4	A	5	0	0	0	0
All	All	3032	0	3013	83	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 14.

All (83) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:72:LEU:O	1:A:76:THR:HG22	1.37	1.18
1:A:141:LEU:HD11	1:A:145:LYS:HE3	1.54	0.89
3:A:410:13P:H32	3:A:411:13P:H31	1.55	0.88
1:A:293:MET:CE	1:A:301:LYS:HD2	2.05	0.87
1:A:72:LEU:O	1:A:76:THR:CG2	2.27	0.78
1:A:379:ASP:HB3	1:A:383:TYR:CE2	2.19	0.77
1:A:105:ASN:OD1	1:A:107:ARG:HG3	1.86	0.75
1:A:107:ARG:HH12	1:A:221:ARG:NH2	1.86	0.73
1:A:133:LYS:HG2	1:A:297:ASP:O	1.89	0.72
1:A:366:ALA:O	1:A:367:LYS:HB2	1.90	0.72
1:A:107:ARG:HH12	1:A:221:ARG:HH22	1.35	0.70
1:A:268:SER:HA	1:A:351:LEU:HD12	1.73	0.69
3:A:410:13P:C3	3:A:411:13P:H31	2.22	0.69
1:A:293:MET:HE1	1:A:301:LYS:HD2	1.75	0.69
1:A:293:MET:HE2	1:A:301:LYS:HD2	1.75	0.68
3:A:410:13P:H32	3:A:411:13P:C3	2.25	0.66
1:A:365:LEU:HA	1:A:367:LYS:O	1.96	0.66
1:A:356:MET:O	1:A:357:GLU:HB2	1.97	0.65
1:A:265:MET:SD	1:A:266:ARG:HG3	2.36	0.65
1:A:265:MET:HB2	1:A:296:PHE:O	1.99	0.63
1:A:293:MET:HE1	1:A:301:LYS:HB2	1.80	0.63
1:A:379:ASP:HB3	1:A:383:TYR:CD2	2.37	0.60
3:A:410:13P:C3	3:A:411:13P:C3	2.79	0.60
1:A:354:GLU:N	1:A:356:MET:O	2.36	0.58
1:A:293:MET:CE	1:A:301:LYS:CD	2.81	0.58
1:A:136:PRO:HA	1:A:235:PRO:HG3	1.85	0.57
1:A:126:ILE:HG22	1:A:240:ARG:HG2	1.87	0.57
1:A:60:HIS:CD2	1:A:69:ILE:HG13	2.40	0.56
1:A:47:VAL:O	1:A:287:ILE:HA	2.07	0.55
1:A:293:MET:HE1	1:A:301:LYS:CD	2.37	0.54
1:A:27:GLU:HG2	1:A:285:ARG:HH22	1.73	0.54
1:A:272:PRO:HG3	1:A:336:ILE:HB	1.91	0.53
1:A:49:ASN:ND2	1:A:290:GLY:H	2.07	0.53
1:A:366:ALA:O	1:A:367:LYS:CB	2.58	0.52
1:A:171:LEU:HD23	1:A:219:VAL:HG23	1.92	0.51
1:A:108:GLY:C	1:A:230:VAL:HG21	2.31	0.51
1:A:293:MET:HE2	1:A:301:LYS:CD	2.41	0.50
1:A:368:VAL:HG12	1:A:369:LYS:N	2.26	0.50
1:A:129:PHE:HB2	1:A:237:LEU:HB3	1.93	0.50
1:A:371:TYR:CD2	1:A:371:TYR:N	2.80	0.49
1:A:298:GLY:HA3	1:A:299:PRO:C	2.33	0.49
1:A:365:LEU:CA	1:A:367:LYS:O	2.61	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:4:THR:OG1	1:A:117:GLU:HG3	2.13	0.48
1:A:5:VAL:HG12	1:A:238:LEU:HD11	1.94	0.48
1:A:175:GLU:HB3	1:A:177:LYS:HE3	1.96	0.47
1:A:22:HIS:HB3	1:A:25:MET:HG3	1.95	0.47
1:A:194:ILE:HD13	1:A:202:ILE:HD11	1.96	0.47
1:A:181:LEU:HD13	1:A:190:LEU:HA	1.95	0.47
1:A:63:GLY:HA2	1:A:117:GLU:HG3	1.95	0.47
1:A:331:GLN:O	1:A:335:GLN:HG3	2.15	0.47
1:A:136:PRO:HG2	1:A:232:LYS:HB2	1.96	0.47
1:A:92:GLY:O	1:A:95:GLN:HB2	2.15	0.46
1:A:221:ARG:NE	1:A:221:ARG:H	2.14	0.45
1:A:138:ALA:HB1	1:A:261:VAL:HG11	1.99	0.45
1:A:364:ILE:HG13	1:A:364:ILE:H	1.48	0.45
1:A:230:VAL:HG12	1:A:231:GLY:N	2.31	0.45
1:A:298:GLY:CA	1:A:299:PRO:O	2.65	0.44
1:A:365:LEU:C	1:A:367:LYS:N	2.71	0.44
1:A:201:ILE:HG22	1:A:219:VAL:HG11	1.99	0.44
1:A:176:HIS:N	1:A:176:HIS:ND1	2.66	0.44
1:A:356:MET:O	1:A:357:GLU:CB	2.64	0.44
1:A:382:LYS:HA	1:A:385:GLU:HB2	1.98	0.44
1:A:27:GLU:CG	1:A:285:ARG:HH22	2.30	0.44
3:A:410:13P:O3	3:A:411:13P:H32	2.18	0.43
1:A:298:GLY:CA	1:A:299:PRO:C	2.87	0.43
1:A:379:ASP:HB3	1:A:383:TYR:HE2	1.79	0.43
1:A:32:LYS:HG2	1:A:76:THR:HB	2.01	0.43
1:A:185:GLU:OE2	1:A:208:ARG:NH2	2.37	0.43
1:A:141:LEU:HD12	1:A:141:LEU:O	2.19	0.42
1:A:203:ARG:HH12	1:A:221:ARG:HH12	1.68	0.42
1:A:27:GLU:HG2	1:A:285:ARG:NH2	2.35	0.42
1:A:149:ASP:HA	1:A:150:PRO:HD3	1.74	0.41
1:A:136:PRO:CG	1:A:232:LYS:HB2	2.50	0.41
1:A:384:ILE:O	1:A:388:VAL:HG23	2.19	0.41
1:A:319:ASP:C	1:A:319:ASP:OD1	2.58	0.41
1:A:316:GLY:HA3	1:A:317:GLY:HA2	1.81	0.41
1:A:379:ASP:O	1:A:380:ARG:C	2.59	0.41
1:A:182:LYS:HD2	1:A:208:ARG:HH21	1.86	0.41
1:A:265:MET:HB3	1:A:296:PHE:CE2	2.56	0.40
1:A:18:HIS:O	1:A:133:LYS:NZ	2.47	0.40
1:A:199:ARG:HD3	1:A:200:TYR:CE2	2.57	0.40
1:A:5:VAL:HG11	1:A:238:LEU:HD21	2.03	0.40
1:A:101:ALA:O	1:A:102:PHE:CB	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/407 (93%)	332 (88%)	34 (9%)	13 (3%)	6	31

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	101	ALA
1	A	102	PHE
1	A	103	SER
1	A	354	GLU
1	A	365	LEU
1	A	367	LYS
1	A	232	LYS
1	A	357	GLU
1	A	363	LYS
1	A	386	ALA
1	A	359	THR
1	A	104	GLY
1	A	234	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/331 (94%)	282 (91%)	29 (9%)	13	45

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	76	THR
1	A	78	LYS
1	A	95	GLN
1	A	99	LYS
1	A	100	ASP
1	A	113	VAL
1	A	115	GLU
1	A	117	GLU
1	A	127	ILE
1	A	161	SER
1	A	176	HIS
1	A	182	LYS
1	A	219	VAL
1	A	221	ARG
1	A	232	LYS
1	A	233	ASP
1	A	234	ASP
1	A	238	LEU
1	A	243	SER
1	A	284	GLU
1	A	286	ARG
1	A	331	GLN
1	A	341	ARG
1	A	355	GLU
1	A	367	LYS
1	A	369	LYS
1	A	382	LYS
1	A	389	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	49	ASN
1	A	95	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	13P	A	410	2	9,9,9	0.58	0	12,12,12	0.98	0
3	13P	A	411	-	9,9,9	0.53	0	12,12,12	2.09	2 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	13P	A	410	2	-	0/8/8/8	0/0/0/0
3	13P	A	411	-	-	1/8/8/8	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	411	13P	O1-C1-C2	-5.85	101.22	110.53
3	A	411	13P	O3P-P-O2P	2.12	115.88	107.61

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	411	13P	O2-C2-C3-O3

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/407 (94%)	-0.38	0 100 100	25, 37, 69, 78	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	13P	A	410	10/10	0.16	1.37	64,70,74,74	1
3	13P	A	411	10/10	0.17	1.02	37,38,40,41	10
2	MG	A	409	1/1	0.06	-3.58	50,50,50,50	0
2	MG	A	408	1/1	0.04	-7.81	13,13,13,13	0

6.5 Other polymers ⓘ

There are no such residues in this entry.