



# wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 10:52 PM GMT

PDB ID : 3TBE  
Title : The crystal structure of the complex of Streptococcus agalactiae sortase C1 and MTSET  
Authors : Khare, B.  
Deposited on : 2011-08-05  
Resolution : 2.85 Å (reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1524 (2.90-2.82)
Clashscore	79885	1879 (2.90-2.82)
Ramachandran outliers	78287	1824 (2.90-2.82)
Sidechain outliers	78261	1827 (2.90-2.82)
RSRZ outliers	66119	1526 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	230	
1	B	230	
1	C	230	
1	D	230	
1	E	230	
1	F	230	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	ETM	B	422	-	X
2	ETM	C	422	-	X

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Geometry</b>	<b>Electron density</b>
2	ETM	D	422	-	X
2	ETM	E	422	-	X
2	ETM	F	422	-	X

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 9008 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sortase family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	194	1509	959	266	281	3	0	0	0
1	B	193	1478	940	261	274	3	0	0	0
1	C	189	1455	924	255	273	3	0	0	0
1	D	193	1473	936	262	272	3	0	0	0
1	E	193	1483	944	261	275	3	0	0	0
1	F	188	1434	914	253	264	3	0	0	0

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	EXPRESSION TAG	UNP Q8E0S7
A	-9	ARG	-	EXPRESSION TAG	UNP Q8E0S7
A	-8	GLY	-	EXPRESSION TAG	UNP Q8E0S7
A	-7	SER	-	EXPRESSION TAG	UNP Q8E0S7
A	-6	HIS	-	EXPRESSION TAG	UNP Q8E0S7
A	-5	HIS	-	EXPRESSION TAG	UNP Q8E0S7
A	-4	HIS	-	EXPRESSION TAG	UNP Q8E0S7
A	-3	HIS	-	EXPRESSION TAG	UNP Q8E0S7
A	-2	HIS	-	EXPRESSION TAG	UNP Q8E0S7
A	-1	HIS	-	EXPRESSION TAG	UNP Q8E0S7
A	0	GLY	-	EXPRESSION TAG	UNP Q8E0S7
A	1	SER	-	EXPRESSION TAG	UNP Q8E0S7
B	-10	MET	-	EXPRESSION TAG	UNP Q8E0S7
B	-9	ARG	-	EXPRESSION TAG	UNP Q8E0S7
B	-8	GLY	-	EXPRESSION TAG	UNP Q8E0S7
B	-7	SER	-	EXPRESSION TAG	UNP Q8E0S7
B	-6	HIS	-	EXPRESSION TAG	UNP Q8E0S7

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	HIS	-	EXPRESSION TAG	UNP Q8E0S7
B	-4	HIS	-	EXPRESSION TAG	UNP Q8E0S7
B	-3	HIS	-	EXPRESSION TAG	UNP Q8E0S7
B	-2	HIS	-	EXPRESSION TAG	UNP Q8E0S7
B	-1	HIS	-	EXPRESSION TAG	UNP Q8E0S7
B	0	GLY	-	EXPRESSION TAG	UNP Q8E0S7
B	1	SER	-	EXPRESSION TAG	UNP Q8E0S7
C	-10	MET	-	EXPRESSION TAG	UNP Q8E0S7
C	-9	ARG	-	EXPRESSION TAG	UNP Q8E0S7
C	-8	GLY	-	EXPRESSION TAG	UNP Q8E0S7
C	-7	SER	-	EXPRESSION TAG	UNP Q8E0S7
C	-6	HIS	-	EXPRESSION TAG	UNP Q8E0S7
C	-5	HIS	-	EXPRESSION TAG	UNP Q8E0S7
C	-4	HIS	-	EXPRESSION TAG	UNP Q8E0S7
C	-3	HIS	-	EXPRESSION TAG	UNP Q8E0S7
C	-2	HIS	-	EXPRESSION TAG	UNP Q8E0S7
C	-1	HIS	-	EXPRESSION TAG	UNP Q8E0S7
C	0	GLY	-	EXPRESSION TAG	UNP Q8E0S7
C	1	SER	-	EXPRESSION TAG	UNP Q8E0S7
D	-10	MET	-	EXPRESSION TAG	UNP Q8E0S7
D	-9	ARG	-	EXPRESSION TAG	UNP Q8E0S7
D	-8	GLY	-	EXPRESSION TAG	UNP Q8E0S7
D	-7	SER	-	EXPRESSION TAG	UNP Q8E0S7
D	-6	HIS	-	EXPRESSION TAG	UNP Q8E0S7
D	-5	HIS	-	EXPRESSION TAG	UNP Q8E0S7
D	-4	HIS	-	EXPRESSION TAG	UNP Q8E0S7
D	-3	HIS	-	EXPRESSION TAG	UNP Q8E0S7
D	-2	HIS	-	EXPRESSION TAG	UNP Q8E0S7
D	-1	HIS	-	EXPRESSION TAG	UNP Q8E0S7
D	0	GLY	-	EXPRESSION TAG	UNP Q8E0S7
D	1	SER	-	EXPRESSION TAG	UNP Q8E0S7
E	-10	MET	-	EXPRESSION TAG	UNP Q8E0S7
E	-9	ARG	-	EXPRESSION TAG	UNP Q8E0S7
E	-8	GLY	-	EXPRESSION TAG	UNP Q8E0S7
E	-7	SER	-	EXPRESSION TAG	UNP Q8E0S7
E	-6	HIS	-	EXPRESSION TAG	UNP Q8E0S7
E	-5	HIS	-	EXPRESSION TAG	UNP Q8E0S7
E	-4	HIS	-	EXPRESSION TAG	UNP Q8E0S7
E	-3	HIS	-	EXPRESSION TAG	UNP Q8E0S7
E	-2	HIS	-	EXPRESSION TAG	UNP Q8E0S7
E	-1	HIS	-	EXPRESSION TAG	UNP Q8E0S7
E	0	GLY	-	EXPRESSION TAG	UNP Q8E0S7

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	1	SER	-	EXPRESSION TAG	UNP Q8E0S7
F	-10	MET	-	EXPRESSION TAG	UNP Q8E0S7
F	-9	ARG	-	EXPRESSION TAG	UNP Q8E0S7
F	-8	GLY	-	EXPRESSION TAG	UNP Q8E0S7
F	-7	SER	-	EXPRESSION TAG	UNP Q8E0S7
F	-6	HIS	-	EXPRESSION TAG	UNP Q8E0S7
F	-5	HIS	-	EXPRESSION TAG	UNP Q8E0S7
F	-4	HIS	-	EXPRESSION TAG	UNP Q8E0S7
F	-3	HIS	-	EXPRESSION TAG	UNP Q8E0S7
F	-2	HIS	-	EXPRESSION TAG	UNP Q8E0S7
F	-1	HIS	-	EXPRESSION TAG	UNP Q8E0S7
F	0	GLY	-	EXPRESSION TAG	UNP Q8E0S7
F	1	SER	-	EXPRESSION TAG	UNP Q8E0S7

- Molecule 2 is 2-(TRIMETHYLAMMONIUM)ETHYLTHIOL (three-letter code: ETM) (formula: C<sub>5</sub>H<sub>14</sub>NS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	S	0	0
			7	5	1	1		
2	B	1	Total	C	N	S	0	0
			7	5	1	1		
2	C	1	Total	C	N	S	0	0
			7	5	1	1		
2	D	1	Total	C	N	S	0	0
			7	5	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	E	1	Total	C	N	S	0	0
			7	5	1	1		
2	F	1	Total	C	N	S	0	0
			7	5	1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	10	Total O 10 10	0	0
5	B	17	Total O 17 17	0	0
5	C	11	Total O 11 11	0	0
5	D	36	Total O 36 36	0	0
5	E	19	Total O 19 19	0	0
5	F	15	Total O 15 15	0	0




## Chain D:




- Molecule 1: Sortase family protein

## Chain E:



- Molecule 1: Sortase family protein

## Chain F:



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	122.41Å 70.65Å 194.44Å 90.00° 90.43° 90.00°	Depositor
Resolution (Å)	27.71 – 2.85 27.71 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.3 (27.71-2.85) 97.9 (27.71-2.85)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.16 (at 2.85Å)	Xtriage
Refinement program	CNS, REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.250 , 0.296 0.258 , 0.297	Depositor DCC
$R_{free}$ test set	1956 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	68.8	Xtriage
Anisotropy	0.338	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 9.9	EDS
Estimated twinning fraction	0.004 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.008 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.448 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.438 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.002 for -h,-k,l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 38931 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9008	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4, ETM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.79	0/1536	0.81	1/2084 (0.0%)
1	B	0.83	1/1503 (0.1%)	0.78	1/2041 (0.0%)
1	C	0.88	0/1480	0.93	6/2010 (0.3%)
1	D	0.87	0/1497	0.92	6/2032 (0.3%)
1	E	0.84	0/1509	0.89	6/2048 (0.3%)
1	F	0.77	0/1460	0.84	5/1987 (0.3%)
All	All	0.83	1/8985 (0.0%)	0.86	25/12202 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	63	TYR	CE1-CZ	-5.47	1.31	1.38

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	193	ARG	NE-CZ-NH1	-11.41	114.59	120.30
1	C	65	ARG	NE-CZ-NH1	10.37	125.48	120.30
1	E	193	ARG	NE-CZ-NH2	9.02	124.81	120.30
1	F	193	ARG	NE-CZ-NH2	-8.24	116.18	120.30
1	D	65	ARG	NE-CZ-NH2	8.22	124.41	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	123	ARG	Sidechain

## 5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1509	0	1509	61	0
1	B	1478	0	1459	56	0
1	C	1455	0	1437	57	0
1	D	1473	0	1461	54	0
1	E	1483	0	1468	43	0
1	F	1434	0	1413	26	0
2	A	7	0	13	3	0
2	B	7	0	13	0	0
2	C	7	0	14	1	0
2	D	7	0	13	0	0
2	E	7	0	13	2	0
2	F	7	0	13	0	0
3	B	10	0	0	2	0
3	C	5	0	0	0	0
3	D	5	0	0	2	0
3	F	5	0	0	0	0
4	C	1	0	0	2	0
5	A	10	0	0	0	0
5	B	17	0	0	2	0
5	C	11	0	0	0	0
5	D	36	0	0	10	0
5	E	19	0	0	0	0
5	F	15	0	0	0	0
All	All	9008	0	8826	291	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

The worst 5 of 291 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:E:129:LYS:O	1:E:132:THR:HG22	1.42	1.18
1:B:129:LYS:O	1:B:132:THR:HG22	1.46	1.15
1:A:204:VAL:HG12	1:A:205:GLU:O	1.48	1.11
1:E:123:ARG:HG3	1:E:132:THR:HA	1.36	1.06
1:D:32:ASN:OD1	5:D:241:HOH:O	1.71	1.05

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/230 (82%)	173 (92%)	13 (7%)	2 (1%)	21	57
1	B	185/230 (80%)	171 (92%)	12 (6%)	2 (1%)	21	57
1	C	183/230 (80%)	172 (94%)	10 (6%)	1 (0%)	38	77
1	D	185/230 (80%)	178 (96%)	6 (3%)	1 (0%)	38	77
1	E	187/230 (81%)	178 (95%)	8 (4%)	1 (0%)	38	77
1	F	182/230 (79%)	170 (93%)	12 (7%)	0	100	100
All	All	1110/1380 (80%)	1042 (94%)	61 (6%)	7 (1%)	33	73

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	38	ALA
1	E	5	ASN
1	A	46	ALA
1	B	46	ALA
1	A	206	LYS

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	159/193 (82%)	141 (89%)	18 (11%)	9 23
1	B	151/193 (78%)	135 (89%)	16 (11%)	10 26
1	C	150/193 (78%)	127 (85%)	23 (15%)	4 10
1	D	150/193 (78%)	132 (88%)	18 (12%)	7 19
1	E	151/193 (78%)	134 (89%)	17 (11%)	9 23
1	F	146/193 (76%)	136 (93%)	10 (7%)	22 53
All	All	907/1158 (78%)	805 (89%)	102 (11%)	9 23

5 of 102 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	80	ARG
1	D	14	THR
1	F	22	ASN
1	C	82	ASN
1	C	134	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	133	ASN
1	D	82	ASN
1	F	75	HIS
1	C	166	GLN
1	A	190	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ETM	A	422	1	6,6,6	0.60	0	8,8,8	0.50	0
3	SO4	B	220	-	4,4,4	0.54	0	6,6,6	0.26	0
3	SO4	B	221	-	4,4,4	0.13	0	6,6,6	0.40	0
2	ETM	B	422	1	6,6,6	0.72	0	8,8,8	0.58	0
3	SO4	C	221	-	4,4,4	0.13	0	6,6,6	0.20	0
2	ETM	C	422	1	6,6,6	0.64	0	8,8,8	0.45	0
3	SO4	D	220	-	4,4,4	0.40	0	6,6,6	0.17	0
2	ETM	D	422	1	6,6,6	0.76	0	8,8,8	0.57	0
2	ETM	E	422	1	6,6,6	0.62	0	8,8,8	0.42	0
3	SO4	F	220	-	4,4,4	0.41	0	6,6,6	0.41	0
2	ETM	F	422	1	6,6,6	0.51	0	8,8,8	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ETM	A	422	1	-	0/4/4/4	0/0/0/0
3	SO4	B	220	-	-	0/0/0/0	0/0/0/0
3	SO4	B	221	-	-	0/0/0/0	0/0/0/0
2	ETM	B	422	1	-	0/4/4/4	0/0/0/0
3	SO4	C	221	-	-	0/0/0/0	0/0/0/0
2	ETM	C	422	1	-	0/4/4/4	0/0/0/0
3	SO4	D	220	-	-	0/0/0/0	0/0/0/0
2	ETM	D	422	1	-	0/4/4/4	0/0/0/0
2	ETM	E	422	1	-	0/4/4/4	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	F	220	-	-	0/0/0/0	0/0/0/0
2	ETM	F	422	1	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	194/230 (84%)	0.30	5 (2%) 53 63	24, 43, 89, 96	0
1	B	193/230 (83%)	0.13	3 (1%) 68 77	25, 43, 85, 122	0
1	C	189/230 (82%)	0.17	13 (6%) 17 19	26, 45, 89, 93	0
1	D	193/230 (83%)	0.22	2 (1%) 79 86	26, 44, 88, 115	0
1	E	193/230 (83%)	0.28	10 (5%) 26 32	28, 49, 95, 106	0
1	F	188/230 (81%)	0.35	8 (4%) 34 40	29, 49, 93, 109	0
All	All	1150/1380 (83%)	0.24	41 (3%) 41 48	24, 46, 90, 122	0

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	47	LEU	5.3
1	C	124	GLY	4.8
1	E	46	ALA	4.1
1	C	126	PRO	3.8
1	C	9	PHE	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	ETM	F	422	7/7	0.79	5.66	182,184,185,185	0
2	ETM	D	422	7/7	0.43	5.18	85,85,86,88	0
2	ETM	E	422	7/7	0.40	4.40	121,123,124,124	0
2	ETM	B	422	7/7	0.37	3.61	99,99,100,101	0
2	ETM	C	422	7/7	0.36	2.75	144,146,146,146	0
2	ETM	A	422	7/7	0.30	1.22	111,115,116,116	0
3	SO4	B	220	5/5	0.18	-0.29	48,48,49,49	0
4	CL	C	220	1/1	0.21	-0.56	71,71,71,71	0
3	SO4	F	220	5/5	0.15	-0.68	55,56,57,57	0
3	SO4	D	220	5/5	0.18	-0.80	50,51,51,51	0
3	SO4	C	221	5/5	0.12	-3.00	93,94,94,95	0
3	SO4	B	221	5/5	0.18	-3.62	89,89,90,90	0

## 6.5 Other polymers (i)

There are no such residues in this entry.