



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 01:14 PM GMT

PDB ID : 1TH4
Title : crystal structure of NADPH depleted bovine liver catalase complexed with 3-amino-1,2,4-triazole
Authors : Sugadev, R.; Ponnuswamy, M.N.; Kumaran, D.; Swaminathan, S.; Sekar, K.
Deposited on : 2004-06-01
Resolution : 2.98 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1468 (3.00-2.96)
Clashscore	79885	1894 (3.00-2.96)
Ramachandran outliers	78287	1826 (3.00-2.96)
Sidechain outliers	78261	1829 (3.00-2.96)
RSRZ outliers	66119	1469 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	506	
1	B	506	
1	C	506	
1	D	506	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	HEM	A	2000	-	X
3	HEM	B	2001	-	X
3	HEM	C	2002	-	X

2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 16915 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catalase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	499	Total	C	N	O	S	0	0	0
			4017	2548	715	740	14			
1	B	499	Total	C	N	O	S	0	0	0
			4017	2548	715	740	14			
1	C	499	Total	C	N	O	S	1	0	0
			4017	2548	715	740	14			
1	D	499	Total	C	N	O	S	0	0	0
			4017	2548	715	740	14			

- Molecule 2 is 3-AMINO-1,2,4-TRIAZOLE (three-letter code: 3TR) (formula: C₂H₄N₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	C	N	0	0
			6	2	4		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	154	Total	O	0	0
			154	154		
4	B	145	Total	O	0	0
			145	145		
4	C	188	Total	O	0	0
			188	188		
4	D	182	Total	O	0	0
			182	182		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Catalase


Chain A:



• Molecule 1: Catalase


Chain B:






• Molecule 1: Catalase


Chain C:



• Molecule 1: Catalase

Chain D:





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.03Å 141.09Å 231.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.90 – 2.98 44.37 – 2.98	Depositor EDS
% Data completeness (in resolution range)	57.6 (39.90-2.98) 57.6 (44.37-2.98)	Depositor EDS
R_{merge}	0.30	Depositor
R_{sym}	0.30	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.73 (at 2.96Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.235 , 0.292 0.228 , 0.285	Depositor DCC
R_{free} test set	1001 reflections (3.10%)	DCC
Wilson B-factor (Å ²)	53.3	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 23.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 33297 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	16915	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, 3TR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	5/4137 (0.1%)	1.78	61/5619 (1.1%)
1	B	0.99	6/4137 (0.1%)	1.45	31/5619 (0.6%)
1	C	0.66	8/4137 (0.2%)	1.49	33/5619 (0.6%)
1	D	0.61	1/4137 (0.0%)	0.92	9/5619 (0.2%)
All	All	0.73	20/16548 (0.1%)	1.44	134/22476 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	2
1	B	1	3
1	C	2	2
1	D	1	0
All	All	5	7

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	319	ARG	CD-NE	49.16	2.30	1.46
1	D	413	HIS	CA-CB	-20.41	1.09	1.53
1	C	202	ARG	NE-CZ	16.91	1.55	1.33
1	B	319	ARG	NE-CZ	16.04	1.53	1.33
1	A	304	HIS	CA-CB	12.45	1.81	1.53

The worst 5 of 134 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	357	TYR	CB-CG-CD2	-68.04	80.18	121.00
1	C	202	ARG	NE-CZ-NH2	-52.67	93.97	120.30
1	B	319	ARG	NE-CZ-NH1	-46.60	97.00	120.30
1	C	202	ARG	NE-CZ-NH1	44.10	142.35	120.30
1	B	319	ARG	CG-CD-NE	-41.12	25.46	111.80

All (5) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	395	ASP	CA
1	B	395	ASP	CA
1	C	453	GLU	CA
1	C	501	GLU	CA
1	D	371	GLN	CA

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	357	TYR	Sidechain
1	A	421	ARG	Sidechain
1	B	126	ARG	Sidechain
1	B	319	ARG	Sidechain
1	B	394	MET	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4017	0	3838	473	1
1	B	4017	0	3839	431	0
1	C	4017	0	3840	429	0
1	D	4017	0	3839	389	0
2	D	6	0	3	12	0
3	A	43	0	30	79	0
3	B	43	0	30	39	0
3	C	43	0	30	49	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	43	0	30	25	0
4	A	154	0	0	30	0
4	B	145	0	0	28	0
4	C	188	0	0	17	1
4	D	182	0	0	14	0
All	All	16915	0	15479	1562	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 49.

The worst 5 of 1562 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:304:HIS:CB	1:A:304:HIS:CA	1.81	1.58
1:A:100:GLU:HG2	1:A:101:HIS:CE1	1.43	1.52
3:A:2000:HEM:CBC	3:A:2000:HEM:CAC	1.87	1.47
1:A:353:ARG:HD2	3:A:2000:HEM:CHC	1.53	1.37
1:C:111:ARG:NE	3:C:2002:HEM:O1D	1.57	1.36

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:289:GLU:OE1	4:C:2061:HOH:O[3_655]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/506 (98%)	400 (80%)	80 (16%)	17 (3%)	6	29
1	B	497/506 (98%)	408 (82%)	65 (13%)	24 (5%)	4	19
1	C	497/506 (98%)	398 (80%)	82 (16%)	17 (3%)	6	29
1	D	497/506 (98%)	402 (81%)	74 (15%)	21 (4%)	4	23
All	All	1988/2024 (98%)	1608 (81%)	301 (15%)	79 (4%)	5	24

5 of 79 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	100	GLU
1	B	19	ALA
1	B	100	GLU
1	B	331	LEU
1	B	394	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	431/437 (99%)	396 (92%)	35 (8%)	17	51
1	B	431/437 (99%)	387 (90%)	44 (10%)	11	37
1	C	431/437 (99%)	394 (91%)	37 (9%)	15	48
1	D	431/437 (99%)	403 (94%)	28 (6%)	24	64
All	All	1724/1748 (99%)	1580 (92%)	144 (8%)	16	50

5 of 144 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	374	VAL
1	C	22	LYS
1	D	378	TYR
1	B	395	ASP
1	B	438	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 74 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	429	GLN
1	C	32	ASN
1	D	371	GLN
1	B	435	ASN
1	C	3	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	HEM	A	2000	1	49,50,50	2.84	17 (34%)	46,82,82	1.37	5 (10%)
3	HEM	B	2001	1	49,50,50	2.85	18 (36%)	46,82,82	2.92	9 (19%)
3	HEM	C	2002	1	49,50,50	2.14	13 (26%)	46,82,82	1.24	4 (8%)
3	HEM	D	2003	1,2	49,50,50	3.91	19 (38%)	46,82,82	4.92	11 (23%)
2	3TR	D	3074	1,3	6,6,6	2.92	3 (50%)	6,7,7	2.32	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	2000	1	-	0/14/114/114	0/0/8/8
3	HEM	B	2001	1	-	0/14/114/114	0/0/8/8
3	HEM	C	2002	1	-	0/14/114/114	0/0/8/8
3	HEM	D	2003	1,2	-	0/14/114/114	0/0/8/8

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3TR	D	3074	1,3	-	0/0/0/0	0/1/1/1

The worst 5 of 70 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2003	HEM	CHD-C4C	16.04	1.67	1.36
3	D	2003	HEM	CHC-C4B	-12.27	1.10	1.39
3	A	2000	HEM	CBC-CAC	10.05	1.87	1.28
3	D	2003	HEM	CHD-C1D	-9.14	1.17	1.39
3	B	2001	HEM	FE-NC	8.98	2.31	1.97

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2003	HEM	CBA-CAA-C2A	28.41	162.74	112.69
3	B	2001	HEM	CHD-C1D-ND	15.05	137.09	124.58
3	D	2003	HEM	CHC-C4B-NB	-8.65	117.40	124.58
3	D	2003	HEM	C1A-CHA-C4D	-6.93	118.35	127.47
3	B	2001	HEM	C1D-CHD-C4C	-6.87	108.49	126.57

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	499/506 (98%)	-0.36	2 (0%)	90	51	10, 40, 76, 97	0
1	B	499/506 (98%)	-0.25	6 (1%)	75	32	16, 44, 78, 102	0
1	C	499/506 (98%)	-0.13	5 (1%)	79	34	16, 41, 81, 100	1 (0%)
1	D	499/506 (98%)	-0.27	5 (1%)	79	34	15, 45, 84, 101	0
All	All	1996/2024 (98%)	-0.26	18 (0%)	81	36	10, 43, 80, 102	1 (0%)

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	421	ARG	4.1
1	B	19	ALA	3.4
1	B	20	ALA	3.4
1	C	448	LYS	3.1
1	D	224	ALA	2.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	HEM	A	2000	43/43	0.41	5.71	80,94,101,101	0
3	HEM	C	2002	43/43	0.37	5.17	41,58,72,92	0
3	HEM	B	2001	43/43	0.29	4.12	21,56,83,101	0
2	3TR	D	3074	6/6	0.20	0.69	89,96,99,101	0
3	HEM	D	2003	43/43	0.17	-0.03	20,38,56,59	0

6.5 Other polymers ⓘ

There are no such residues in this entry.