



# Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 03:57 PM GMT

PDB ID : 1TLO  
Title : High resolution crystal structure of calpain I protease core in complex with E64  
Authors : Moldoveanu, T.; Campbell, R.L.; Cuerrier, D.; Davies, P.L.  
Deposited on : 2004-06-09  
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	339	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2674 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calpain 1, large [catalytic] subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	0	0
			2594	1662	436	487	9			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	MET	-	INITIATING METHIONINE	UNP P97571
A	357	LEU	-	EXPRESSION TAG	UNP P97571
A	358	GLU	-	EXPRESSION TAG	UNP P97571
A	359	HIS	-	EXPRESSION TAG	UNP P97571
A	360	HIS	-	EXPRESSION TAG	UNP P97571
A	361	HIS	-	EXPRESSION TAG	UNP P97571
A	362	HIS	-	EXPRESSION TAG	UNP P97571
A	363	HIS	-	EXPRESSION TAG	UNP P97571
A	364	HIS	-	EXPRESSION TAG	UNP P97571

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		

- Molecule 3 is N-[N-[1-HYDROXYCARBOXYETHYL-CARBONYL]LEUCYLAMINO-BUTYL]-GUANIDINE (three-letter code: E64) (formula: C<sub>15</sub>H<sub>30</sub>N<sub>5</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			25	15	5	5		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	53	Total	O	0	0
			53	53		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Calpain 1, large [catalytic] subunit

Chain A: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	40.52Å 69.76Å 108.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.90 29.47 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-1.90) 98.3 (29.47-1.90)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.62 (at 1.91Å)	Xtriage
Refinement program	CNS 1.1, XFIT	Depositor
R, $R_{free}$	0.238 , 0.272 0.239 , 0.270	Depositor DCC
$R_{free}$ test set	1193 reflections (4.85%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.4	Xtriage
Anisotropy	0.532	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 45.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 24593 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2674	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, E64

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/2661	0.58	0/3610

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2594	0	2501	49	0
2	A	2	0	0	0	0
3	A	25	0	26	8	0
4	A	53	0	0	0	0
All	All	2674	0	2527	49	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

All (49) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:248:LEU:HG	1:A:350:ILE:HD12	1.53	0.89
1:A:184:ASN:H	1:A:184:ASN:HD22	1.28	0.80
1:A:234:LEU:HD12	1:A:338:PHE:HZ	1.48	0.77
1:A:40:GLN:HE22	1:A:197:LYS:NZ	1.88	0.72
1:A:349:GLU:O	1:A:350:ILE:HD13	1.92	0.68
1:A:209:CYS:HB3	1:A:212:GLU:HG3	1.76	0.67
1:A:248:LEU:CG	1:A:350:ILE:HD12	2.28	0.63
1:A:40:GLN:HE22	1:A:197:LYS:HZ3	1.47	0.62
1:A:263:ILE:N	1:A:263:ILE:HD12	2.15	0.61
1:A:228:GLN:HE22	1:A:346:THR:HA	1.66	0.60
1:A:353:LEU:H	1:A:353:LEU:HD22	1.65	0.60
1:A:211:SER:O	1:A:215:GLU:HG2	2.01	0.60
1:A:273:ALA:HB2	3:A:400:E64:H101	1.84	0.58
1:A:234:LEU:HD12	1:A:338:PHE:CZ	2.36	0.57
1:A:282:VAL:O	1:A:288:ARG:HA	2.04	0.57
1:A:209:CYS:HB2	3:A:400:E64:N4	2.19	0.56
1:A:212:GLU:HA	1:A:215:GLU:OE1	2.06	0.56
1:A:272:HIS:ND1	3:A:400:E64:O2	2.39	0.55
1:A:40:GLN:NE2	1:A:197:LYS:HZ3	2.04	0.55
1:A:353:LEU:N	1:A:353:LEU:HD13	2.22	0.54
1:A:256:ASP:OD1	1:A:258:ARG:HG2	2.07	0.53
1:A:154:TRP:CZ3	1:A:158:GLU:HA	2.44	0.53
1:A:184:ASN:N	1:A:184:ASN:HD22	1.96	0.52
1:A:127:ASN:C	1:A:127:ASN:HD22	2.14	0.52
1:A:318:ASP:OD1	1:A:321:GLU:HG3	2.09	0.52
1:A:109:GLN:HB2	1:A:118:LEU:HD23	1.92	0.51
1:A:300:GLU:HG3	1:A:301:VAL:HG23	1.93	0.50
1:A:80:THR:HA	1:A:83:ILE:HD12	1.95	0.49
1:A:184:ASN:ND2	1:A:184:ASN:H	2.05	0.48
1:A:282:VAL:HA	1:A:288:ARG:HH12	1.77	0.48
1:A:306:PRO:O	1:A:307:TRP:HB2	2.14	0.47
1:A:281:GLN:N	1:A:281:GLN:NE2	2.63	0.47
1:A:124:LEU:HD11	1:A:130:ILE:HG22	1.97	0.46
1:A:208:GLY:HA2	3:A:400:E64:C16	2.46	0.45
1:A:228:GLN:O	1:A:229:LYS:HD3	2.16	0.45
1:A:281:GLN:H	1:A:281:GLN:NE2	2.15	0.44
1:A:178:VAL:CG1	1:A:189:ALA:HA	2.48	0.44
1:A:115:CYS:HB3	3:A:400:E64:O1	2.12	0.44
1:A:209:CYS:N	3:A:400:E64:N4	2.65	0.44
1:A:208:GLY:C	3:A:400:E64:N4	2.71	0.43
1:A:300:GLU:O	1:A:301:VAL:HB	2.19	0.43
1:A:209:CYS:CB	3:A:400:E64:HN42	2.31	0.42
1:A:182:GLN:HG3	1:A:184:ASN:ND2	2.34	0.42

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:61:PHE:HB2	1:A:197:LYS:HG3	2.01	0.42
1:A:303:TRP:CZ2	1:A:305:GLY:HA3	2.55	0.42
1:A:263:ILE:N	1:A:263:ILE:CD1	2.83	0.41
1:A:235:TYR:O	1:A:239:LEU:HG	2.21	0.41
1:A:308:SER:OG	1:A:311:SER:HB2	2.21	0.40
1:A:124:LEU:CD1	1:A:130:ILE:HG22	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	320/339 (94%)	310 (97%)	9 (3%)	1 (0%)	50 37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	301	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	276/291 (95%)	268 (97%)	8 (3%)	55 44

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	127	ASN
1	A	133	ARG
1	A	184	ASN
1	A	256	ASP
1	A	281	GLN
1	A	353	LEU
1	A	354	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	40	GLN
1	A	52	ASN
1	A	94	ASN
1	A	96	GLN
1	A	127	ASN
1	A	152	GLN
1	A	184	ASN
1	A	228	GLN
1	A	236	GLN
1	A	281	GLN
1	A	290	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	E64	A	400	1	24,24,24	2.75	10 (41%)	30,30,30	2.55	14 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	E64	A	400	1	-	0/29/29/29	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	400	E64	C16-N3	8.68	1.49	1.33
3	A	400	E64	C16-N5	5.14	1.44	1.32
3	A	400	E64	C2-C3	4.81	1.60	1.52
3	A	400	E64	O3-C3	-3.48	1.34	1.42
3	A	400	E64	C3-C4	3.06	1.56	1.52
3	A	400	E64	C16-N4	2.70	1.37	1.32
3	A	400	E64	O2-C1	-2.55	1.21	1.30
3	A	400	E64	O1-C1	2.26	1.30	1.22
3	A	400	E64	C6-N1	-2.15	1.41	1.45
3	A	400	E64	O4-C4	2.07	1.27	1.23

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	400	E64	C14-C15-N3	-4.68	98.10	112.21
3	A	400	E64	N4-C16-N3	-4.46	106.42	119.87
3	A	400	E64	N3-C16-N5	4.46	133.71	120.69
3	A	400	E64	C15-N3-C16	4.41	134.14	124.55
3	A	400	E64	C13-C12-N2	-4.18	99.63	112.21
3	A	400	E64	O4-C4-C3	3.85	128.25	120.66
3	A	400	E64	C8-C7-C6	3.46	125.44	115.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	400	E64	O5-C11-C6	-2.83	114.09	120.31
3	A	400	E64	O2-C1-C2	2.68	123.47	114.20
3	A	400	E64	C12-N2-C11	2.53	127.78	122.57
3	A	400	E64	C6-N1-C4	2.48	127.11	121.63
3	A	400	E64	O1-C1-C2	-2.44	113.97	122.62
3	A	400	E64	O4-C4-N1	-2.15	118.79	122.93
3	A	400	E64	C7-C6-N1	2.11	115.76	110.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	322/339 (94%)	1.12	51 (15%) 3 2	20, 31, 53, 66	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	257	ILE	12.3
1	A	256	ASP	10.6
1	A	255	SER	9.5
1	A	261	GLU	8.2
1	A	259	ASP	8.0
1	A	260	LEU	7.7
1	A	258	ARG	6.6
1	A	320	TYR	6.5
1	A	265	PHE	5.7
1	A	354	THR	5.4
1	A	323	GLU	5.0
1	A	76	ASN	4.9
1	A	319	PRO	4.1
1	A	318	ASP	3.9
1	A	137	TYR	3.9
1	A	244	ARG	3.6
1	A	353	LEU	3.5
1	A	254	ILE	3.4
1	A	288	ARG	3.2
1	A	312	TYR	3.2
1	A	243	GLU	3.2
1	A	240	LYS	3.2
1	A	262	ALA	3.0
1	A	121	ILE	3.0
1	A	322	ARG	3.0
1	A	75	PRO	2.9
1	A	143	GLU	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	158	GLU	2.8
1	A	119	ALA	2.8
1	A	328	LYS	2.7
1	A	33	ASN	2.7
1	A	129	THR	2.7
1	A	232	SER	2.7
1	A	132	HIS	2.6
1	A	310	ASN	2.6
1	A	276	VAL	2.6
1	A	309	ASP	2.4
1	A	324	GLN	2.3
1	A	223	GLU	2.3
1	A	128	GLU	2.3
1	A	226	ASP	2.3
1	A	182	GLN	2.3
1	A	286	GLY	2.2
1	A	191	LEU	2.2
1	A	250	CYS	2.1
1	A	120	ALA	2.1
1	A	315	ASN	2.1
1	A	107	ILE	2.1
1	A	215	GLU	2.1
1	A	100	ASP	2.1
1	A	118	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	E64	A	400	25/25	0.33	0.92	41,46,51,52	0
2	CA	A	365	1/1	0.06	-1.73	25,25,25,25	0
2	CA	A	366	1/1	0.07	-2.19	30,30,30,30	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.