



wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 12:42 AM GMT

PDB ID : 1TOC
Title : STRUCTURE OF SERINE PROTEINASE
Authors : Van De Locht, A.; Huber, R.; Bode, W.
Deposited on : 1996-07-20
Resolution : 3.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1078 (3.16-3.04)
Ramachandran outliers	78287	1044 (3.16-3.04)
Sidechain outliers	78261	1044 (3.16-3.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	49	
1	C	49	
1	E	49	
1	G	49	
2	B	259	
2	D	259	
2	F	259	
2	H	259	
3	R	120	
3	S	120	
3	T	120	
3	U	120	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13467 atoms, of which 608 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THROMBIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	28	Total	C	H	N	O	S	10	0	0
			244	148	10	37	48	1			
1	C	28	Total	C	H	N	O	S	10	0	0
			244	148	10	37	48	1			
1	E	28	Total	C	H	N	O	S	10	0	0
			244	148	10	37	48	1			
1	G	28	Total	C	H	N	O	S	21	0	0
			244	148	10	37	48	1			

- Molecule 2 is a protein called THROMBIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	259	Total	C	H	N	O	S	147	0	0
			2206	1337	112	376	369	12			
2	D	259	Total	C	H	N	O	S	160	0	0
			2206	1337	112	376	369	12			
2	F	259	Total	C	H	N	O	S	135	0	0
			2206	1337	112	376	369	12			
2	H	259	Total	C	H	N	O	S	145	0	0
			2206	1337	112	376	369	12			

- Molecule 3 is a protein called ORNITHODORIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	R	120	Total	C	H	N	O	S	53	0	0
			909	521	30	147	197	14			
3	S	120	Total	C	H	N	O	S	50	0	0
			909	521	30	147	197	14			
3	T	120	Total	C	H	N	O	S	51	0	0
			909	521	30	147	197	14			
3	U	120	Total	C	H	N	O	S	47	0	0
			909	521	30	147	197	14			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	8	Total 8	O 8	0	0
4	D	7	Total 7	O 7	0	0
4	F	8	Total 8	O 8	0	0
4	H	8	Total 8	O 8	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: THROMBIN

Chain A: 



• Molecule 1: THROMBIN

Chain C: 




• Molecule 1: THROMBIN

Chain E: 




• Molecule 1: THROMBIN

Chain G: 




• Molecule 2: THROMBIN

Chain B: 




- Molecule 2: THROMBIN

Chain D:




- Molecule 2: THROMBIN

Chain F:




- Molecule 2: THROMBIN


Chain H:



- Molecule 3: ORNITHODORIN


Chain R:






• Molecule 3: ORNITHODORIN

Chain S:




• Molecule 3: ORNITHODORIN

Chain T:



• Molecule 3: ORNITHODORIN

Chain U:



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	125.30Å 89.90Å 101.40Å 90.00° 108.40° 90.00°	Depositor
Resolution (Å)	10.00 – 3.10	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-3.10)	Depositor
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.230 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	13467	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/237	0.65	0/316
1	C	0.49	0/237	0.67	0/316
1	E	0.56	0/237	0.70	0/316
1	G	0.52	0/237	0.67	0/316
2	B	0.57	0/2148	0.75	0/2905
2	D	0.57	0/2148	0.74	0/2905
2	F	0.54	0/2148	0.75	0/2905
2	H	0.56	0/2148	0.78	0/2905
3	R	0.64	0/895	0.75	0/1215
3	S	0.63	0/895	0.86	1/1215 (0.1%)
3	T	0.62	0/895	0.79	0/1215
3	U	0.61	0/895	0.75	0/1215
All	All	0.58	0/13120	0.76	1/17744 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	S	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	S	3	VAL	CB-CA-C	-6.82	98.43	111.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	S	22	TYR	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	234	10	217	13	0
1	C	234	10	217	18	0
1	E	234	10	217	9	0
1	G	234	10	217	13	0
2	B	2094	112	1985	111	0
2	D	2094	112	1985	128	0
2	F	2094	112	1985	121	0
2	H	2094	112	1985	124	0
3	R	879	30	738	65	0
3	S	879	30	738	60	0
3	T	879	30	738	62	0
3	U	879	30	738	59	0
4	B	8	0	0	1	0
4	D	7	0	0	1	0
4	F	8	0	0	2	0
4	H	8	0	0	2	0
All	All	12859	608	11760	687	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 29.

The worst 5 of 687 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:H:81:LYS:HE3	2:H:113:GLU:HB3	1.33	1.08
2:D:99:LEU:HD11	3:S:1:LEU:HD13	1.42	0.98
2:B:217:GLU:HB3	3:R:4:LEU:HD21	1.48	0.95
3:S:28:THR:HG23	3:S:30:LEU:HD21	1.48	0.93
3:S:70:PRO:HD2	3:S:81:TYR:CD2	2.02	0.93

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	26/49 (53%)	16 (62%)	6 (23%)	4 (15%)	0	1
1	C	26/49 (53%)	16 (62%)	5 (19%)	5 (19%)	0	0
1	E	26/49 (53%)	16 (62%)	6 (23%)	4 (15%)	0	1
1	G	26/49 (53%)	16 (62%)	5 (19%)	5 (19%)	0	0
2	B	257/259 (99%)	218 (85%)	34 (13%)	5 (2%)	12	51
2	D	257/259 (99%)	217 (84%)	35 (14%)	5 (2%)	12	51
2	F	257/259 (99%)	216 (84%)	35 (14%)	6 (2%)	10	45
2	H	257/259 (99%)	217 (84%)	35 (14%)	5 (2%)	12	51
3	R	118/120 (98%)	93 (79%)	20 (17%)	5 (4%)	4	27
3	S	118/120 (98%)	91 (77%)	21 (18%)	6 (5%)	3	22
3	T	118/120 (98%)	91 (77%)	21 (18%)	6 (5%)	3	22
3	U	118/120 (98%)	91 (77%)	22 (19%)	5 (4%)	4	27
All	All	1604/1712 (94%)	1298 (81%)	245 (15%)	61 (4%)	5	30

5 of 61 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	77(A)	ARG
2	B	149(E)	GLU
1	C	2	GLY
2	D	77(A)	ARG
2	D	149(E)	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	26/43 (60%)	24 (92%)	2 (8%)	18	57
1	C	26/43 (60%)	23 (88%)	3 (12%)	8	31
1	E	26/43 (60%)	23 (88%)	3 (12%)	8	31
1	G	26/43 (60%)	23 (88%)	3 (12%)	8	31
2	B	226/226 (100%)	214 (95%)	12 (5%)	32	72
2	D	226/226 (100%)	209 (92%)	17 (8%)	19	58
2	F	226/226 (100%)	214 (95%)	12 (5%)	32	72
2	H	226/226 (100%)	215 (95%)	11 (5%)	35	76
3	R	101/101 (100%)	82 (81%)	19 (19%)	2	9
3	S	101/101 (100%)	81 (80%)	20 (20%)	2	8
3	T	101/101 (100%)	83 (82%)	18 (18%)	2	10
3	U	101/101 (100%)	85 (84%)	16 (16%)	4	14
All	All	1412/1480 (95%)	1276 (90%)	136 (10%)	12	42

5 of 136 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	S	61	MET
2	F	27	SER
3	U	37	SER
3	S	73	CYS
3	S	95	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
3	S	15	ASN
2	F	57	HIS
3	U	14	ASN
3	S	47	GLN
2	F	159	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

6.4 Ligands

EDS was not executed - this section will therefore be empty.

6.5 Other polymers

EDS was not executed - this section will therefore be empty.