



Full wwPDB X-ray Structure Validation Report

Aug 22, 2014 – 03:26 AM EDT

PDB ID : 4TT3
Title : The Pathway of Binding of the Intrinsically Disordered Mitochondrial Inhibitor Protein to F1-ATPase
Authors : Bason, J.V.; Montgomery, M.G.; Leslie, A.G.W.; Walker, J.E.
Deposited on : 2014-06-19
Resolution : 3.21 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable23489
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable23489

1 Overall quality at a glance

The reported resolution of this entry is 3.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1205 (3.30-3.14)
Clashscore	79885	1072 (3.28-3.16)
Ramachandran outliers	78287	1052 (3.28-3.16)
Sidechain outliers	78261	1051 (3.28-3.16)
RSRZ outliers	66119	1206 (3.30-3.14)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	510	
1	B	510	
1	C	510	
2	D	480	
2	E	480	
2	F	480	
3	G	273	
4	H	66	
4	I	66	
4	J	66	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
7	GOL	A	603	-	X
7	GOL	B	603	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 24110 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP synthase subunit alpha, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	487	Total	C	N	O	S	0	0	0
			3715	2341	656	706	12			
1	B	486	Total	C	N	O	S	0	0	0
			3701	2330	654	705	12			
1	C	483	Total	C	N	O	S	0	0	0
			3684	2323	651	698	12			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	481	GLY	SER	see sequence details	UNP P19483
B	481	GLY	SER	see sequence details	UNP P19483
C	481	GLY	SER	see sequence details	UNP P19483

- Molecule 2 is a protein called ATP synthase subunit beta, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	469	Total	C	N	O	S	0	0	0
			3558	2254	605	688	11			
2	E	470	Total	C	N	O	S	0	0	0
			3563	2257	606	689	11			
2	F	469	Total	C	N	O	S	0	0	0
			3558	2254	605	688	11			

- Molecule 3 is a protein called ATP synthase subunit gamma, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	183	Total	C	N	O	S	0	0	0
			1430	894	260	269	7			


- Molecule 4 is a protein called ATPase inhibitor, mitochondrial.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	H	40	Total	C	N	O	0	0	0
			311	189	64	58			
4	I	28	Total	C	N	O	0	0	0
			236	144	49	43			
4	J	19	Total	C	N	O	0	0	0
			162	102	35	25			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	39	ALA	LYS	engineered mutation	UNP P01096
H	61	HIS	-	expression tag	UNP P01096
H	62	HIS	-	expression tag	UNP P01096
H	63	HIS	-	expression tag	UNP P01096
H	64	HIS	-	expression tag	UNP P01096
H	65	HIS	-	expression tag	UNP P01096
H	66	HIS	-	expression tag	UNP P01096
I	39	ALA	LYS	engineered mutation	UNP P01096
I	61	HIS	-	expression tag	UNP P01096
I	62	HIS	-	expression tag	UNP P01096
I	63	HIS	-	expression tag	UNP P01096
I	64	HIS	-	expression tag	UNP P01096
I	65	HIS	-	expression tag	UNP P01096
I	66	HIS	-	expression tag	UNP P01096
J	39	ALA	LYS	engineered mutation	UNP P01096
J	61	HIS	-	expression tag	UNP P01096
J	62	HIS	-	expression tag	UNP P01096
J	63	HIS	-	expression tag	UNP P01096
J	64	HIS	-	expression tag	UNP P01096
J	65	HIS	-	expression tag	UNP P01096
J	66	HIS	-	expression tag	UNP P01096

- Molecule 5 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
5	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
5	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		
6	A	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		
6	F	1	Total	Mg	0	0
			1	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	F	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	3	Total	O	0	0
			3	3		
9	B	3	Total	O	0	0
			3	3		
9	C	4	Total	O	0	0
			4	4		
9	D	5	Total	O	0	0
			5	5		
9	E	5	Total	O	0	0
			5	5		
9	F	8	Total	O	0	0
			8	8		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATP synthase subunit alpha, mitochondrial

Chain A: 




- Molecule 1: ATP synthase subunit alpha, mitochondrial

Chain B: 



- Molecule 1: ATP synthase subunit alpha, mitochondrial

Chain C: 



- Molecule 2: ATP synthase subunit beta, mitochondrial

Chain D: 



- Molecule 2: ATP synthase subunit beta, mitochondrial

Chain E: 




- Molecule 2: ATP synthase subunit beta, mitochondrial


Chain F: 




- Molecule 3: ATP synthase subunit gamma, mitochondrial




- Molecule 4: ATPase inhibitor, mitochondrial



- Molecule 4: ATPase inhibitor, mitochondrial



- Molecule 4: ATPase inhibitor, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	108.39Å 154.54Å 272.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	136.02 – 3.21 67.19 – 3.21	Depositor EDS
% Data completeness (in resolution range)	96.7 (136.02-3.21) 96.7 (67.19-3.21)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.242 , 0.283 0.243 , 0.280	Depositor DCC
R_{free} test set	3688 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	90.0	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 37.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 72985 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	24110	wwPDB-VP
Average B, all atoms (Å ²)	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, ATP, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/3766	0.42	0/5080
1	B	0.25	0/3749	0.43	0/5057
1	C	0.25	0/3733	0.41	0/5035
2	D	0.25	0/3616	0.42	0/4906
2	E	0.25	0/3621	0.43	0/4913
2	F	0.26	0/3616	0.44	1/4906 (0.0%)
3	G	0.26	0/1441	0.39	0/1923
4	H	0.29	0/315	0.37	0/417
4	I	0.31	0/239	0.40	0/315
4	J	0.30	0/165	0.42	0/218
All	All	0.25	0/24261	0.42	1/32770 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	F	1	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	477	HIS	CB-CA-C	6.21	122.83	110.40

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	F	477	HIS	CA

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3715	0	3814	4	0
1	B	3701	0	3809	4	0
1	C	3684	0	3793	2	0
2	D	3558	0	3605	2	0
2	E	3563	0	3611	5	0
2	F	3558	0	3605	2	0
3	G	1430	0	1495	0	0
4	H	311	0	298	0	0
4	I	236	0	229	0	0
4	J	162	0	163	0	0
5	A	31	0	12	0	0
5	B	31	0	12	0	0
5	C	31	0	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
7	A	6	0	8	0	0
7	B	6	0	8	0	0
8	D	27	0	12	0	0
8	F	27	0	12	0	0
9	A	3	0	0	0	0
9	B	3	0	0	0	0
9	C	4	0	0	0	0
9	D	5	0	0	0	0
9	E	5	0	0	0	0
9	F	8	0	0	0	0
All	All	24110	0	24498	19	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 0.

All (19) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:34:ILE:HD13	1:B:39:ALA:HB2	1.79	0.64
1:B:179:ALA:HB1	1:B:267:ILE:HD13	1.82	0.61
1:A:327:ILE:HD11	1:A:342:VAL:HG21	1.88	0.55
2:E:97:VAL:HG13	2:E:98:ILE:HG23	1.92	0.52
1:A:179:ALA:HB1	1:A:267:ILE:HD13	1.96	0.47
1:C:327:ILE:HD11	1:C:342:VAL:HG21	1.97	0.46
1:A:129:VAL:HG21	1:A:245:LEU:HD11	1.99	0.45
2:E:256:ASP:HA	2:E:257:ASN:HA	1.75	0.44
2:E:321:ALA:HB3	2:E:322:PRO:CD	2.48	0.44
2:D:374:VAL:HG13	2:D:410:ILE:HG21	1.99	0.43
2:E:345:TYR:HA	2:E:346:PRO:C	2.39	0.43
2:F:256:ASP:HA	2:F:257:ASN:HA	1.79	0.43
2:D:434:LEU:O	2:D:438:ILE:HD12	2.19	0.42
2:E:170:ILE:HG21	2:E:215:VAL:HG22	2.01	0.42
2:F:13:ILE:HD12	2:F:73:GLN:HB3	2.01	0.42
1:A:338:ILE:HB	1:A:339:PRO:HD3	2.00	0.42
1:B:383:MET:HG3	1:B:438:ILE:HD11	2.01	0.42
1:B:258:ARG:NH1	1:B:308:ARG:O	2.54	0.41
1:C:294:TYR:HB3	1:C:298:VAL:HG11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	485/510 (95%)	470 (97%)	15 (3%)	0	100	100
1	B	482/510 (94%)	463 (96%)	19 (4%)	0	100	100
1	C	479/510 (94%)	462 (96%)	17 (4%)	0	100	100
2	D	467/480 (97%)	442 (95%)	24 (5%)	1 (0%)	56	94
2	E	468/480 (98%)	445 (95%)	23 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	F	467/480 (97%)	439 (94%)	26 (6%)	2 (0%)	43	88
3	G	171/273 (63%)	160 (94%)	11 (6%)	0	100	100
4	H	38/66 (58%)	38 (100%)	0	0	100	100
4	I	26/66 (39%)	26 (100%)	0	0	100	100
4	J	17/66 (26%)	16 (94%)	0	1 (6%)	2	21
All	All	3100/3441 (90%)	2961 (96%)	135 (4%)	4 (0%)	59	95

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	J	32	ARG
2	D	352	ASP
2	F	352	ASP
2	F	161	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/412 (95%)	393 (100%)	0	100	100
1	B	393/412 (95%)	393 (100%)	0	100	100
1	C	390/412 (95%)	389 (100%)	1 (0%)	96	99
2	D	379/386 (98%)	379 (100%)	0	100	100
2	E	379/386 (98%)	379 (100%)	0	100	100
2	F	379/386 (98%)	379 (100%)	0	100	100
3	G	156/231 (68%)	156 (100%)	0	100	100
4	H	26/49 (53%)	26 (100%)	0	100	100
4	I	21/49 (43%)	21 (100%)	0	100	100
4	J	14/49 (29%)	14 (100%)	0	100	100
All	All	2530/2772 (91%)	2529 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	416	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	E	379	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 5 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ATP	A	601	6	33,33,33	1.06	2 (6%)	52,52,52	2.07	9 (17%)
7	GOL	A	603	-	5,5,5	0.24	0	5,5,5	0.19	0
5	ATP	B	601	6	33,33,33	1.06	2 (6%)	52,52,52	2.10	9 (17%)
7	GOL	B	603	-	5,5,5	0.20	0	5,5,5	0.29	0
5	ATP	C	600	6	33,33,33	1.08	2 (6%)	52,52,52	2.06	9 (17%)
8	ADP	D	600	6	29,29,29	1.08	2 (6%)	45,45,45	2.14	8 (17%)
8	ADP	F	600	6	29,29,29	1.11	2 (6%)	45,45,45	2.13	8 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ATP	A	601	6	-	0/22/38/38	0/3/3/3
7	GOL	A	603	-	-	0/4/4/4	0/0/0/0
5	ATP	B	601	6	-	0/22/38/38	0/3/3/3
7	GOL	B	603	-	-	0/4/4/4	0/0/0/0
5	ATP	C	600	6	-	0/22/38/38	0/3/3/3
8	ADP	D	600	6	-	0/16/32/32	0/3/3/3
8	ADP	F	600	6	-	0/16/32/32	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	600	ATP	C5-C4	3.18	1.47	1.40
5	B	601	ATP	C5-C4	3.17	1.47	1.40
8	F	600	ADP	C5-C4	3.15	1.47	1.40
5	A	601	ATP	C5-C4	3.11	1.47	1.40
8	D	600	ADP	C5-C4	3.07	1.47	1.40
5	A	601	ATP	C4-N9	-2.65	1.33	1.37
8	D	600	ADP	C4-N9	-2.62	1.33	1.37
8	F	600	ADP	C4-N9	-2.60	1.34	1.37
5	B	601	ATP	C4-N9	-2.57	1.34	1.37
5	C	600	ATP	C4-N9	-2.57	1.34	1.37

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	601	ATP	C5-C4-N3	-7.51	118.66	125.98
8	F	600	ADP	C5-C4-N3	-7.45	118.71	125.98
5	A	601	ATP	C5-C4-N3	-7.41	118.75	125.98
5	C	600	ATP	C5-C4-N3	-7.40	118.77	125.98
8	D	600	ADP	C5-C4-N3	-7.26	118.90	125.98
5	A	601	ATP	N3-C2-N1	-6.97	122.76	128.89
8	D	600	ADP	N3-C2-N1	-6.89	122.83	128.89
5	C	600	ATP	N3-C2-N1	-6.88	122.83	128.89
8	F	600	ADP	N3-C2-N1	-6.87	122.85	128.89
5	B	601	ATP	N3-C2-N1	-6.81	122.90	128.89
5	A	601	ATP	N3-C4-N9	5.64	135.07	125.39
8	F	600	ADP	N3-C4-N9	5.64	135.07	125.39
5	B	601	ATP	N3-C4-N9	5.62	135.04	125.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	600	ATP	N3-C4-N9	5.59	134.97	125.39
8	D	600	ADP	N3-C4-N9	5.48	134.79	125.39
5	B	601	ATP	PA-O3A-PB	-3.97	120.94	131.93
5	A	601	ATP	PB-O3B-PG	-3.94	121.03	131.93
5	B	601	ATP	PB-O3B-PG	-3.77	121.49	131.93
8	D	600	ADP	PA-O3A-PB	-3.67	121.77	131.93
5	C	600	ATP	PA-O3A-PB	-3.58	122.01	131.93
5	A	601	ATP	PA-O3A-PB	-3.39	122.53	131.93
8	D	600	ADP	C3'-C2'-C1'	3.32	106.12	100.92
5	C	600	ATP	PB-O3B-PG	-3.28	122.85	131.93
5	C	600	ATP	C3'-C2'-C1'	3.28	106.06	100.92
5	B	601	ATP	C3'-C2'-C1'	3.24	106.01	100.92
8	F	600	ADP	C3'-C2'-C1'	3.19	105.92	100.92
8	F	600	ADP	PA-O3A-PB	-3.15	123.20	131.93
5	B	601	ATP	C4-C5-N7	-3.06	106.45	109.41
5	C	600	ATP	C4-C5-N7	-2.97	106.54	109.41
8	D	600	ADP	C4-C5-N7	-2.96	106.55	109.41
8	F	600	ADP	C4-C5-N7	-2.90	106.60	109.41
5	A	601	ATP	C4-C5-N7	-2.87	106.64	109.41
5	A	601	ATP	C3'-C2'-C1'	2.81	105.33	100.92
5	B	601	ATP	C2-N3-C4	2.81	121.36	113.27
5	C	600	ATP	C2-N3-C4	2.81	121.36	113.27
5	A	601	ATP	C2-N3-C4	2.80	121.32	113.27
8	D	600	ADP	C2-N3-C4	2.79	121.29	113.27
8	F	600	ADP	C2-N3-C4	2.79	121.29	113.27
5	A	601	ATP	C8-N9-C4	2.17	108.72	106.96
5	C	600	ATP	C8-N9-C4	2.13	108.69	106.96
8	D	600	ADP	C8-N9-C4	2.08	108.65	106.96
8	F	600	ADP	C8-N9-C4	2.05	108.63	106.96
5	B	601	ATP	C8-N9-C4	2.03	108.61	106.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	487/510 (95%)	-0.04	0 100 100	58, 80, 115, 164	0
1	B	486/510 (95%)	-0.09	0 100 100	55, 75, 107, 130	0
1	C	483/510 (94%)	0.06	0 100 100	67, 97, 122, 151	0
2	D	469/480 (97%)	0.16	1 (0%) 93 67	74, 102, 131, 144	0
2	E	470/480 (97%)	0.07	4 (0%) 81 35	56, 78, 126, 164	0
2	F	469/480 (97%)	0.07	1 (0%) 93 67	66, 95, 118, 141	0
3	G	183/273 (67%)	0.82	20 (10%) 6 2	73, 135, 167, 178	0
4	H	40/66 (60%)	0.14	0 100 100	92, 111, 132, 138	0
4	I	28/66 (42%)	0.84	2 (7%) 16 4	133, 140, 149, 153	0
4	J	19/66 (28%)	1.06	1 (5%) 25 5	148, 151, 158, 159	0
All	All	3134/3441 (91%)	0.10	29 (0%) 81 35	55, 90, 140, 178	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	393	MET	4.9
3	G	149	LEU	3.5
2	E	391	LEU	3.2
2	E	394	ASP	3.0
3	G	213	ILE	3.0
3	G	108	VAL	3.0
3	G	112	ILE	3.0
2	F	477	HIS	2.9
3	G	136	PRO	2.8
3	G	215	TYR	2.6
3	G	207	TYR	2.5
3	G	217	LEU	2.5
4	I	48	HIS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	G	138	PHE	2.4
2	D	477	HIS	2.4
3	G	166	ARG	2.4
3	G	209	LEU	2.3
4	I	46	LYS	2.3
3	G	212	ILE	2.1
3	G	146	LEU	2.1
2	E	392	GLY	2.1
3	G	145	ALA	2.1
3	G	89	MET	2.1
4	J	37	ARG	2.1
3	G	210	ALA	2.1
3	G	148	LEU	2.1
3	G	137	THR	2.0
3	G	211	ASN	2.0
3	G	143	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	GOL	B	603	6/6	0.34	3.02	75,76,76,77	0
7	GOL	A	603	6/6	0.31	2.02	72,73,73,74	0
6	MG	C	601	1/1	0.22	1.80	83,83,83,83	0
8	ADP	F	600	27/27	0.22	0.08	76,82,86,86	0
5	ATP	C	600	31/31	0.23	-0.19	83,93,100,101	4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	ATP	B	601	31/31	0.18	-0.44	63,72,78,78	0
8	ADP	D	600	27/27	0.21	-0.73	85,88,90,91	0
5	ATP	A	601	31/31	0.18	-1.05	72,74,76,76	4
6	MG	F	601	1/1	0.16	-1.47	78,78,78,78	0
6	MG	D	601	1/1	0.16	-1.47	88,88,88,88	0
6	MG	A	602	1/1	0.15	-2.12	75,75,75,75	0
6	MG	B	602	1/1	0.17	-3.74	63,63,63,63	0

6.5 Other polymers

There are no such residues in this entry.