



# Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 01:44 AM GMT

PDB ID : 1TXN  
Title : Crystal structure of coproporphyrinogen III oxidase  
Authors : Kumaran, D.; Swaminathan, S.; Burley, S.K.; New York SGX Research Center  
for Structural Genomics (NYSGXRC)  
Deposited on : 2004-07-05  
Resolution : 1.70 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2456 (1.70-1.70)
Clashscore	79885	2929 (1.70-1.70)
Ramachandran outliers	78287	2878 (1.70-1.70)
Sidechain outliers	78261	2878 (1.70-1.70)
RSRZ outliers	66119	2456 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	328	
1	B	328	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	GOL	A	500	X	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4636 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coproporphyrinogen III oxidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	Se	0	0	0
			2112	1342	369	392	3	6			
1	B	256	Total	C	N	O	S	Se	0	0	0
			2087	1327	365	386	3	6			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P11353
A	16	MSE	MET	MODIFIED RESIDUE	UNP P11353
A	56	MSE	MET	MODIFIED RESIDUE	UNP P11353
A	86	MSE	MET	MODIFIED RESIDUE	UNP P11353
A	118	MSE	MET	MODIFIED RESIDUE	UNP P11353
A	227	MSE	MET	MODIFIED RESIDUE	UNP P11353
A	249	MSE	MET	MODIFIED RESIDUE	UNP P11353
A	293	MSE	MET	MODIFIED RESIDUE	UNP P11353
B	1	MSE	MET	MODIFIED RESIDUE	UNP P11353
B	16	MSE	MET	MODIFIED RESIDUE	UNP P11353
B	56	MSE	MET	MODIFIED RESIDUE	UNP P11353
B	86	MSE	MET	MODIFIED RESIDUE	UNP P11353
B	118	MSE	MET	MODIFIED RESIDUE	UNP P11353
B	227	MSE	MET	MODIFIED RESIDUE	UNP P11353
B	249	MSE	MET	MODIFIED RESIDUE	UNP P11353
B	293	MSE	MET	MODIFIED RESIDUE	UNP P11353

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	207	Total	O	0	0
			207	207		
3	B	224	Total	O	0	0
			224	224		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Coproporphyrinogen III oxidase



#### • Molecule 1: Coproporphyrinogen III oxidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.75Å 71.75Å 117.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.70 45.51 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.4 (50.00-1.70) 93.4 (45.51-1.70)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.77 (at 1.70Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.211 , 0.245 0.210 , 0.244	Depositor DCC
$R_{free}$ test set	3713 reflections (6.09%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.1	Xtriage
Anisotropy	0.320	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 24.7	EDS
Estimated twinning fraction	0.168 for h,-k,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	1 of 61508 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4636	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/2166	0.60	1/2923 (0.0%)
1	B	0.32	0/2140	0.59	1/2887 (0.0%)
All	All	0.32	0/4306	0.60	2/5810 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	SER	N-CA-C	-6.54	93.34	111.00
1	B	160	SER	N-CA-C	-6.09	94.57	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2112	0	2014	38	0
1	B	2087	0	1987	47	0
2	A	6	0	4	2	0
3	A	207	0	0	2	0
3	B	224	0	0	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4636	0	4005	81	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

All (81) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:6:ASP:H	1:A:7:PRO:HD2	1.31	0.93
1:B:12:ILE:HG23	1:B:225:LEU:HD21	1.51	0.93
1:B:15:GLN:HE22	1:B:222:GLN:HE22	1.23	0.86
1:A:202:ARG:HG3	1:A:204:GLU:HG3	1.66	0.77
1:B:129:THR:HB	1:B:263:ARG:HH22	1.52	0.75
1:B:6:ASP:N	1:B:7:PRO:HD3	2.05	0.70
1:A:15:GLN:HE22	1:A:222:GLN:HE22	1.37	0.70
1:B:6:ASP:H	1:B:7:PRO:HD3	1.55	0.70
1:A:6:ASP:H	1:A:7:PRO:CD	2.06	0.66
1:A:133:ASN:C	1:A:133:ASN:HD22	1.99	0.65
1:B:143:ASP:OD1	1:B:145:THR:HG22	1.95	0.65
1:A:133:ASN:HD22	1:A:134:TYR:N	1.95	0.65
1:B:221:PRO:O	1:B:225:LEU:HG	1.97	0.65
1:B:267:VAL:HA	1:B:270:ASN:OD1	1.98	0.62
1:B:95:LEU:HB3	1:B:96:PRO:HD3	1.80	0.62
1:A:173:LEU:HD12	1:A:238:SER:HB3	1.81	0.61
1:B:264:GLY:O	1:B:268:GLU:HG3	2.03	0.59
1:A:192:TRP:CE3	1:A:195:GLU:HG3	2.38	0.58
1:B:170:PHE:O	1:B:174:HIS:HD2	1.87	0.58
1:B:16:MSE:HE2	1:B:16:MSE:HA	1.86	0.57
1:A:213:PHE:HA	2:A:500:GOL:H12	1.86	0.57
1:B:78:LEU:HD11	1:B:137:PHE:HZ	1.70	0.56
1:B:206:ARG:HH11	1:B:206:ARG:HG2	1.71	0.56
1:A:254:GLU:HG2	1:B:76:GLY:HA3	1.88	0.56
1:B:174:HIS:HE1	3:B:333:HOH:O	1.90	0.55
1:B:129:THR:CB	1:B:263:ARG:HH22	2.18	0.55
1:B:83:VAL:O	1:B:87:LYS:HG3	2.07	0.54
1:B:259:GLN:HE22	1:B:262:ARG:HE	1.57	0.53
1:A:174:HIS:HE1	3:A:521:HOH:O	1.92	0.52
1:A:206:ARG:CZ	1:B:88:ALA:HB3	2.40	0.52
1:B:66:LYS:HG2	1:B:269:PHE:HB2	1.91	0.52
1:B:165:GLU:HG2	1:B:245:ARG:NH2	2.25	0.51
1:A:170:PHE:O	1:A:174:HIS:HD2	1.93	0.51
1:B:173:LEU:HD12	1:B:238:SER:HB3	1.91	0.51
1:A:52:GLY:H	1:A:73:VAL:H	1.59	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:219:ARG:HD3	3:B:421:HOH:O	2.11	0.50
1:A:186:TYR:HB3	1:A:187:PRO:HD3	1.94	0.50
1:A:3:ALA:N	1:A:4:PRO:CD	2.75	0.50
1:B:206:ARG:NH1	1:B:206:ARG:HG2	2.26	0.49
1:A:6:ASP:N	1:A:7:PRO:HD2	2.13	0.49
1:A:128:PRO:HD3	3:A:515:HOH:O	2.11	0.49
1:A:76:GLY:HA3	1:B:254:GLU:HG2	1.94	0.49
1:B:46:ARG:HD2	1:B:49:ASP:HA	1.95	0.48
1:B:28:THR:O	1:B:32:GLU:HG3	2.14	0.48
1:A:35:ASP:OD1	1:A:36:THR:N	2.43	0.47
1:B:6:ASP:N	1:B:7:PRO:CD	2.77	0.47
1:A:182:ASP:HB3	1:A:185:LEU:HG	1.96	0.47
1:B:16:MSE:HE3	1:B:225:LEU:HD23	1.95	0.47
1:A:214:ASP:H	2:A:500:GOL:H11	1.81	0.46
1:B:182:ASP:HB3	1:B:185:LEU:HG	1.98	0.46
1:B:195:GLU:O	1:B:199:ILE:HG13	2.16	0.45
1:A:15:GLN:NE2	1:A:222:GLN:HE22	2.11	0.45
1:B:186:TYR:HB3	1:B:187:PRO:HD3	1.98	0.45
1:A:77:GLN:HG2	1:A:110:LYS:HG2	1.98	0.45
1:A:162:LEU:HD23	1:A:206:ARG:CZ	2.46	0.45
1:A:48:ASN:HA	1:A:48:ASN:HD22	1.60	0.45
1:B:86:MSE:HE3	3:B:513:HOH:O	2.17	0.45
1:A:3:ALA:H	1:A:4:PRO:CD	2.30	0.44
1:B:259:GLN:NE2	1:B:262:ARG:HE	2.16	0.44
1:B:16:MSE:HE3	1:B:151:PHE:HE2	1.82	0.43
1:A:94:ARG:HG2	1:A:96:PRO:HD2	2.00	0.43
1:B:118:MSE:CG	1:B:132:LEU:HB3	2.49	0.43
1:A:42:ASP:HB3	1:A:55:SER:HB3	2.00	0.43
1:A:118:MSE:HG2	1:A:132:LEU:HB3	2.01	0.43
1:A:149:TRP:CZ2	1:A:221:PRO:HD3	2.54	0.42
1:B:70:ASN:O	1:B:116:LEU:HA	2.19	0.42
1:A:245:ARG:HG3	1:A:246:ARG:HG2	2.02	0.42
1:A:94:ARG:HH21	1:A:147:GLN:CD	2.22	0.42
1:B:13:ARG:HD3	1:B:138:GLU:OE2	2.19	0.42
1:B:80:PRO:HA	1:B:95:LEU:HD13	2.02	0.42
1:A:133:ASN:ND2	1:A:133:ASN:C	2.71	0.42
1:B:7:PRO:O	1:B:8:ARG:HG3	2.20	0.42
1:B:160:SER:HA	1:B:262:ARG:HH21	1.84	0.42
1:A:94:ARG:H	1:A:94:ARG:HD3	1.85	0.41
1:B:132:LEU:HD23	1:B:132:LEU:C	2.41	0.41
1:A:267:VAL:HB	1:B:119:VAL:HG12	2.03	0.41
1:B:121:HIS:HB3	1:B:265:ARG:NH2	2.35	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:118:MSE:CG	1:A:132:LEU:HB3	2.51	0.41
1:A:162:LEU:HD22	1:A:206:ARG:HB2	2.03	0.40
1:B:266:TYR:HA	1:B:269:PHE:CE1	2.56	0.40
1:B:222:GLN:HA	1:B:225:LEU:HD12	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/328 (78%)	246 (96%)	7 (3%)	2 (1%)	27	8
1	B	250/328 (76%)	237 (95%)	10 (4%)	3 (1%)	19	3
All	All	505/656 (77%)	483 (96%)	17 (3%)	5 (1%)	22	5

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	6	ASP
1	B	7	PRO
1	A	3	ALA
1	B	8	ARG
1	B	6	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/276 (81%)	217 (97%)	6 (3%)	57	33

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	220/276 (80%)	218 (99%)	2 (1%)	87	79
All	All	443/552 (80%)	435 (98%)	8 (2%)	71	53

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	THR
1	A	48	ASN
1	A	78	LEU
1	A	94	ARG
1	A	133	ASN
1	A	194	ASP
1	B	9	ASN
1	B	128	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	24	GLN
1	A	40	HIS
1	A	48	ASN
1	A	133	ASN
1	A	141	ASN
1	A	174	HIS
1	A	256	GLN
1	B	9	ASN
1	B	15	GLN
1	B	29	GLN
1	B	121	HIS
1	B	147	GLN
1	B	174	HIS
1	B	256	GLN
1	B	259	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	500	-	5,5,5	4.54	5 (100%)	5,5,5	5.73	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	500	-	-	0/4/4/4	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	GOL	C3-C2	-7.69	1.20	1.52
2	A	500	GOL	O1-C1	4.38	1.61	1.42
2	A	500	GOL	O3-C3	3.10	1.55	1.42
2	A	500	GOL	C1-C2	-2.83	1.40	1.52
2	A	500	GOL	O2-C2	-2.65	1.35	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	500	GOL	O3-C3-C2	10.38	160.37	109.71
2	A	500	GOL	O2-C2-C3	6.72	138.82	108.22
2	A	500	GOL	O1-C1-C2	3.30	125.81	109.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	259/328 (78%)	1.30	43 (16%) 2 3	8, 16, 39, 43	0
1	B	256/328 (78%)	1.31	39 (15%) 3 4	8, 16, 39, 43	0
All	All	515/656 (78%)	1.30	82 (15%) 3 4	8, 16, 39, 43	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	3	ALA	18.2
1	B	44	TRP	18.1
1	A	4	PRO	16.5
1	A	3	ALA	16.1
1	A	44	TRP	16.0
1	B	51	GLY	15.2
1	A	2	PRO	14.9
1	A	5	GLN	14.7
1	B	4	PRO	14.1
1	B	2	PRO	13.8
1	A	45	THR	12.7
1	B	6	ASP	12.4
1	B	48	ASN	12.0
1	B	45	THR	11.7
1	A	95	LEU	11.2
1	A	46	ARG	11.2
1	B	47	GLY	10.7
1	B	50	GLY	10.6
1	B	269	PHE	10.5
1	B	7	PRO	10.4
1	A	266	TYR	10.0
1	A	49	ASP	9.9
1	A	6	ASP	9.8
1	B	95	LEU	9.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	49	ASP	9.6
1	A	52	GLY	8.9
1	B	5	GLN	8.7
1	B	266	TYR	8.7
1	A	7	PRO	8.4
1	B	270	ASN	8.3
1	B	202	ARG	8.0
1	A	53	GLY	7.7
1	B	52	GLY	7.6
1	B	46	ARG	7.5
1	A	48	ASN	6.7
1	B	43	THR	6.4
1	A	50	GLY	6.3
1	A	47	GLY	6.3
1	A	269	PHE	6.3
1	A	8	ARG	6.2
1	A	51	GLY	6.0
1	B	265	ARG	6.0
1	A	94	ARG	5.8
1	A	43	THR	5.7
1	A	96	PRO	5.3
1	A	267	VAL	5.1
1	B	206	ARG	5.0
1	B	8	ARG	4.9
1	B	96	PRO	4.7
1	B	107	ASP	4.6
1	A	107	ASP	4.6
1	A	268	GLU	4.4
1	B	198	TYR	4.3
1	B	268	GLU	4.1
1	A	89	ASP	4.0
1	A	264	GLY	4.0
1	A	270	ASN	3.8
1	B	94	ARG	3.6
1	A	88	ALA	3.3
1	A	54	THR	3.2
1	B	54	THR	3.2
1	A	265	ARG	3.2
1	B	197	PHE	3.1
1	B	53	GLY	3.0
1	B	225	LEU	2.8
1	B	267	VAL	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	87	LYS	2.6
1	A	198	TYR	2.6
1	A	108	GLY	2.6
1	B	201	HIS	2.6
1	A	77	GLN	2.5
1	A	84	SER	2.5
1	A	42	ASP	2.5
1	A	203	LYS	2.4
1	B	200	THR	2.4
1	A	196	TYR	2.3
1	B	9	ASN	2.3
1	B	87	LYS	2.2
1	B	42	ASP	2.2
1	A	41	ALA	2.2
1	A	81	ALA	2.2
1	A	142	GLN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	A	500	6/6	0.23	5.41	19,24,31,32	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.