



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 05:00 AM GMT

PDB ID : 1TYH
Title : Crystal Structure of Transcriptional Activator tenA from Bacillus subtilis
Authors : Eswaramoorthy, S.; Swaminathan, S.; Burley, S.K.; New York SGX Research
Center for Structural Genomics (NYSGXRC)
Deposited on : 2004-07-07
Resolution : 2.54 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	4284 (2.58-2.50)
Ramachandran outliers	78287	4193 (2.58-2.50)
Sidechain outliers	78261	4195 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	248	
1	B	248	
1	D	248	
1	E	248	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7252 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional activator tenA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S	Se	0	0	0
			1783	1154	287	334	3	5			
1	B	221	Total	C	N	O	S	Se	0	0	0
			1783	1154	287	334	3	5			
1	D	221	Total	C	N	O	S	Se	0	0	0
			1791	1158	287	338	3	5			
1	E	221	Total	C	N	O	S	Se	0	0	0
			1783	1154	287	334	3	5			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	CLONING ARTIFACT	UNP P25052
A	0	SER	-	CLONING ARTIFACT	UNP P25052
A	1	LEU	-	CLONING ARTIFACT	UNP P25052
A	71	MSE	MET	MODIFIED RESIDUE	UNP P25052
A	83	MSE	MET	MODIFIED RESIDUE	UNP P25052
A	116	MSE	MET	MODIFIED RESIDUE	UNP P25052
A	194	MSE	MET	MODIFIED RESIDUE	UNP P25052
A	211	MSE	MET	MODIFIED RESIDUE	UNP P25052
A	237	GLU	-	EXPRESSION TAG	UNP P25052
A	238	GLY	-	EXPRESSION TAG	UNP P25052
A	239	GLY	-	EXPRESSION TAG	UNP P25052
A	240	SER	-	EXPRESSION TAG	UNP P25052
A	241	HIS	-	EXPRESSION TAG	UNP P25052
A	242	HIS	-	EXPRESSION TAG	UNP P25052
A	243	HIS	-	EXPRESSION TAG	UNP P25052
A	244	HIS	-	EXPRESSION TAG	UNP P25052
A	245	HIS	-	EXPRESSION TAG	UNP P25052
A	246	HIS	-	EXPRESSION TAG	UNP P25052
B	-1	MSE	-	CLONING ARTIFACT	UNP P25052
B	0	SER	-	CLONING ARTIFACT	UNP P25052
B	1	LEU	-	CLONING ARTIFACT	UNP P25052

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	71	MSE	MET	MODIFIED RESIDUE	UNP P25052
B	83	MSE	MET	MODIFIED RESIDUE	UNP P25052
B	116	MSE	MET	MODIFIED RESIDUE	UNP P25052
B	194	MSE	MET	MODIFIED RESIDUE	UNP P25052
B	211	MSE	MET	MODIFIED RESIDUE	UNP P25052
B	237	GLU	-	EXPRESSION TAG	UNP P25052
B	238	GLY	-	EXPRESSION TAG	UNP P25052
B	239	GLY	-	EXPRESSION TAG	UNP P25052
B	240	SER	-	EXPRESSION TAG	UNP P25052
B	241	HIS	-	EXPRESSION TAG	UNP P25052
B	242	HIS	-	EXPRESSION TAG	UNP P25052
B	243	HIS	-	EXPRESSION TAG	UNP P25052
B	244	HIS	-	EXPRESSION TAG	UNP P25052
B	245	HIS	-	EXPRESSION TAG	UNP P25052
B	246	HIS	-	EXPRESSION TAG	UNP P25052
D	-1	MSE	-	CLONING ARTIFACT	UNP P25052
D	0	SER	-	CLONING ARTIFACT	UNP P25052
D	1	LEU	-	CLONING ARTIFACT	UNP P25052
D	71	MSE	MET	MODIFIED RESIDUE	UNP P25052
D	83	MSE	MET	MODIFIED RESIDUE	UNP P25052
D	116	MSE	MET	MODIFIED RESIDUE	UNP P25052
D	194	MSE	MET	MODIFIED RESIDUE	UNP P25052
D	211	MSE	MET	MODIFIED RESIDUE	UNP P25052
D	237	GLU	-	EXPRESSION TAG	UNP P25052
D	238	GLY	-	EXPRESSION TAG	UNP P25052
D	239	GLY	-	EXPRESSION TAG	UNP P25052
D	240	SER	-	EXPRESSION TAG	UNP P25052
D	241	HIS	-	EXPRESSION TAG	UNP P25052
D	242	HIS	-	EXPRESSION TAG	UNP P25052
D	243	HIS	-	EXPRESSION TAG	UNP P25052
D	244	HIS	-	EXPRESSION TAG	UNP P25052
D	245	HIS	-	EXPRESSION TAG	UNP P25052
D	246	HIS	-	EXPRESSION TAG	UNP P25052
E	-1	MSE	-	CLONING ARTIFACT	UNP P25052
E	0	SER	-	CLONING ARTIFACT	UNP P25052
E	1	LEU	-	CLONING ARTIFACT	UNP P25052
E	71	MSE	MET	MODIFIED RESIDUE	UNP P25052
E	83	MSE	MET	MODIFIED RESIDUE	UNP P25052
E	116	MSE	MET	MODIFIED RESIDUE	UNP P25052
E	194	MSE	MET	MODIFIED RESIDUE	UNP P25052
E	211	MSE	MET	MODIFIED RESIDUE	UNP P25052
E	237	GLU	-	EXPRESSION TAG	UNP P25052

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	238	GLY	-	EXPRESSION TAG	UNP P25052
E	239	GLY	-	EXPRESSION TAG	UNP P25052
E	240	SER	-	EXPRESSION TAG	UNP P25052
E	241	HIS	-	EXPRESSION TAG	UNP P25052
E	242	HIS	-	EXPRESSION TAG	UNP P25052
E	243	HIS	-	EXPRESSION TAG	UNP P25052
E	244	HIS	-	EXPRESSION TAG	UNP P25052
E	245	HIS	-	EXPRESSION TAG	UNP P25052
E	246	HIS	-	EXPRESSION TAG	UNP P25052

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	36	Total O 36 36	0	0
2	B	34	Total O 34 34	0	0
2	D	17	Total O 17 17	0	0
2	E	25	Total O 25 25	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Transcriptional activator tenA




• Molecule 1: Transcriptional activator tenA




• Molecule 1: Transcriptional activator tenA





● Molecule 1: Transcriptional activator tenA

Chain E:



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	58.43Å 58.43Å 297.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.54	Depositor
% Data completeness (in resolution range)	96.9 (50.00-2.54)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.244 , 0.304	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7252	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/1836	0.60	1/2483 (0.0%)
1	B	0.44	0/1836	0.60	1/2483 (0.0%)
1	D	0.46	0/1844	0.62	0/2493
1	E	0.45	0/1836	0.62	0/2483
All	All	0.45	0/7352	0.61	2/9942 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	214	ARG	N-CA-C	-5.92	95.01	111.00
1	A	214	ARG	N-CA-C	-5.90	95.06	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1783	0	1625	50	0
1	B	1783	0	1625	56	0
1	D	1791	0	1633	144	0
1	E	1783	0	1625	136	0
2	A	36	0	0	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	34	0	0	7	0
2	D	17	0	0	5	0
2	E	25	0	0	7	0
All	All	7252	0	6508	374	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 27.

The worst 5 of 374 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:99:GLU:HG3	2:A:268:HOH:O	1.46	1.11
1:E:151:PRO:HG2	1:E:157:GLN:OE1	1.70	0.90
1:A:220:ASP:HB2	2:A:272:HOH:O	1.71	0.90
1:D:155:ILE:HG23	1:D:156:TYR:CD1	2.09	0.88
1:A:86:HIS:CB	2:A:271:HOH:O	2.22	0.87

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/248 (88%)	208 (95%)	9 (4%)	2 (1%)	25	41
1	B	219/248 (88%)	206 (94%)	8 (4%)	5 (2%)	10	13
1	D	219/248 (88%)	180 (82%)	30 (14%)	9 (4%)	4	4
1	E	219/248 (88%)	181 (83%)	30 (14%)	8 (4%)	5	6
All	All	876/992 (88%)	775 (88%)	77 (9%)	24 (3%)	8	10

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	30	GLY
1	D	95	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	187	THR
1	E	95	ILE
1	E	96	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/201 (87%)	170 (97%)	5 (3%)	55	80
1	B	175/201 (87%)	169 (97%)	6 (3%)	49	75
1	D	177/201 (88%)	161 (91%)	16 (9%)	14	24
1	E	175/201 (87%)	167 (95%)	8 (5%)	37	60
All	All	702/804 (87%)	667 (95%)	35 (5%)	34	56

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	55	GLN
1	D	139	TYR
1	E	185	ASN
1	D	63	LYS
1	D	97	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	55	GLN
1	D	148	HIS
1	E	55	GLN
1	D	25	GLN
1	E	25	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.