



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 26, 2014 – 04:56 PM GMT

PDB ID : 3U3Q
Title : The S-SAD phased crystal structure of the ecto-domain of Death Receptor 6 (DR6)
Authors : Ru, H.; Zhao, L.X.; Ding, W.; Jiao, L.Y.; Shaw, N.; Zhang, L.G.; Hung, L.W.; Matsugaki, N.; Wakatsuki, S.; Liu, Z.J.
Deposited on : 2011-10-06
Resolution : 2.70 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1557 (2.70-2.70)
Clashscore	79885	1939 (2.70-2.70)
Ramachandran outliers	78287	1905 (2.70-2.70)
Sidechain outliers	78261	1905 (2.70-2.70)
RSRZ outliers	66119	1559 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	313	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1375 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor receptor superfamily member 21.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	164	Total	C	N	O	S	0	1	0
			1233	754	220	238	21			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	349	HIS	-	EXPRESSION TAG	UNP O75509
A	350	HIS	-	EXPRESSION TAG	UNP O75509
A	351	HIS	-	EXPRESSION TAG	UNP O75509
A	352	HIS	-	EXPRESSION TAG	UNP O75509
A	353	HIS	-	EXPRESSION TAG	UNP O75509
A	354	HIS	-	EXPRESSION TAG	UNP O75509

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	142	Total	O	0	0
			142	142		

4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.95Å 77.95Å 186.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.70 38.15 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.70) 99.9 (38.15-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.96 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.195 , 0.235 0.195 , 0.235	Depositor DCC
R_{free} test set	471 reflections (5.05%)	DCC
Wilson B-factor (Å ²)	45.8	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 9806 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1375	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	2/1262 (0.2%)	1.04	3/1717 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	115	TRP	CD2-CE2	6.18	1.48	1.41
1	A	154	TRP	CD2-CE2	5.35	1.47	1.41

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	LEU	CB-CG-CD1	-6.82	99.41	111.00
1	A	98	ARG	NE-CZ-NH1	6.04	123.32	120.30
1	A	191	ASP	CB-CG-OD2	5.48	123.23	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1233	0	1190	26	0
2	A	142	0	0	9	0
All	All	1375	0	1190	26	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 11.

All (26) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:201:LYS:HB3	1:A:210:VAL:HG13	1.68	0.75
1:A:193:LEU:C	1:A:195:GLN:H	2.05	0.60
1:A:193:LEU:O	1:A:195:GLN:N	2.36	0.57
1:A:103:ILE:HG12	2:A:501:HOH:O	2.05	0.55
1:A:191:ASP:HB3	1:A:194:SER:OG	2.06	0.54
1:A:92:PRO:HD2	1:A:95:THR:OG1	2.09	0.52
1:A:90:SER:HB2	2:A:403:HOH:O	2.09	0.52
1:A:179[A]:VAL:HG11	2:A:511:HOH:O	2.09	0.52
1:A:195:GLN:O	1:A:196:ASN:HB2	2.11	0.51
1:A:90:SER:CB	2:A:403:HOH:O	2.58	0.51
1:A:170:GLN:OE1	2:A:503:HOH:O	2.19	0.50
1:A:191:ASP:O	1:A:195:GLN:HB2	2.11	0.50
1:A:201:LYS:HB3	1:A:210:VAL:CG1	2.42	0.49
1:A:103:ILE:CD1	1:A:104:GLU:H	2.25	0.49
1:A:54:TYR:CE1	1:A:67:CYS:HB2	2.47	0.49
1:A:126:LEU:HB3	2:A:420:HOH:O	2.13	0.49
1:A:185:LYS:NZ	2:A:500:HOH:O	2.47	0.47
1:A:158:LYS:HB3	1:A:167:ARG:HB2	1.98	0.45
1:A:193:LEU:C	1:A:195:GLN:N	2.70	0.45
1:A:192:CYS:O	1:A:195:GLN:HB3	2.18	0.44
1:A:159:LYS:HG3	2:A:474:HOH:O	2.17	0.44
1:A:201:LYS:CB	1:A:210:VAL:HG13	2.43	0.44
1:A:167:ARG:NE	2:A:510:HOH:O	2.46	0.43
1:A:74:THR:HG22	1:A:90:SER:HA	2.01	0.43
1:A:103:ILE:HD12	1:A:104:GLU:H	1.83	0.42
1:A:143:THR:HB	1:A:144:CYS:H	1.29	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/313 (52%)	152 (93%)	7 (4%)	4 (2%)	9	21

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	194	SER
1	A	125	ALA
1	A	193	LEU
1	A	142	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/279 (52%)	134 (92%)	11 (8%)	19	41

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	ARG
1	A	103	ILE
1	A	105	LYS
1	A	129	ARG
1	A	143	THR
1	A	164	GLU
1	A	173	ARG
1	A	193	LEU
1	A	210	VAL
1	A	213	THR
1	A	214	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	196	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	164/313 (52%)	-0.21	5 (3%) 48 54	19, 38, 84, 108	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	115	TRP	3.0
1	A	200	ILE	3.0
1	A	141	ASN	2.2
1	A	142	ALA	2.1
1	A	213	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.