



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 04:56 PM GMT

PDB ID : 3U3Y
Title : Mouse TREX1 D200H mutant
Authors : Bailey, S.L.; Harvey, S.; Perrino, F.W.; Hollis, T.
Deposited on : 2011-10-06
Resolution : 2.28 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3861 (2.30-2.26)
Clashscore	79885	4801 (2.30-2.26)
Ramachandran outliers	78287	4729 (2.30-2.26)
Sidechain outliers	78261	4728 (2.30-2.26)
RSRZ outliers	66119	3864 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	314	
1	B	314	
2	C	4	
2	D	4	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	BU1	A	315	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7123 atoms, of which 3388 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Three prime repair exonuclease 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	220	Total	C	H	N	O	S	0	1	0
			3408	1082	1702	301	314	9			
1	A	215	Total	C	H	N	O	S	0	1	0
			3352	1066	1676	295	306	9			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	200	HIS	ASP	ENGINEERED MUTATION	UNP Q91XB0
A	200	HIS	ASP	ENGINEERED MUTATION	UNP Q91XB0

- Molecule 2 is a DNA chain called 5'-D(*GP*AP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	4	Total	C	N	O	P	0	0	0
			81	39	18	21	3			
2	D	4	Total	C	N	O	P	0	0	0
			81	39	18	21	3			

- Molecule 3 is 1,4-BUTANEDIOL (three-letter code: BU1) (formula: C₄H₁₀O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			16	4	10	2		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	76	Total	O	0	0
			76	76		
5	A	88	Total	O	0	0
			88	88		
5	C	7	Total	O	0	0
			7	7		
5	D	12	Total	O	0	0
			12	12		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Three prime repair exonuclease 1



- Molecule 1: Three prime repair exonuclease 1



- Molecule 2: 5'-D(*GP*AP*CP*G)-3'



- Molecule 2: 5'-D(*GP*AP*CP*G)-3'

Chain D:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.07Å 57.86Å 68.23Å 90.00° 108.11° 90.00°	Depositor
Resolution (Å)	22.72 – 2.28 22.72 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.2 (22.72-2.28) 99.4 (22.72-2.28)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.42 (at 2.28Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.213 , 0.275 0.209 , 0.278	Depositor DCC
R_{free} test set	1135 reflections (5.16%)	DCC
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.455	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 38.5	EDS
Estimated twinning fraction	0.012 for l,-k,h	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 22059 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7123	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, BU1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	1/1721 (0.1%)	0.58	1/2345 (0.0%)
1	B	0.36	0/1751	0.54	0/2386
2	C	0.74	0/91	1.14	0/139
2	D	0.74	0/91	1.20	0/139
All	All	0.40	1/3654 (0.0%)	0.61	1/5009 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	54	PRO	C-N	-5.47	1.23	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	164	GLN	N-CA-CB	-5.52	100.67	110.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	99	ARG	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1676	1676	0	49	0
1	B	1706	1702	0	44	0
2	C	81	0	46	0	0
2	D	81	0	46	2	0
3	A	6	10	0	10	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	88	0	0	1	0
5	B	76	0	0	1	0
5	C	7	0	0	0	0
5	D	12	0	0	1	0
All	All	3735	3388	92	86	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

All (86) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:202:LEU:HD12	3:A:315:BU1:H21	1.58	0.85
1:A:212:PRO:O	1:A:216:LEU:HD22	1.90	0.71
1:A:192:THR:HB	3:A:315:BU1:C3	2.21	0.71
1:A:164:GLN:N	1:A:164:GLN:OE1	2.24	0.70
1:A:159:LEU:HD23	1:A:182:ILE:HG21	1.73	0.69
1:A:54:PRO:HD3	1:A:188:TRP:CZ3	2.28	0.69
1:B:159:LEU:HD23	1:B:182:ILE:HG23	1.75	0.67
1:B:198:GLU:OE1	1:A:66:LYS:NZ	2.28	0.64
1:A:54:PRO:HD3	1:A:188:TRP:CH2	2.33	0.64
1:A:153:VAL:HG23	1:A:218:TRP:CZ3	2.33	0.64
1:B:185:ARG:HG3	1:B:185:ARG:HH11	1.66	0.61
1:A:6:LEU:HB2	1:A:7:PRO:HD3	1.84	0.59
1:B:44:LEU:O	1:B:212:PRO:HB3	2.02	0.59
1:B:221:GLU:HG3	1:B:222:HIS:ND1	2.17	0.59
1:A:193:ASP:HB3	3:A:315:BU1:H12	1.86	0.57
1:B:107:LEU:HD11	1:A:107:LEU:CD1	2.34	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:202:LEU:HB2	3:A:315:BU1:H41	1.86	0.56
1:B:107:LEU:CD1	1:A:107:LEU:CD1	2.83	0.56
1:B:175:LYS:HD2	1:B:185:ARG:HD3	1.87	0.56
1:B:131:PHE:HB2	1:B:132:PRO:HD3	1.88	0.56
1:A:109:ARG:HB2	1:A:146:PRO:HB3	1.87	0.56
1:A:131:PHE:HB2	1:A:132:PRO:HD3	1.88	0.56
1:B:40:HIS:HD2	1:B:116:PRO:HG2	1.71	0.55
1:B:37:LEU:CD2	1:B:39:VAL:HG23	2.37	0.55
1:A:140:ARG:HD2	5:A:337:HOH:O	2.07	0.54
1:A:35:CYS:HG	1:A:198:GLU:HA	1.73	0.54
1:A:192:THR:HB	3:A:315:BU1:C4	2.38	0.54
1:B:128:ARG:NH1	2:D:1:DG:N7	2.57	0.53
1:B:114:ARG:NH2	1:A:99:ARG:O	2.41	0.53
1:A:202:LEU:CD1	3:A:315:BU1:H21	2.36	0.52
1:A:58:PRO:HD3	1:A:187:TYR:CE2	2.45	0.52
1:A:182:ILE:HG22	1:A:186:LEU:HD21	1.92	0.51
1:B:131:PHE:CE1	1:B:152:CYS:HB2	2.46	0.51
1:B:183:TYR:CD2	1:B:203:THR:HG22	2.45	0.51
1:A:212:PRO:O	1:A:216:LEU:CD2	2.58	0.51
1:A:42:ARG:HA	1:A:45:GLU:HG2	1.91	0.51
1:B:44:LEU:O	1:B:212:PRO:CB	2.59	0.50
1:B:91:GLU:HG3	5:B:377:HOH:O	2.12	0.50
1:B:24:LEU:HB3	1:B:25:PRO:HD2	1.92	0.50
1:B:95:GLN:OE1	1:A:43:ALA:HB1	2.13	0.49
1:B:63:VAL:HB	1:A:95:GLN:OE1	2.12	0.49
1:A:159:LEU:HD12	1:A:207:ILE:CG2	2.43	0.48
1:B:153:VAL:CG1	1:B:223:ALA:HA	2.44	0.48
1:A:153:VAL:HG23	1:A:218:TRP:HZ3	1.77	0.48
1:A:51:GLN:HB2	1:A:52:GLY:HA3	1.94	0.48
1:A:40:HIS:CE1	1:A:42:ARG:HG2	2.49	0.47
1:B:37:LEU:C	1:B:37:LEU:HD21	2.35	0.47
1:B:107:LEU:CD1	1:A:107:LEU:HD13	2.45	0.47
1:A:225:PRO:HB2	1:A:228:THR:HG23	1.96	0.47
1:B:34:LEU:HD23	1:B:34:LEU:C	2.34	0.47
1:A:39:VAL:HG12	1:A:61:PRO:HG2	1.97	0.46
1:B:224:ARG:HH21	1:B:229:VAL:HG22	1.80	0.46
1:B:152:CYS:O	1:B:153:VAL:HG11	2.15	0.45
1:A:199:GLY:HA2	3:A:315:BU1:H12	1.97	0.45
1:B:98:GLN:O	1:A:114:ARG:NH2	2.48	0.45
1:A:192:THR:HB	3:A:315:BU1:H31	1.95	0.45
1:B:40:HIS:CD2	1:B:116:PRO:HG2	2.52	0.44
1:B:193:ASP:OD1	1:B:196:THR:OG1	2.30	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:54:PRO:HD3	1:A:188:TRP:CE3	2.52	0.44
1:B:131:PHE:CZ	1:B:152:CYS:HB2	2.53	0.43
1:B:153:VAL:HG11	1:B:223:ALA:HA	2.00	0.43
1:B:30:GLU:OE1	1:B:74:GLY:N	2.46	0.43
1:B:185:ARG:NH1	1:B:185:ARG:HG3	2.32	0.43
1:B:24:LEU:HB3	1:B:25:PRO:CD	2.49	0.43
1:B:29:PRO:O	1:B:233:TYR:OH	2.37	0.43
1:A:159:LEU:CD1	1:A:207:ILE:CG2	2.97	0.43
1:B:163:GLU:O	1:B:166:SER:N	2.48	0.43
1:B:186:LEU:HD12	1:B:210:TRP:CG	2.55	0.42
1:B:182:ILE:O	1:B:186:LEU:HG	2.20	0.42
1:A:218:TRP:CD1	1:A:222:HIS:CE1	3.07	0.42
1:B:138:LEU:O	1:B:141:LEU:HB2	2.19	0.42
1:B:129:TYR:O	1:B:132:PRO:HD2	2.20	0.42
1:B:14:LEU:HD13	1:B:216:LEU:HD21	2.01	0.42
1:A:13:THR:OG1	1:A:116:PRO:HD2	2.20	0.42
1:A:20[B]:GLU:OE1	1:A:85:THR:HB	2.19	0.42
1:A:217:GLN:HA	1:A:217:GLN:OE1	2.19	0.41
1:B:6:LEU:HG	1:B:9:GLY:O	2.21	0.41
1:A:193:ASP:CB	3:A:315:BU1:H32	2.51	0.41
1:A:159:LEU:HD12	1:A:207:ILE:HG23	2.02	0.41
1:A:40:HIS:ND1	1:A:42:ARG:HG2	2.35	0.41
1:B:216:LEU:HA	1:B:216:LEU:HD23	1.96	0.41
1:A:164:GLN:CD	1:A:164:GLN:N	2.74	0.40
1:B:6:LEU:HD22	1:B:9:GLY:C	2.40	0.40
2:D:3:DC:H5	5:D:177:HOH:O	2.00	0.40
1:A:193:ASP:HB3	3:A:315:BU1:H32	2.03	0.40
1:A:44:LEU:CB	1:A:216:LEU:HD23	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/314 (67%)	205 (98%)	5 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	215/314 (68%)	213 (99%)	2 (1%)	0	100	100
All	All	425/628 (68%)	418 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/263 (70%)	171 (92%)	14 (8%)	19	21
1	B	188/263 (72%)	177 (94%)	11 (6%)	28	33
All	All	373/526 (71%)	348 (93%)	25 (7%)	23	27

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	6	LEU
1	B	34	LEU
1	B	44	LEU
1	B	83	GLU
1	B	113	GLN
1	B	114	ARG
1	B	141	LEU
1	B	143	THR
1	B	159	LEU
1	B	213	GLN
1	B	217	GLN
1	A	6	LEU
1	A	36	LEU
1	A	42	ARG
1	A	51	GLN
1	A	59	ARG
1	A	99	ARG
1	A	112	LEU
1	A	114	ARG
1	A	153	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	159	LEU
1	A	175	LYS
1	A	186	LEU
1	A	207	ILE
1	A	217	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	189	GLN
1	A	51	GLN
1	A	103	ASN
1	A	117	GLN
1	A	125	ASN
1	A	135	GLN
1	A	209	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BU1	A	315	-	5,5,5	0.33	0	4,4,4	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BU1	A	315	-	-	0/3/3/3	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/314 (68%)	0.91	37 (17%) 2 3	20, 38, 66, 99	0
1	B	220/314 (70%)	0.90	38 (17%) 2 3	19, 42, 72, 88	0
2	C	4/4 (100%)	0.55	1 (25%) 1 2	28, 36, 43, 49	0
2	D	4/4 (100%)	0.33	0 100 100	36, 37, 37, 50	0
All	All	443/636 (69%)	0.90	76 (17%) 2 3	19, 40, 69, 99	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	165	ALA	6.3
1	B	94	VAL	5.7
1	B	164	GLN	5.6
1	A	142	SER	5.3
1	A	188	TRP	5.1
1	A	143	THR	4.4
1	B	229	VAL	4.4
1	A	79	PRO	4.3
1	A	217	GLN	4.2
1	B	208	CYS	4.1
1	A	208	CYS	4.1
1	B	228	THR	4.1
1	B	142	SER	4.0
1	B	143	THR	4.0
1	A	17	LEU	3.9
1	B	106	ILE	3.8
1	A	205	LEU	3.6
1	A	207	ILE	3.6
1	A	174	ARG	3.5
1	A	204	LEU	3.4
1	B	49	ILE	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	6	LEU	3.3
1	A	175	LYS	3.3
1	B	148	ASP	3.3
1	A	162	LEU	3.1
1	B	134	LEU	3.0
1	A	141	LEU	3.0
1	B	59	ARG	3.0
1	A	155	SER	3.0
1	B	219	VAL	2.9
2	C	1	DG	2.8
1	B	141	LEU	2.8
1	B	84	ILE	2.8
1	A	16	PHE	2.8
1	B	166	SER	2.8
1	A	57	VAL	2.8
1	A	84	ILE	2.8
1	A	53	HIS	2.7
1	A	94	VAL	2.7
1	B	227	SER	2.6
1	A	124	HIS	2.5
1	A	144	PRO	2.5
1	A	214	ALA	2.5
1	A	184	THR	2.5
1	B	44	LEU	2.5
1	B	220	ASP	2.5
1	A	218	TRP	2.4
1	B	16	PHE	2.4
1	A	26	SER	2.4
1	A	229	VAL	2.4
1	A	80	GLY	2.3
1	B	174	ARG	2.3
1	B	217	GLN	2.3
1	B	118	PRO	2.3
1	B	119	CYS	2.3
1	B	144	PRO	2.3
1	A	122	VAL	2.2
1	B	34	LEU	2.2
1	B	155	SER	2.2
1	A	37	LEU	2.2
1	A	54	PRO	2.2
1	A	56	PRO	2.2
1	B	80	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	46	ASN	2.1
1	B	50	SER	2.1
1	B	184	THR	2.1
1	B	230	LYS	2.1
1	A	14	LEU	2.1
1	B	177	TYR	2.1
1	B	214	ALA	2.1
1	B	188	TRP	2.1
1	A	34	LEU	2.1
1	B	221	GLU	2.1
1	A	19	LEU	2.0
1	A	52	GLY	2.0
1	B	122	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	BU1	A	315	6/6	0.44	9.02	30,39,42,50	0
4	CA	C	5	1/1	0.06	-2.39	47,47,47,47	0
4	CA	D	5	1/1	0.05	-3.71	35,35,35,35	0

6.5 Other polymers ⓘ

There are no such residues in this entry.