



wwPDB X-ray Structure Validation Summary Report i

Sep 25, 2014 – 07:42 PM EDT

PDB ID : 4U4J
Title : Crystal structure of Salmonella alpha-2-macroglobulinmutant Y1175G
Authors : Wong, S.G.; Dessen, A.
Deposited on : 2014-07-23
Resolution : 2.90 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.16 November 2013
Xtriage (Phenix)	:	dev-1439
EDS	:	stable23828
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable23828

1 Overall quality at a glance


The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1053 (2.90-2.90)
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)
RSRZ outliers	66119	1054 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	1647	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 23627 atoms, of which 11754 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative inner membrane lipoprotein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	1538	Total	C	H	N	O	S	0	0	0
			23627	7480	11754	2091	2279	23			


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	initiating methionine	UNP Q8ZN46
A	-1	GLY	-	expression tag	UNP Q8ZN46
A	0	SER	-	expression tag	UNP Q8ZN46
A	1	SER	-	expression tag	UNP Q8ZN46
A	2	HIS	-	expression tag	UNP Q8ZN46
A	3	HIS	-	expression tag	UNP Q8ZN46
A	4	HIS	-	expression tag	UNP Q8ZN46
A	5	HIS	-	expression tag	UNP Q8ZN46
A	6	HIS	-	expression tag	UNP Q8ZN46
A	7	HIS	-	expression tag	UNP Q8ZN46
A	8	SER	-	expression tag	UNP Q8ZN46
A	9	SER	-	expression tag	UNP Q8ZN46
A	10	GLY	-	expression tag	UNP Q8ZN46
A	11	LEU	-	expression tag	UNP Q8ZN46
A	12	VAL	-	expression tag	UNP Q8ZN46
A	13	PRO	-	expression tag	UNP Q8ZN46
A	14	ARG	-	expression tag	UNP Q8ZN46
A	15	GLY	-	expression tag	UNP Q8ZN46
A	16	SER	-	expression tag	UNP Q8ZN46
A	17	HIS	-	expression tag	UNP Q8ZN46
A	18	MET	-	expression tag	UNP Q8ZN46
A	98	ALA	LYS	engineered mutation	UNP Q8ZN46
A	99	ALA	LYS	engineered mutation	UNP Q8ZN46
A	1175	GLY	TYR	engineered mutation	UNP Q8ZN46

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative inner membrane lipoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	254.24Å 82.16Å 98.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.99 – 2.90 58.99 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (58.99-2.90) 99.9 (58.99-2.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 2.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.244 , 0.284 0.244 , 0.284	Depositor DCC
R_{free} test set	2338 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	87.1	Xtriage
Anisotropy	0.461	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 46790 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	23627	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/12113	0.43	1/16503 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	316	LEU	CA-CB-CG	7.46	132.47	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	252	LYS	Peptide
1	A	808	LYS	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11873	11754	18	59	0
All	All	11873	11754	18	59	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

The worst 5 of 59 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:87:GLN:N	1:A:87:GLN:OE1	2.19	0.76
1:A:1366:GLU:O	1:A:1368:GLN:N	2.19	0.75
1:A:1368:GLN:OE1	1:A:1369:TRP:N	2.30	0.65
1:A:463:LEU:O	1:A:465:ARG:NH1	2.34	0.60
1:A:317:ASN:O	1:A:319:LYS:N	2.33	0.60

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1526/1647 (93%)	1375 (90%)	106 (7%)	45 (3%)	7	28

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	321	GLN
1	A	556	ASP
1	A	587	SER
1	A	651	LYS
1	A	758	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1259/1347 (94%)	1227 (98%)	32 (2%)	60 91

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	334	VAL
1	A	555	LEU
1	A	1368	GLN
1	A	439	ARG
1	A	586	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	758	ASN
1	A	956	HIS
1	A	1091	GLN
1	A	1328	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1538/1647 (93%)	0.67	130 (8%) 11 13	51, 90, 137, 213	0

The worst 5 of 130 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	928	ILE	8.8
1	A	85	PRO	7.2
1	A	588	GLY	6.6
1	A	57	LYS	6.1
1	A	929	GLU	6.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.