



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 10:17 AM GMT

PDB ID : 3U7G
Title : Crystal structure of mPNKP catalytic fragment (D170A) bound to single-stranded DNA (TCCTAp)
Authors : Coquelle, N.; Havali, Z.; Bernstein, N.; Green, R.; Glover, J.N.M.
Deposited on : 2011-10-13
Resolution : 2.10 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance



The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	379	
2	X	5	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	GOL	A	1	-	X
4	GOL	A	526	-	X
4	GOL	A	527	-	X
5	MG	A	529	-	X

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 6141 atoms, of which 2946 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional polynucleotide phosphatase/kinase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	377	Total	C	H	N	O	S	Se	0	0	0
			5847	1878	2922	508	527	7	5			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	170	ALA	ASP	ENGINEERED MUTATION	UNP Q9JLV6

- Molecule 2 is a DNA chain called DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	2	Total	C	N	O	P	0	0	0
			41	20	7	12	2			

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	X	1	Total	O	P	0	0
			4	3	1		
3	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	A	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	196	Total	O	0	0
			196	196		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	X	5	Total	O	0	0
			5	5		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional polynucleotide phosphatase/kinase



- Molecule 2: DNA



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	77.16Å 43.03Å 127.27Å 90.00° 107.12° 90.00°	Depositor
Resolution (Å)	40.54 – 2.10 40.54 – 2.10	Depositor EDS
% Data completeness (in resolution range)	90.8 (40.54-2.10) 90.8 (40.54-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 2.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, R_{free}	0.207 , 0.260 0.203 , 0.255	Depositor DCC
R_{free} test set	1082 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.668	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 35.1	EDS
Estimated twinning fraction	0.033 for h,-k,-h-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 21483 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6141	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2996	0.53	0/4057
2	X	1.09	0/45	1.25	0/67
All	All	0.41	0/3041	0.55	0/4124

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2925	2922	0	51	0
2	X	41	0	23	1	0
3	A	5	0	0	0	0
3	X	4	0	0	0	0
4	A	18	24	0	3	0
5	A	1	0	0	0	0
6	A	196	0	0	12	0
6	X	5	0	0	0	0
All	All	3195	2946	23	51	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (51) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:492:GLU:OE2	6:A:89:HOH:O	1.90	0.90
1:A:422:ASN:ND2	6:A:101:HOH:O	2.05	0.89
1:A:354:PRO:O	6:A:80:HOH:O	2.01	0.78
1:A:357:SER:OG	6:A:93:HOH:O	2.08	0.71
1:A:492:GLU:O	6:A:103:HOH:O	2.10	0.70
1:A:304:ASP:O	6:A:73:HOH:O	2.14	0.64
1:A:430:ARG:NE	6:A:89:HOH:O	2.33	0.62
1:A:447:PHE:CE1	1:A:498:LEU:HD12	2.38	0.59
1:A:427:VAL:HA	6:A:89:HOH:O	2.03	0.58
1:A:410:ALA:O	1:A:413:ARG:HG2	2.03	0.57
1:A:281:GLU:HG2	6:A:539:HOH:O	2.05	0.56
1:A:408:CYS:HA	1:A:418:VAL:HG21	1.87	0.56
1:A:235:VAL:O	1:A:239:LEU:HG	2.06	0.55
1:A:162:PRO:HB2	1:A:276:ILE:HD12	1.89	0.55
1:A:286:VAL:HA	1:A:321:ALA:O	2.07	0.53
1:A:359:LEU:HG	1:A:360:LEU:HD12	1.90	0.52
1:A:149:LEU:N	1:A:149:LEU:CD1	2.73	0.51
1:A:360:LEU:HD21	1:A:388:ALA:CB	2.41	0.51
1:A:418:VAL:HB	6:A:82:HOH:O	2.10	0.50
1:A:394:ASN:OD1	1:A:395:ARG:N	2.44	0.50
1:A:359:LEU:HD21	1:A:384:HIS:HB3	1.95	0.49
1:A:160:VAL:HA	4:A:526:GOL:O3	2.12	0.49
1:A:236:GLU:O	1:A:240:GLU:HG2	2.13	0.49
1:A:430:ARG:HB2	6:A:89:HOH:O	2.12	0.48
1:A:211:LYS:HA	4:A:526:GOL:H31	1.96	0.47
1:A:360:LEU:HD21	1:A:388:ALA:HB3	1.97	0.47
1:A:503:ARG:HD3	6:A:68:HOH:O	2.15	0.47
1:A:424:ASN:N	1:A:425:PRO:HD3	2.30	0.47
1:A:364:PRO:HA	1:A:417:ARG:HG2	1.98	0.46
1:A:434:ILE:HD11	1:A:494:PHE:CZ	2.50	0.46
1:A:489:THR:O	1:A:492:GLU:CG	2.65	0.45
1:A:232:LYS:O	1:A:236:GLU:HG3	2.16	0.45
1:A:168:ALA:HA	1:A:213:VAL:O	2.17	0.45
1:A:295:ASN:HA	1:A:300:ARG:O	2.17	0.45
1:A:489:THR:O	1:A:492:GLU:HG2	2.17	0.45
1:A:453:ILE:O	1:A:457:ARG:HG3	2.17	0.44
1:A:322:THR:HB	1:A:323:PRO:HD2	1.98	0.44
1:A:459:ASN:O	1:A:463:ARG:HG3	2.17	0.44
1:A:475:ASP:O	1:A:478:MSE:N	2.51	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:359:LEU:HG	1:A:360:LEU:CD1	2.48	0.43
1:A:427:VAL:N	1:A:428:PRO:CD	2.82	0.43
1:A:217:ASN:O	2:X:402:DA:H5''	2.19	0.42
1:A:373:PRO:HB2	1:A:473:VAL:HG11	2.02	0.42
1:A:488:PRO:HA	1:A:492:GLU:OE1	2.20	0.42
1:A:429:SER:O	1:A:432:ARG:HB3	2.19	0.41
1:A:365:GLU:OE2	1:A:445:ARG:NH2	2.51	0.41
1:A:149:LEU:HD12	1:A:149:LEU:N	2.35	0.41
1:A:411:ALA:HB3	1:A:418:VAL:CG2	2.50	0.41
1:A:342:PRO:HG3	4:A:1:GOL:H11	2.03	0.41
1:A:405:VAL:HG21	1:A:432:ARG:HD2	2.03	0.40
1:A:184:PHE:HA	1:A:185:PRO:HD3	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	373/379 (98%)	360 (96%)	11 (3%)	2 (0%)	38	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	398	LEU
1	A	298	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	311/311 (100%)	307 (99%)	4 (1%)	80 85

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	149	LEU
1	A	383	GLU
1	A	492	GLU
1	A	509	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	271	GLN
1	A	422	ASN
1	A	513	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	1	-	5,5,5	0.48	0	5,5,5	0.40	0
4	GOL	A	526	-	5,5,5	0.32	0	5,5,5	0.37	0
4	GOL	A	527	-	5,5,5	0.24	0	5,5,5	0.40	0
3	PO4	A	528	-	4,4,4	0.33	0	6,6,6	0.31	0
3	PO4	X	523	2,5	1,3,4	10.32	1 (100%)	0,3,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1	-	-	0/4/4/4	0/0/0/0
4	GOL	A	526	-	-	0/4/4/4	0/0/0/0
4	GOL	A	527	-	-	0/4/4/4	0/0/0/0
3	PO4	A	528	-	-	0/0/0/0	0/0/0/0
3	PO4	X	523	2,5	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	X	523	PO4	P-O1	10.32	1.58	1.46

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	377/379 (99%)	0.57	42 (11%) 6 7	8, 27, 66, 101	0
2	X	2/5 (40%)	-0.03	0 100 100	20, 20, 20, 42	0
All	All	379/384 (98%)	0.57	42 (11%) 6 7	8, 27, 66, 101	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	434	ILE	6.2
1	A	442	VAL	5.9
1	A	405	VAL	5.8
1	A	418	VAL	5.6
1	A	401	TRP	5.6
1	A	366	VAL	5.2
1	A	420	ILE	4.3
1	A	412	LEU	4.1
1	A	419	VAL	4.0
1	A	393	VAL	3.8
1	A	367	VAL	3.8
1	A	368	VAL	3.7
1	A	415	GLY	3.6
1	A	397	THR	3.3
1	A	144	LEU	3.2
1	A	411	ALA	3.0
1	A	413	ARG	2.9
1	A	416	LYS	2.8
1	A	427	VAL	2.8
1	A	497	ILE	2.8
1	A	431	ALA	2.8
1	A	443	PRO	2.8
1	A	422	ASN	2.8
1	A	391	VAL	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	381	ILE	2.7
1	A	473	VAL	2.7
1	A	441	GLY	2.6
1	A	370	VAL	2.6
1	A	404	CYS	2.6
1	A	360	LEU	2.5
1	A	385	LEU	2.5
1	A	505	GLN	2.4
1	A	432	ARG	2.3
1	A	477	VAL	2.3
1	A	504	LEU	2.2
1	A	398	LEU	2.2
1	A	392	HIS	2.2
1	A	396	ASP	2.2
1	A	493	GLY	2.1
1	A	345	ILE	2.0
1	A	348	ALA	2.0
1	A	438	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	MG	A	529	1/1	0.28	12.54	30,30,30,30	1
4	GOL	A	1	6/6	0.23	4.05	21,27,32,36	0
4	GOL	A	526	6/6	0.21	3.59	47,57,59,60	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	GOL	A	527	6/6	0.18	2.07	38,47,55,56	0
3	PO4	X	523	4/5	0.10	-0.27	12,14,14,15	0
3	PO4	A	528	5/5	0.10	-0.87	22,24,27,31	0

6.5 Other polymers ⓘ

There are no such residues in this entry.