



wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 09:50 AM GMT

PDB ID : 3UBJ
Title : Influenza hemagglutinin from the 2009 pandemic in complex with ligand LSTa
Authors : Xu, R.; Wilson, I.A.
Deposited on : 2011-10-24
Resolution : 2.25 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1108 (2.28-2.24)
Clashscore	79885	1326 (2.28-2.24)
Ramachandran outliers	78287	1291 (2.28-2.24)
Sidechain outliers	78261	1291 (2.28-2.24)
RSRZ outliers	66119	1110 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	329	
1	C	329	
1	E	329	
1	G	329	
1	I	329	
1	K	329	
2	B	177	
2	D	177	
2	F	177	
2	H	177	
2	J	177	
2	L	177	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 24423 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2522	1595	432	482	13			
1	C	320	Total	C	N	O	S	0	0	0
			2501	1582	429	477	13			
1	E	319	Total	C	N	O	S	0	0	0
			2492	1578	428	473	13			
1	G	323	Total	C	N	O	S	0	0	0
			2522	1595	432	482	13			
1	I	320	Total	C	N	O	S	0	0	0
			2501	1582	429	477	13			
1	K	320	Total	C	N	O	S	0	0	0
			2504	1585	429	477	13			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
A	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
A	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
A	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1
C	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
C	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
C	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
C	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1
E	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
E	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
E	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
E	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1
G	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
G	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
G	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
G	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1
I	9	PRO	-	EXPRESSION TAG	UNP C3W5S1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
I	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
I	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1
K	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
K	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
K	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
K	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1

- Molecule 2 is a protein called Hemagglutinin HA2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	169	Total	C	N	O	S	0	0	0
			1360	855	229	270	6			
2	D	170	Total	C	N	O	S	0	0	0
			1371	861	233	271	6			
2	F	170	Total	C	N	O	S	0	0	0
			1371	861	233	271	6			
2	H	172	Total	C	N	O	S	0	0	0
			1389	871	235	277	6			
2	J	170	Total	C	N	O	S	0	0	0
			1371	861	233	271	6			
2	L	169	Total	C	N	O	S	0	0	0
			1360	855	229	270	6			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	175	SER	-	EXPRESSION TAG	UNP C3W5S1
B	176	GLY	-	EXPRESSION TAG	UNP C3W5S1
B	177	ARG	-	EXPRESSION TAG	UNP C3W5S1
D	175	SER	-	EXPRESSION TAG	UNP C3W5S1
D	176	GLY	-	EXPRESSION TAG	UNP C3W5S1
D	177	ARG	-	EXPRESSION TAG	UNP C3W5S1
F	175	SER	-	EXPRESSION TAG	UNP C3W5S1
F	176	GLY	-	EXPRESSION TAG	UNP C3W5S1
F	177	ARG	-	EXPRESSION TAG	UNP C3W5S1
H	175	SER	-	EXPRESSION TAG	UNP C3W5S1
H	176	GLY	-	EXPRESSION TAG	UNP C3W5S1
H	177	ARG	-	EXPRESSION TAG	UNP C3W5S1
J	175	SER	-	EXPRESSION TAG	UNP C3W5S1
J	176	GLY	-	EXPRESSION TAG	UNP C3W5S1
J	177	ARG	-	EXPRESSION TAG	UNP C3W5S1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	175	SER	-	EXPRESSION TAG	UNP C3W5S1
L	176	GLY	-	EXPRESSION TAG	UNP C3W5S1
L	177	ARG	-	EXPRESSION TAG	UNP C3W5S1


- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	E	2	Total	C	N	O	0	0
			28	16	2	10		
3	K	2	Total	C	N	O	0	0
			28	16	2	10		

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
A	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
A	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
A	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1
E	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
E	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
E	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
E	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1
K	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
K	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
K	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
K	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	E	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	I	1	Total	C	N	O	0	0
			14	8	1	5		
4	I	1	Total	C	N	O	0	0
			14	8	1	5		
4	I	1	Total	C	N	O	0	0
			14	8	1	5		
4	K	1	Total	C	N	O	0	0
			14	8	1	5		
4	K	1	Total	C	N	O	0	0
			14	8	1	5		
4	K	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	2	Total	C	N	O	0	0
			31	17	1	13		
5	G	2	Total	C	N	O	0	0
			31	17	1	13		
5	I	2	Total	C	N	O	0	0
			31	17	1	13		
5	K	2	Total	C	N	O	0	0
			31	17	1	13		

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
C	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
C	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
C	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1
G	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
G	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
G	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
G	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1
I	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
I	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
I	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
I	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1
K	9	PRO	-	EXPRESSION TAG	UNP C3W5S1
K	10	GLY	-	EXPRESSION TAG	UNP C3W5S1
K	205	CYS	GLY	ENGINEERED MUTATION	UNP C3W5S1
K	220	CYS	ARG	ENGINEERED MUTATION	UNP C3W5S1

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	98	Total	O	0	0
			98	98		
6	B	47	Total	O	0	0
			47	47		
6	C	71	Total	O	0	0
			71	71		
6	D	52	Total	O	0	0
			52	52		
6	E	91	Total	O	0	0
			91	91		
6	F	40	Total	O	0	0
			40	40		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	G	84	Total 84	O 84	0	0
6	H	54	Total 54	O 54	0	0
6	I	104	Total 104	O 104	0	0
6	J	43	Total 43	O 43	0	0
6	K	65	Total 65	O 65	0	0
6	L	34	Total 34	O 34	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Hemagglutinin HA1

Chain A:




• Molecule 1: Hemagglutinin HA1

Chain C:




• Molecule 1: Hemagglutinin HA1


Chain E:




- Molecule 1: Hemagglutinin HA1




- Molecule 1: Hemagglutinin HA1



- Molecule 1: Hemagglutinin HA1




- Molecule 2: Hemagglutinin HA2




- Molecule 2: Hemagglutinin HA2

Chain D:




• Molecule 2: Hemagglutinin HA2

Chain F:




• Molecule 2: Hemagglutinin HA2

Chain H:




• Molecule 2: Hemagglutinin HA2

Chain J:



• Molecule 2: Hemagglutinin HA2

Chain L:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	66.85Å 116.09Å 118.40Å 61.05° 77.00° 80.39°	Depositor
Resolution (Å)	49.32 – 2.25 49.31 – 2.24	Depositor EDS
% Data completeness (in resolution range)	91.0 (49.32-2.25) 90.4 (49.31-2.24)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.25Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.202 , 0.252 0.199 , 0.250	Depositor DCC
R_{free} test set	6543 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 131422 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	24423	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GAL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.25	0/2586	0.45	0/3516
1	C	0.24	0/2564	0.43	0/3484
1	E	0.25	0/2556	0.46	0/3473
1	G	0.25	0/2586	0.43	0/3516
1	I	0.25	0/2564	0.45	0/3484
1	K	0.24	0/2567	0.43	0/3488
2	B	0.24	0/1388	0.38	0/1871
2	D	0.25	0/1399	0.40	0/1885
2	F	0.25	0/1399	0.39	0/1885
2	H	0.24	0/1417	0.39	0/1909
2	J	0.24	0/1399	0.38	0/1885
2	L	0.24	0/1388	0.37	0/1871
All	All	0.25	0/23813	0.42	0/32267

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2522	0	2466	57	0
1	C	2501	0	2444	50	0
1	E	2492	0	2437	64	0
1	G	2522	0	2465	55	0
1	I	2501	0	2441	45	0
1	K	2504	0	2446	62	0
2	B	1360	0	1284	37	0
2	D	1371	0	1296	33	0
2	F	1371	0	1297	31	0
2	H	1389	0	1309	34	0
2	J	1371	0	1297	35	0
2	L	1360	0	1284	26	0
3	A	28	0	25	0	0
3	E	28	0	25	1	0
3	K	28	0	25	1	0
4	A	14	0	13	0	0
4	D	14	0	13	0	0
4	E	14	0	13	0	0
4	G	42	0	39	0	0
4	I	42	0	39	1	0
4	K	42	0	39	1	0
5	C	31	0	26	1	0
5	G	31	0	26	1	0
5	I	31	0	26	3	0
5	K	31	0	26	2	0
6	A	98	0	0	2	0
6	B	47	0	0	1	0
6	C	71	0	0	0	0
6	D	52	0	0	0	0
6	E	91	0	0	1	0
6	F	40	0	0	2	0
6	G	84	0	0	0	0
6	H	54	0	0	1	0
6	I	104	0	0	1	0
6	J	43	0	0	2	0
6	K	65	0	0	0	0
6	L	34	0	0	1	0
All	All	24423	0	22801	464	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

The worst 5 of 464 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:283:THR:HG22	1:C:285:LYS:H	1.19	1.08
2:H:97:GLU:HG2	2:L:58:LYS:HD2	1.44	0.96
1:A:79:LEU:HA	1:A:81:THR:N	1.84	0.93
1:I:283:THR:HG22	1:I:285:LYS:H	1.35	0.91
1:A:283:THR:HG22	1:A:285:LYS:H	1.39	0.86

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/329 (98%)	301 (94%)	15 (5%)	5 (2%)	14	8
1	C	316/329 (96%)	299 (95%)	15 (5%)	2 (1%)	33	33
1	E	315/329 (96%)	299 (95%)	14 (4%)	2 (1%)	33	33
1	G	321/329 (98%)	301 (94%)	19 (6%)	1 (0%)	50	54
1	I	316/329 (96%)	298 (94%)	16 (5%)	2 (1%)	33	33
1	K	316/329 (96%)	298 (94%)	17 (5%)	1 (0%)	50	54
2	B	167/177 (94%)	159 (95%)	7 (4%)	1 (1%)	33	33
2	D	168/177 (95%)	154 (92%)	11 (6%)	3 (2%)	13	6
2	F	168/177 (95%)	156 (93%)	11 (6%)	1 (1%)	33	33
2	H	170/177 (96%)	164 (96%)	6 (4%)	0	100	100
2	J	168/177 (95%)	156 (93%)	7 (4%)	5 (3%)	7	2
2	L	167/177 (94%)	156 (93%)	10 (6%)	1 (1%)	33	33
All	All	2913/3036 (96%)	2741 (94%)	148 (5%)	24 (1%)	27	24

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	127	LYS
1	C	94	ASN
1	A	81	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	265	GLY
1	C	264	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/290 (98%)	279 (98%)	6 (2%)	66	75
1	C	282/290 (97%)	270 (96%)	12 (4%)	40	45
1	E	281/290 (97%)	274 (98%)	7 (2%)	60	70
1	G	285/290 (98%)	279 (98%)	6 (2%)	66	75
1	I	282/290 (97%)	273 (97%)	9 (3%)	51	60
1	K	283/290 (98%)	272 (96%)	11 (4%)	43	51
2	B	145/152 (95%)	138 (95%)	7 (5%)	35	39
2	D	146/152 (96%)	140 (96%)	6 (4%)	41	47
2	F	146/152 (96%)	141 (97%)	5 (3%)	49	57
2	H	148/152 (97%)	144 (97%)	4 (3%)	57	67
2	J	146/152 (96%)	142 (97%)	4 (3%)	57	67
2	L	145/152 (95%)	141 (97%)	4 (3%)	56	65
All	All	2574/2652 (97%)	2493 (97%)	81 (3%)	52	61

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	295	GLN
1	G	253	VAL
1	K	133(A)	LYS
1	E	320	LEU
2	F	163	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	27	GLN
2	H	79	ASN
1	K	18	HIS
2	F	95	ASN
2	J	27	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	330	1,3	12,14,15	0.70	1 (8%)	15,19,21	1.00	1 (6%)
3	NAG	A	331	3	12,14,15	0.70	1 (8%)	15,19,21	0.95	1 (6%)
5	SIA	C	330	5	20,20,21	3.80	1 (5%)	23,28,31	2.59	5 (21%)
5	GAL	C	331	5	10,11,12	0.83	1 (10%)	11,15,17	1.07	1 (9%)
3	NAG	E	331	1,3	12,14,15	0.71	1 (8%)	15,19,21	0.90	0
3	NAG	E	332	3	12,14,15	0.65	0	15,19,21	1.03	2 (13%)
5	SIA	G	401	5	20,20,21	3.84	1 (5%)	23,28,31	2.64	6 (26%)
5	GAL	G	402	5	10,11,12	0.78	0	11,15,17	1.32	1 (9%)
5	SIA	I	501	5	20,20,21	3.72	1 (5%)	23,28,31	2.63	4 (17%)
5	GAL	I	502	5	10,11,12	0.83	1 (10%)	11,15,17	0.87	0
5	SIA	K	601	5	20,20,21	3.81	1 (5%)	23,28,31	2.50	4 (17%)
5	GAL	K	602	5	10,11,12	0.80	1 (10%)	11,15,17	0.87	1 (9%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	K	631	1,3	12,14,15	0.64	0	15,19,21	0.91	1 (6%)
3	NAG	K	632	3	12,14,15	0.61	0	15,19,21	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	330	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	331	3	-	0/6/23/26	0/1/1/1
5	SIA	C	330	5	-	0/15/34/38	0/1/1/1
5	GAL	C	331	5	-	0/2/19/22	0/1/1/1
3	NAG	E	331	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	332	3	-	0/6/23/26	0/1/1/1
5	SIA	G	401	5	-	0/15/34/38	0/1/1/1
5	GAL	G	402	5	-	0/2/19/22	0/1/1/1
5	SIA	I	501	5	-	0/15/34/38	0/1/1/1
5	GAL	I	502	5	-	0/2/19/22	0/1/1/1
5	SIA	K	601	5	-	0/15/34/38	0/1/1/1
5	GAL	K	602	5	-	0/2/19/22	0/1/1/1
3	NAG	K	631	1,3	-	0/6/23/26	0/1/1/1
3	NAG	K	632	3	-	0/6/23/26	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	401	SIA	O6-C2	16.93	1.43	1.28
5	K	601	SIA	O6-C2	16.79	1.43	1.28
5	C	330	SIA	O6-C2	16.78	1.43	1.28
5	I	501	SIA	O6-C2	16.39	1.43	1.28
3	E	331	NAG	O5-C5	-2.15	1.41	1.45

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	I	501	SIA	O6-C2-C3	-11.29	110.17	124.91
5	K	601	SIA	O6-C2-C3	-10.82	110.77	124.91
5	G	401	SIA	O6-C2-C3	-10.81	110.79	124.91
5	C	330	SIA	O6-C2-C3	-10.68	110.96	124.91
5	G	401	SIA	O6-C6-C5	3.80	115.05	110.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAG	A	332	1	12,14,15	0.67	0	15,19,21	0.83	0
4	NAG	D	261	2	12,14,15	0.63	0	15,19,21	1.04	1 (6%)
4	NAG	E	341	1	12,14,15	0.64	0	15,19,21	0.96	0
4	NAG	G	411	1	12,14,15	0.63	0	15,19,21	0.92	1 (6%)
4	NAG	G	431	1	12,14,15	0.64	0	15,19,21	0.91	1 (6%)
4	NAG	G	451	1	12,14,15	0.65	0	15,19,21	0.94	0
4	NAG	I	521	1	12,14,15	0.64	0	15,19,21	1.00	1 (6%)
4	NAG	I	531	1	12,14,15	0.59	0	15,19,21	0.95	1 (6%)
4	NAG	I	541	1	12,14,15	0.58	0	15,19,21	1.02	0
4	NAG	K	621	1	12,14,15	0.59	0	15,19,21	0.92	1 (6%)
4	NAG	K	641	1	12,14,15	0.63	0	15,19,21	0.85	1 (6%)
4	NAG	K	651	1	12,14,15	0.66	0	15,19,21	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	332	1	-	0/6/23/26	0/1/1/1
4	NAG	D	261	2	-	0/6/23/26	0/1/1/1
4	NAG	E	341	1	-	0/6/23/26	0/1/1/1
4	NAG	G	411	1	-	0/6/23/26	0/1/1/1
4	NAG	G	431	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	451	1	-	0/6/23/26	0/1/1/1
4	NAG	I	521	1	-	0/6/23/26	0/1/1/1
4	NAG	I	531	1	-	0/6/23/26	0/1/1/1
4	NAG	I	541	1	-	0/6/23/26	0/1/1/1
4	NAG	K	621	1	-	0/6/23/26	0/1/1/1
4	NAG	K	641	1	-	0/6/23/26	0/1/1/1
4	NAG	K	651	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	261	NAG	O5-C5-C6	3.18	110.32	106.98
4	I	521	NAG	O5-C5-C6	3.14	110.28	106.98
4	K	621	NAG	O5-C5-C6	2.78	109.90	106.98
4	G	411	NAG	O5-C5-C6	2.56	109.67	106.98
4	G	431	NAG	O5-C5-C6	2.32	109.41	106.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/329 (98%)	-0.17	7 (2%) 59 65	20, 34, 57, 75	0
1	C	320/329 (97%)	-0.16	6 (1%) 64 70	27, 41, 59, 81	0
1	E	319/329 (96%)	-0.20	4 (1%) 74 80	22, 35, 50, 74	0
1	G	323/329 (98%)	-0.21	1 (0%) 91 95	21, 35, 51, 64	0
1	I	320/329 (97%)	-0.34	5 (1%) 68 75	22, 34, 54, 76	0
1	K	320/329 (97%)	-0.13	5 (1%) 68 75	25, 40, 58, 84	0
2	B	169/177 (95%)	0.70	22 (13%) 4 5	22, 46, 95, 111	0
2	D	170/177 (96%)	0.04	4 (2%) 56 63	22, 47, 65, 81	0
2	F	170/177 (96%)	-0.04	3 (1%) 65 72	22, 42, 61, 71	0
2	H	172/177 (97%)	-0.10	2 (1%) 75 81	23, 42, 62, 74	0
2	J	170/177 (96%)	0.47	21 (12%) 5 5	21, 47, 81, 87	0
2	L	169/177 (95%)	0.32	14 (8%) 11 14	23, 47, 83, 99	0
All	All	2945/3036 (97%)	-0.05	94 (3%) 45 52	20, 38, 70, 111	0

The worst 5 of 94 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	166	ALA	6.9
2	B	160	PRO	6.5
2	B	163	SER	6.2
2	L	168	LEU	5.7
1	E	290	THR	5.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	E	331	14/15	0.24	-	58,64,68,70	0
5	GAL	G	402	11/12	0.36	-	69,74,80,80	0
3	NAG	A	331	14/15	0.13	-	46,53,62,66	0
5	GAL	I	502	11/12	0.13	-	50,60,65,65	0
3	NAG	E	332	14/15	0.31	-	62,76,83,86	0
5	SIA	I	501	20/21	0.10	-	35,40,46,48	0
3	NAG	K	632	14/15	0.19	-	69,73,82,83	0
5	SIA	G	401	20/21	0.25	-	42,54,60,65	0
5	GAL	C	331	11/12	0.12	-	57,66,70,70	0
3	NAG	K	631	14/15	0.14	-	49,58,66,69	0
3	NAG	A	330	14/15	0.07	-	34,43,47,49	0
5	SIA	K	601	20/21	0.13	-	42,56,66,68	0
5	GAL	K	602	11/12	0.21	-	70,81,89,90	0
5	SIA	C	330	20/21	0.10	-	36,46,54,55	0

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	NAG	G	411	14/15	0.29	-	64,73,79,80	0
4	NAG	G	451	14/15	0.37	-	59,86,95,99	0
4	NAG	I	521	14/15	0.19	-	73,87,91,93	0
4	NAG	A	332	14/15	0.35	-	55,63,71,72	0
4	NAG	K	621	14/15	0.18	-	58,69,78,82	0
4	NAG	G	431	14/15	0.09	-	38,52,61,67	0
4	NAG	I	541	14/15	0.14	-	54,66,70,76	0
4	NAG	K	651	14/15	0.17	-	60,77,87,88	0
4	NAG	K	641	14/15	0.19	-	55,61,68,74	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NAG	D	261	14/15	0.15	-	57,65,72,73	0
4	NAG	E	341	14/15	0.25	-	68,75,83,85	0
4	NAG	I	531	14/15	0.14	-	40,54,58,59	0

6.5 Other polymers ⓘ

There are no such residues in this entry.