



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 09:42 AM GMT

PDB ID : 3UBP  
Title : DIAMIDOPHOSPHATE INHIBITED BACILLUS PASTEURII UREASE  
Authors : Benini, S.; Rypniewski, W.R.; Wilson, K.S.; Miletto, S.; Mangani, S.; Ciurli, S.  
Deposited on : 1998-12-16  
Resolution : 2.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	101	
2	B	126	
3	C	570	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6903 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (UREASE GAMMA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	101	Total	C	N	O	S	8	0	0
			781	494	133	148	6			

- Molecule 2 is a protein called PROTEIN (UREASE BETA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	122	Total	C	N	O	S	16	0	0
			951	589	171	190	1			

- Molecule 3 is a protein called PROTEIN (UREASE ALPHA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	570	Total	C	N	O	S	19	0	0
			4323	2714	743	843	23			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	19	GLU	ARG	VARIANT	UNP P41020
C	28	TRP	-	INSERTION	UNP P41020
C	29	ILE	GLY	VARIANT	UNP P41020
C	36	THR	TYR	VARIANT	UNP P41020
C	37	THR	TYR	VARIANT	UNP P41020
C	38	TYR	LEU	VARIANT	UNP P41020
C	263	LEU	VAL	VARIANT	UNP P41020
C	420	ILE	MET	VARIANT	UNP P41020

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Ni	0	0
			2	2		

- Molecule 5 is DIAMIDOPHOSPHATE (three-letter code: 2PA) (formula:  $\text{H}_5\text{N}_2\text{O}_2\text{P}$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	N	O	P	0	0
			5	2	2	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	124	Total	O	0	0
			124	124		
6	B	186	Total	O	0	0
			186	186		
6	C	531	Total	O	0	0
			531	531		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (UREASE GAMMA SUBUNIT)

Chain A: 




- Molecule 2: PROTEIN (UREASE BETA SUBUNIT)

Chain B: 



- Molecule 3: PROTEIN (UREASE ALPHA SUBUNIT)

Chain C: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	131.53Å 131.53Å 188.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	18.00 – 2.00 17.98 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (18.00-2.00) 100.0 (17.98-2.00)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	0.15	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.67 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.158 , 0.200 0.162 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	13.7	Xtriage
Anisotropy	0.413	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 40.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 65263 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6903	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2PA, NI, KCX, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.57	0/790	1.10	6/1065 (0.6%)
2	B	0.75	2/963 (0.2%)	1.19	5/1296 (0.4%)
3	C	0.59	2/4392 (0.0%)	1.17	22/5955 (0.4%)
All	All	0.61	4/6145 (0.1%)	1.16	33/8316 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	4
3	C	0	3
All	All	0	7

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	5	ASN	C-N	-7.08	1.17	1.34
3	C	547	LYS	CD-CE	-5.33	1.38	1.51
3	C	1	MET	N-CA	5.28	1.56	1.46
2	B	16	GLU	CD-OE2	5.20	1.31	1.25

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	566	ARG	NE-CZ-NH2	-13.05	113.78	120.30
3	C	5	ARG	NE-CZ-NH2	-12.70	113.95	120.30
2	B	111	GLU	N-CA-CB	10.86	130.14	110.60
3	C	376	ARG	NE-CZ-NH1	10.20	125.40	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	388	ARG	NE-CZ-NH2	-8.62	115.99	120.30

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	109	ASN	Mainchain
2	B	110	LYS	Peptide
2	B	125	VAL	Mainchain
2	B	8	VAL	Mainchain
3	C	289	ALA	Mainchain

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	781	0	804	13	1
2	B	951	0	936	14	0
3	C	4323	0	4290	48	0
4	C	2	0	0	0	0
5	C	5	0	4	1	0
6	A	124	0	0	8	2
6	B	186	0	0	8	0
6	C	531	0	0	10	6
All	All	6903	0	6034	73	9

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:C:325:LEU:HA	3:C:326:LYS:HE2	1.61	0.82
3:C:536:ASP:HB3	6:C:1392:HOH:O	1.84	0.77
2:B:13:ARG:NE	6:B:280:HOH:O	2.21	0.73
3:C:399:ASP:O	3:C:403:LEU:HG	1.90	0.71

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:109:ASN:ND2	6:B:269:HOH:O	2.24	0.71

The worst 5 of 9 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
6:A:150:HOH:O	6:A:150:HOH:O[2_665]	0.41	1.79
6:A:155:HOH:O	6:A:155:HOH:O[2_665]	0.41	1.79
6:C:1135:HOH:O	6:C:1135:HOH:O[12_565]	0.41	1.79
6:C:1228:HOH:O	6:C:1228:HOH:O[11_555]	1.23	0.97
6:C:1367:HOH:O	6:C:1367:HOH:O[10_665]	1.42	0.78

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/101 (98%)	97 (98%)	2 (2%)	0	100	100
2	B	120/126 (95%)	114 (95%)	5 (4%)	1 (1%)	27	17
3	C	567/570 (100%)	541 (95%)	22 (4%)	4 (1%)	30	20
All	All	786/797 (99%)	752 (96%)	29 (4%)	5 (1%)	33	24

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	397	GLY
2	B	99	ILE
3	C	275	HIS
3	C	283	HIS
3	C	367	MET

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of

similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	83 (98%)	2 (2%)	61	61
2	B	101/105 (96%)	101 (100%)	0	100	100
3	C	460/460 (100%)	447 (97%)	13 (3%)	56	54
All	All	646/650 (99%)	631 (98%)	15 (2%)	63	63

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	253	LEU
3	C	326	LYS
3	C	398	SER
3	C	249	HIS
3	C	395	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	267	ASN
3	C	565	GLN
3	C	328	ASN
3	C	6	GLN
3	C	519	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	KCX	C	220	3,4	11,11,12	7.39	4 (36%)	10,12,14	2.62	5 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KCX	C	220	3,4	-	0/8/10/12	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	220	KCX	O-C	16.74	1.22	1.11
3	C	220	KCX	CX-NZ	16.19	1.48	1.32
3	C	220	KCX	OQ1-CX	6.61	1.33	1.21
3	C	220	KCX	CA-C	3.20	1.54	1.48

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	220	KCX	CE-NZ-CX	-4.88	113.10	121.99
3	C	220	KCX	C-CA-N	-4.10	109.74	113.83
3	C	220	KCX	OQ2-CX-OQ1	-3.34	117.99	122.17
3	C	220	KCX	OQ2-CX-NZ	2.55	119.50	116.33
3	C	220	KCX	CD-CE-NZ	-2.18	105.66	112.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	2PA	C	902	4	4,4,4	4.90	3 (75%)	5,6,6	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	2PA	C	902	4	-	0/0/0/0	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	902	2PA	P2-O1	7.24	1.56	1.47
5	C	902	2PA	P2-N4	5.21	1.70	1.62
5	C	902	2PA	P2-N5	3.70	1.68	1.62

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	101/101 (100%)	-0.81	0 100 100	8, 12, 20, 26	4 (3%)
2	B	121/126 (96%)	-0.67	1 (0%) 83 84	10, 14, 27, 45	3 (2%)
3	C	570/570 (100%)	-0.70	7 (1%) 75 76	6, 12, 25, 53	9 (1%)
All	All	792/797 (99%)	-0.71	8 (1%) 79 80	6, 12, 26, 53	16 (2%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	328	ASN	4.0
3	C	327	GLN	3.9
3	C	397	GLY	3.7
3	C	329	ILE	3.0
3	C	522	ASN	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	KCX	C	220	12/13	0.07	-0.03	5,10,17,18	0

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	2PA	C	902	5/5	0.12	1.96	14,14,18,19	0
4	NI	C	900	1/1	0.04	-2.42	21,21,21,21	0
4	NI	C	901	1/1	0.02	-6.03	16,16,16,16	0

## 6.5 Other polymers

There are no such residues in this entry.