



Full wwPDB X-ray Structure Validation Report i

Feb 28, 2014 – 03:55 AM GMT

PDB ID : 1UDT
Title : Crystal structure of Human Phosphodiesterase 5 complexed with Sildenafil(Viagra)
Authors : Sung, B.-J.; Lee, J.I.; Heo, Y.-S.; Kim, J.H.; Moon, J.; Yoon, J.M.; Hyun, Y.-L.; Kim, E.; Eum, S.J.; Lee, T.G.; Cho, J.M.; Park, S.-Y.; Lee, J.-O.; Jeon, Y.H.; Hwang, K.Y.; Ro, S.
Deposited on : 2003-05-06
Resolution : 2.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	324	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2655 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cGMP-specific 3',5'-cyclic phosphodiesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	0	0	0
			2542	1617	442	465	18			


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 5-{2-ETHOXY-5-[(4-METHYLPIPERAZIN-1-YL)SULFONYL]PHENYL}-1-METHYL-3-PROPYL-1H,6H,7H-PYRAZOLO[4,3-D]PYRIMIDIN-7-ONE (three-letter code: VIA) (formula: C₂₂H₃₀N₆O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			33	22	6	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	78	Total	O	0	0
			78	78		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	60.12Å 155.63Å 89.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.88 – 2.30 19.88 – 2.29	Depositor EDS
% Data completeness (in resolution range)	87.9 (19.88-2.30) 87.0 (19.88-2.29)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.89 (at 2.28Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.193 , 0.251 0.193 , 0.251	Depositor DCC
R_{free} test set	823 reflections (4.89%)	DCC
Wilson B-factor (Å ²)	30.4	Xtriage
Anisotropy	0.377	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 34.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 16903 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2655	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, VIA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/2590	0.56	0/3493

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2542	0	2552	76	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	33	0	30	0	0
5	A	78	0	0	9	0
All	All	2655	0	2582	76	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 15.

All (76) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:740:ARG:HH11	1:A:740:ARG:HA	1.21	1.04
1:A:585:VAL:HA	1:A:590:MET:HE2	1.51	0.92
1:A:860:GLN:CD	1:A:860:GLN:OXT	2.17	0.83
1:A:605:ASN:HD21	1:A:688:GLN:HE21	1.26	0.83
1:A:860:GLN:OXT	1:A:860:GLN:NE2	2.18	0.77
1:A:605:ASN:ND2	1:A:688:GLN:HE21	1.85	0.74
1:A:797:LEU:HB3	1:A:799:ILE:HD13	1.71	0.71
1:A:740:ARG:HA	1:A:740:ARG:NH1	2.03	0.70
1:A:813:ILE:HG12	1:A:814:PRO:HD3	1.74	0.68
1:A:772:TRP:HB3	1:A:773:PRO:HD3	1.78	0.66
1:A:757:ALA:O	1:A:761:THR:HG22	1.97	0.64
1:A:690:LEU:HD13	1:A:713:LEU:HD11	1.80	0.64
1:A:813:ILE:CG1	1:A:814:PRO:HD3	2.29	0.62
1:A:714:LYS:O	1:A:718:GLN:HG3	2.01	0.61
1:A:564:PHE:HB2	1:A:777:ARG:NH2	2.16	0.61
1:A:679:SER:HB3	1:A:682:GLU:HG2	1.85	0.57
1:A:682:GLU:HG3	1:A:724:ASP:HB2	1.85	0.57
1:A:853:TRP:CE3	1:A:856:LEU:HD12	2.39	0.57
1:A:585:VAL:CA	1:A:590:MET:HE2	2.32	0.57
1:A:771:PRO:HD2	1:A:774:ILE:HD12	1.86	0.57
1:A:676:TYR:CZ	1:A:678:HIS:HA	2.40	0.57
1:A:537:THR:N	5:A:16:HOH:O	2.39	0.56
1:A:797:LEU:CB	1:A:799:ILE:HD13	2.36	0.56
1:A:705:SER:OG	1:A:708:GLU:HG3	2.07	0.55
1:A:761:THR:HB	5:A:53:HOH:O	2.07	0.54
1:A:701:LEU:HB2	5:A:22:HOH:O	2.05	0.54
1:A:639:THR:OG1	1:A:642:GLU:HG3	2.07	0.54
1:A:537:THR:O	1:A:541:GLN:HG3	2.07	0.54
1:A:781:LEU:HD11	5:A:4:HOH:O	2.10	0.52
1:A:806:ASN:OD1	1:A:809:LYS:HB2	2.09	0.52
1:A:753:GLU:HG2	5:A:60:HOH:O	2.10	0.52
1:A:679:SER:HB3	1:A:682:GLU:CG	2.40	0.51
1:A:646:LEU:HD11	1:A:756:LEU:HD22	1.92	0.51
1:A:748:ASP:HB3	1:A:751:GLN:HB2	1.92	0.51
1:A:717:LYS:O	1:A:721:LEU:HG	2.11	0.50
1:A:561:PHE:CD2	1:A:623:GLN:HG3	2.47	0.50
1:A:789:GLN:O	1:A:793:GLU:HB2	2.12	0.49
1:A:848:LYS:O	1:A:852:LYS:HG2	2.13	0.48
1:A:854:GLN:O	1:A:858:GLU:HG3	2.14	0.48
1:A:732:ARG:HD2	1:A:736:PHE:CE2	2.47	0.48
1:A:810:LYS:O	1:A:813:ILE:HG23	2.14	0.48
1:A:569:LEU:C	1:A:569:LEU:HD23	2.34	0.48
1:A:826:LEU:HD13	1:A:846:CYS:SG	2.54	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:777:ARG:HD2	5:A:4:HOH:O	2.14	0.47
1:A:741:LYS:HB2	1:A:743:GLN:HG2	1.96	0.47
1:A:583:ASN:O	1:A:587:ASN:HB2	2.14	0.47
1:A:738:LEU:HD13	1:A:751:GLN:OE1	2.15	0.47
1:A:554:LEU:HD13	1:A:574:CYS:HA	1.96	0.47
1:A:806:ASN:HD21	1:A:808:GLU:HB2	1.80	0.46
1:A:548:VAL:HG23	1:A:548:VAL:O	2.15	0.46
1:A:585:VAL:HA	1:A:590:MET:CE	2.37	0.46
1:A:840:PHE:N	1:A:841:PRO:CD	2.78	0.45
1:A:562:SER:HB2	1:A:774:ILE:HD13	1.99	0.45
1:A:830:GLU:HG3	1:A:831:ALA:N	2.31	0.45
1:A:814:PRO:O	1:A:818:VAL:HG23	2.17	0.45
1:A:649:ALA:O	1:A:653:HIS:N	2.50	0.44
1:A:569:LEU:O	1:A:569:LEU:HD23	2.17	0.44
1:A:758:MET:O	1:A:761:THR:HG23	2.17	0.44
1:A:787:PHE:HB2	1:A:807:ARG:HH21	1.83	0.43
1:A:681:MET:HG3	1:A:685:HIS:NE2	2.33	0.43
1:A:682:GLU:HA	1:A:685:HIS:CD2	2.54	0.43
1:A:728:TYR:CZ	1:A:732:ARG:HD3	2.54	0.42
1:A:662:ASN:HB3	1:A:663:SER:H	1.74	0.42
1:A:780:GLU:HB3	5:A:27:HOH:O	2.20	0.41
1:A:800:GLU:HA	1:A:801:PRO:HD3	1.88	0.41
1:A:707:GLU:HG3	1:A:708:GLU:N	2.36	0.41
1:A:544:ALA:O	1:A:592:HIS:HE1	2.04	0.41
1:A:677:CYS:HB2	5:A:23:HOH:O	2.20	0.41
1:A:562:SER:HB2	1:A:774:ILE:CD1	2.51	0.41
1:A:725:LEU:HA	1:A:725:LEU:HD12	1.87	0.41
1:A:634:ILE:O	1:A:634:ILE:HG22	2.20	0.41
1:A:810:LYS:C	1:A:812:LYS:H	2.23	0.40
1:A:809:LYS:HD2	1:A:809:LYS:N	2.36	0.40
1:A:625:MET:HE2	1:A:625:MET:HB2	1.86	0.40
1:A:817:GLN:O	1:A:821:ILE:HG13	2.22	0.40
1:A:585:VAL:HB	5:A:6:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	309/324 (95%)	297 (96%)	11 (4%)	1 (0%)	50 60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	564	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	279/289 (96%)	259 (93%)	20 (7%)	21 25

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	543	LEU
1	A	558	ASP
1	A	563	ASP
1	A	591	LYS
1	A	595	LEU
1	A	605	ASN
1	A	629	LEU
1	A	647	LEU
1	A	682	GLU
1	A	713	LEU
1	A	725	LEU
1	A	740	ARG
1	A	761	THR
1	A	775	GLN
1	A	797	LEU
1	A	806	ASN
1	A	812	LYS
1	A	826	LEU
1	A	828	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	830	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	541	GLN
1	A	583	ASN
1	A	592	HIS
1	A	605	ASN
1	A	635	GLN
1	A	662	ASN
1	A	694	ASN
1	A	699	GLN
1	A	718	GLN
1	A	750	HIS
1	A	798	ASN
1	A	811	ASN
1	A	834	HIS
1	A	849	ASN
1	A	851	GLN
1	A	859	GLN
1	A	860	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	VIA	A	1000	-	36,36,36	4.29	19 (52%)	49,53,53	3.79	23 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	VIA	A	1000	-	-	0/22/32/32	0/2/4/4

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1000	VIA	C7-S10	-10.40	1.60	1.76
4	A	1000	VIA	O12-S10	-9.41	1.31	1.43
4	A	1000	VIA	C19-N14	9.34	1.58	1.47
4	A	1000	VIA	O11-S10	-8.79	1.32	1.43
4	A	1000	VIA	C6-C7	6.85	1.50	1.38
4	A	1000	VIA	C9-C4	6.56	1.50	1.40
4	A	1000	VIA	C9-C21	-6.12	1.42	1.49
4	A	1000	VIA	C21-N26	4.56	1.40	1.32
4	A	1000	VIA	C5-C6	3.96	1.46	1.38
4	A	1000	VIA	C5-C4	3.93	1.48	1.39
4	A	1000	VIA	C8-C7	3.83	1.46	1.39
4	A	1000	VIA	O3-C4	3.81	1.45	1.37
4	A	1000	VIA	C15-N14	3.78	1.52	1.47
4	A	1000	VIA	C30-N29	3.64	1.40	1.34
4	A	1000	VIA	C8-C9	3.42	1.45	1.39
4	A	1000	VIA	C21-N22	3.34	1.40	1.34
4	A	1000	VIA	C16-N17	3.07	1.53	1.46
4	A	1000	VIA	S10-N14	-2.59	1.59	1.63
4	A	1000	VIA	C25-N26	2.48	1.42	1.37

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1000	VIA	C18-C19-N14	10.50	117.33	109.02
4	A	1000	VIA	C16-C15-N14	8.76	115.95	109.02
4	A	1000	VIA	O12-S10-O11	-8.02	105.07	119.38
4	A	1000	VIA	C16-N17-C18	6.96	119.25	109.54
4	A	1000	VIA	C15-C16-N17	6.79	118.31	110.82
4	A	1000	VIA	O11-S10-N14	6.47	113.30	106.71
4	A	1000	VIA	O11-S10-C7	6.20	116.17	107.94
4	A	1000	VIA	O12-S10-N14	5.91	112.73	106.71
4	A	1000	VIA	C7-S10-N14	-5.34	100.49	107.29
4	A	1000	VIA	C30-N29-N28	5.27	109.03	104.48
4	A	1000	VIA	C15-N14-C19	-5.02	105.95	112.16
4	A	1000	VIA	C19-C18-N17	4.87	116.19	110.82
4	A	1000	VIA	C23-C24-C25	-4.35	116.80	119.92
4	A	1000	VIA	C15-N14-S10	3.53	123.43	117.00
4	A	1000	VIA	C2-O3-C4	3.33	125.03	118.02
4	A	1000	VIA	C8-C9-C4	3.22	121.63	118.17
4	A	1000	VIA	C31-N28-C24	3.21	129.92	125.00
4	A	1000	VIA	C33-C32-C30	-3.16	102.78	115.48
4	A	1000	VIA	N22-C21-N26	-2.70	119.74	122.70
4	A	1000	VIA	C6-C7-C8	-2.59	117.43	120.49
4	A	1000	VIA	C9-C21-N26	2.54	123.13	117.87
4	A	1000	VIA	C5-C6-C7	2.45	122.29	119.50
4	A	1000	VIA	C5-C4-C9	-2.02	117.71	121.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/324 (96%)	-0.05	9 (2%) 49 59	20, 36, 66, 90	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	860	GLN	3.1
1	A	664	TYR	2.9
1	A	742	ASN	2.7
1	A	809	LYS	2.7
1	A	741	LYS	2.6
1	A	808	GLU	2.6
1	A	547	VAL	2.4
1	A	707	GLU	2.2
1	A	798	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	VIA	A	1000	33/33	0.15	0.28	23,36,45,52	0
3	MG	A	1002	1/1	0.07	-1.39	27,27,27,27	0
2	ZN	A	1001	1/1	0.05	-3.98	33,33,33,33	0

6.5 Other polymers ⓘ

There are no such residues in this entry.