



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 10:08 PM GMT

PDB ID : 3UFT
Title : Structure of rat nitric oxide synthase heme domain in complex with 6-(((3R,4R)-4-(4-(3-chloro-5-fluorophenoxy)butoxy)pyrrolidin-3-yl)methyl)-4-methylpyridin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2011-11-01
Resolution : 2.08 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3396 (2.10-2.06)
Clashscore	79885	4085 (2.10-2.06)
Ramachandran outliers	78287	4045 (2.10-2.06)
Sidechain outliers	78261	4046 (2.10-2.06)
RSRZ outliers	66119	3397 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	422	
1	B	422	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
5	ACT	A	860	-	X
5	ACT	B	860	-	X
6	ZN	A	900	-	X

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 7013 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	0	0
			3313	2121	566	605	21			
1	B	411	Total	C	N	O	S	0	1	0
			3348	2142	574	610	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 6-({(3R,4R)-4-[4-(3-CHLORO-5-FLUOROPHENOXY)BUTOXY]PYRROLIDIN-3-YL}METHYL)-4-METHYLPYRIDIN-2-AMINE (three-letter code: HW7) (formula: C₂₁H₂₇ClFN₃O₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	Cl	F	N	O	0	0
			28	21	1	1	3	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total	C	Cl	F	N	O	0	0
			28	21	1	1	3	2		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	69	Total	O	0	0
			69	69		
7	B	98	Total	O	0	0
			98	98		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nitric oxide synthase, brain

Chain A: 



- Molecule 1: Nitric oxide synthase, brain

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.84Å 111.22Å 164.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.99 – 2.08 46.99 – 2.08	Depositor EDS
% Data completeness (in resolution range)	99.1 (46.99-2.08) 99.1 (46.99-2.08)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.206 , 0.260 0.230 , 0.272	Depositor DCC
R_{free} test set	2849 reflections (4.95%)	DCC
Wilson B-factor (Å ²)	43.7	Xtriage
Anisotropy	0.391	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 57575 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7013	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HW7, ZN, H4B, HEM, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.84	1/3406 (0.0%)	0.84	1/4621 (0.0%)
1	B	1.02	9/3444 (0.3%)	0.93	4/4669 (0.1%)
All	All	0.94	10/6850 (0.1%)	0.88	5/9290 (0.1%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	421	TRP	CB-CG	7.27	1.63	1.50
1	B	706	TYR	CD1-CE1	6.57	1.49	1.39
1	B	482	TYR	CD2-CE2	-5.88	1.30	1.39
1	B	477	SER	C-O	-5.65	1.12	1.23
1	B	706	TYR	CD2-CE2	5.55	1.47	1.39
1	B	567	VAL	CB-CG1	5.52	1.64	1.52
1	B	582	CYS	CB-SG	5.22	1.91	1.82
1	A	609	GLU	CG-CD	5.12	1.59	1.51
1	B	566	ALA	C-O	5.08	1.33	1.23
1	B	408	ALA	CA-CB	-5.05	1.41	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	489	ASP	CB-CG-OD2	7.64	125.18	118.30
1	A	410	ARG	NE-CZ-NH2	7.59	124.10	120.30
1	B	430	ARG	NE-CZ-NH1	5.81	123.21	120.30
1	B	418	ARG	NE-CZ-NH1	-5.67	117.47	120.30
1	B	596	ARG	NE-CZ-NH2	-5.07	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3313	0	0	20	0
1	B	3348	0	0	11	0
2	A	43	0	0	0	0
2	B	43	0	0	1	0
3	A	17	0	0	0	0
3	B	17	0	0	0	0
4	A	28	0	0	0	0
4	B	28	0	0	1	0
5	A	4	0	0	0	0
5	B	4	0	0	0	0
6	A	1	0	0	0	0
7	A	69	0	0	0	0
7	B	98	0	0	3	0
All	All	7013	0	0	29	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (29) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:750:HEM:O2D	4:B:800:HW7:N01	2.36	0.59
1:A:514:ARG:CG	1:A:514:ARG:NH2	2.68	0.56
1:B:396:LEU:O	1:B:430:ARG:NH1	2.38	0.56
1:A:664:ASN:OD1	1:A:667:ARG:NH2	2.40	0.54
1:A:632:GLU:OE2	1:B:628:GLN:NE2	2.40	0.54
1:A:350:THR:N	1:A:353:GLN:NE2	2.56	0.53
1:A:387:GLU:O	1:A:391:THR:OG1	2.28	0.52
1:B:512:ALA:N	7:B:103:HOH:O	2.42	0.52
1:A:504:ILE:O	1:A:508:GLN:N	2.43	0.51
1:B:481:ARG:CZ	1:B:498:ASN:OD1	2.59	0.51
1:A:628:GLN:NE2	1:B:632:GLU:OE2	2.48	0.47
1:A:476:ASN:N	1:A:476:ASN:OD1	2.49	0.46
1:A:445:HIS:C	1:A:445:HIS:CD2	2.89	0.46
1:A:537:PRO:O	1:A:540:LEU:N	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:317:HIS:O	1:A:320:SER:CB	2.65	0.45
1:A:448:TYR:O	1:A:456:ARG:NH2	2.49	0.45
1:B:455:LEU:N	1:B:455:LEU:CD1	2.79	0.45
1:B:337:LEU:N	7:B:37:HOH:O	2.51	0.44
1:A:597:ASP:OD1	1:A:603:ARG:NH2	2.50	0.43
1:A:544:VAL:N	1:A:561:TRP:O	2.51	0.43
1:A:371:ARG:CG	1:A:371:ARG:NH2	2.82	0.43
1:A:710:PRO:O	1:A:714:HIS:N	2.53	0.42
1:A:467:ASP:OD1	1:A:467:ASP:C	2.58	0.42
1:B:448:TYR:C	1:B:448:TYR:CD2	2.94	0.41
1:A:328:GLU:O	1:B:324:THR:N	2.54	0.41
1:A:547:ARG:NH1	1:A:644:ASP:OD1	2.54	0.40
1:B:548:HIS:NE2	1:B:632:GLU:OE1	2.54	0.40
1:A:431:ASP:C	1:A:431:ASP:OD2	2.59	0.40
1:B:456:ARG:NH2	7:B:90:HOH:O	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/422 (96%)	382 (95%)	21 (5%)	0	100	100
1	B	408/422 (97%)	390 (96%)	18 (4%)	0	100	100
All	All	811/844 (96%)	772 (95%)	39 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/377 (96%)	338 (93%)	25 (7%)	22	17
1	B	367/377 (97%)	357 (97%)	10 (3%)	57	60
All	All	730/754 (97%)	695 (95%)	35 (5%)	35	31

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	328	GLU
1	A	350	THR
1	A	352	ASP
1	A	353	GLN
1	A	371	ARG
1	A	380	ARG
1	A	381	LEU
1	A	423	LYS
1	A	469	LYS
1	A	476	ASN
1	A	481	ARG
1	A	489	ASP
1	A	511	LYS
1	A	514	ARG
1	A	527	ASN
1	A	547	ARG
1	A	550	LYS
1	A	552	ASP
1	A	555	LYS
1	A	569	ASN
1	A	578	GLU
1	A	615	ASP
1	A	645	LYS
1	A	653	SER
1	A	701	THR
1	B	332	MET
1	B	350	THR
1	B	353	GLN
1	B	360	GLU
1	B	392	SER
1	B	480	ILE
1	B	486	LYS
1	B	507	GLN
1	B	547	ARG
1	B	715	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	750	1	49,50,50	3.54	13 (26%)	46,82,82	2.27	11 (23%)
3	H4B	A	760	-	18,18,18	1.10	2 (11%)	24,26,26	2.00	10 (41%)
4	HW7	A	800	-	30,30,30	1.10	3 (10%)	40,40,40	2.08	10 (25%)
5	ACT	A	860	-	1,3,3	0.13	0	0,3,3	0.00	-
2	HEM	B	750	1	49,50,50	2.88	22 (44%)	46,82,82	2.29	12 (26%)
3	H4B	B	760	-	18,18,18	1.76	4 (22%)	24,26,26	2.18	6 (25%)
4	HW7	B	800	-	30,30,30	1.27	3 (10%)	40,40,40	2.14	13 (32%)
5	ACT	B	860	-	1,3,3	1.06	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/14/114/114	0/0/8/8
3	H4B	A	760	-	-	0/8/17/17	0/0/2/2
4	HW7	A	800	-	-	0/13/23/23	0/3/3/3
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/14/114/114	0/0/8/8
3	H4B	B	760	-	-	0/8/17/17	0/0/2/2
4	HW7	B	800	-	-	0/13/23/23	0/3/3/3
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (47) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C3D-C4D	-13.63	1.41	1.44
2	A	750	HEM	C2D-C1D	-10.18	1.42	1.44
2	A	750	HEM	C2B-C1B	9.97	1.47	1.44
2	B	750	HEM	C2D-C1D	9.25	1.46	1.44
2	B	750	HEM	C3D-C4D	6.70	1.46	1.44
2	A	750	HEM	C3B-C2B	-6.38	1.32	1.43
2	B	750	HEM	C3C-C2C	-6.07	1.33	1.43
2	A	750	HEM	C3C-C2C	-5.42	1.34	1.43
2	B	750	HEM	C3B-C2B	-5.38	1.34	1.43
2	A	750	HEM	C3D-C2D	5.03	1.52	1.43
2	B	750	HEM	C4A-C3A	4.97	1.46	1.40
2	B	750	HEM	C3C-CAC	4.73	1.55	1.40
2	A	750	HEM	C3B-CAB	4.59	1.54	1.40
2	B	750	HEM	C3D-C2D	4.35	1.51	1.43
3	B	760	H4B	C2-N2	4.17	1.38	1.32
2	B	750	HEM	C3B-CAB	4.11	1.53	1.40
2	A	750	HEM	C3C-CAC	4.06	1.53	1.40
2	A	750	HEM	FE-NA	3.78	2.08	1.92
4	B	800	HW7	C25-CL2	3.64	1.83	1.74
3	B	760	H4B	C2-N1	3.63	1.38	1.33
4	B	800	HW7	C06-N01	3.44	1.40	1.34
2	B	750	HEM	FE-NA	3.30	2.06	1.92
4	A	800	HW7	C25-CL2	3.24	1.82	1.74
2	B	750	HEM	FE-ND	3.23	2.09	1.97
2	B	750	HEM	CHB-C1B	3.16	1.40	1.35
2	A	750	HEM	CMC-C2C	3.05	1.56	1.47
4	B	800	HW7	C24-C23	3.03	1.43	1.37
2	B	750	HEM	CMB-C2B	2.95	1.56	1.47
2	B	750	HEM	CHA-C4D	2.91	1.40	1.35
3	B	760	H4B	O4-C4	2.88	1.30	1.24
2	B	750	HEM	C1A-NA	2.84	1.42	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	CMC-C2C	2.73	1.55	1.47
2	B	750	HEM	CMD-C2D	2.69	1.55	1.47
2	A	750	HEM	CMB-C2B	2.67	1.55	1.47
3	B	760	H4B	C7-N8	2.59	1.50	1.46
3	A	760	H4B	C4-N3	-2.47	1.33	1.37
2	B	750	HEM	CHD-C4C	2.43	1.40	1.36
4	A	800	HW7	F23-C23	-2.43	1.30	1.36
2	B	750	HEM	C4A-NA	2.43	1.41	1.36
2	B	750	HEM	FE-NC	2.42	2.06	1.97
2	B	750	HEM	CAD-C3D	2.41	1.60	1.51
2	A	750	HEM	CHA-C4D	2.40	1.39	1.35
2	A	750	HEM	CMD-C2D	2.33	1.54	1.47
4	A	800	HW7	C24-C23	2.28	1.41	1.37
2	B	750	HEM	CAA-C2A	2.08	1.55	1.52
2	B	750	HEM	CMA-C3A	2.04	1.55	1.51
3	A	760	H4B	C2-N2	2.04	1.35	1.32

All (62) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	750	HEM	C3B-C4B-NB	-10.18	106.71	114.00
2	A	750	HEM	C3B-C4B-NB	-7.02	108.98	114.00
4	A	800	HW7	C02-N01-C06	6.54	122.82	118.23
2	A	750	HEM	CBA-CAA-C2A	-6.51	101.22	112.69
4	B	800	HW7	C2'-C3'-C4'	-6.43	98.46	103.48
3	B	760	H4B	C4-C4A-C8A	5.64	119.79	114.56
2	A	750	HEM	CBD-CAD-C3D	-5.40	102.58	114.37
2	B	750	HEM	C4A-CHB-C1B	-5.30	120.49	127.47
4	A	800	HW7	C24-C25-CL2	5.13	125.48	119.14
3	B	760	H4B	N2-C2-N3	4.98	123.34	117.86
4	B	800	HW7	C24-C25-CL2	4.94	125.25	119.14
4	B	800	HW7	C21-C26-C25	4.51	122.96	117.70
2	A	750	HEM	C1A-CHA-C4D	-4.45	121.61	127.47
3	A	760	H4B	N2-C2-N3	4.20	122.48	117.86
2	B	750	HEM	C4D-ND-C1D	4.16	109.42	105.16
2	A	750	HEM	C4D-ND-C1D	4.13	109.39	105.16
4	B	800	HW7	C02-N01-C06	4.02	121.05	118.23
2	B	750	HEM	CHC-C4B-NB	3.89	127.81	124.58
2	A	750	HEM	CHA-C4D-ND	3.63	129.29	124.31
4	A	800	HW7	C21-C26-C25	3.57	121.87	117.70
3	A	760	H4B	C4A-C8A-N8	3.45	123.97	119.23
3	A	760	H4B	C4-C4A-C8A	3.40	117.71	114.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	800	HW7	F23-C23-C22	-3.32	113.10	118.22
4	A	800	HW7	C3'-C2'-N1'	-3.23	99.67	105.28
2	A	750	HEM	C2D-C1D-ND	-3.20	109.15	112.93
3	B	760	H4B	C6-C7-N8	-3.14	107.60	111.66
2	B	750	HEM	C2D-C1D-ND	-3.08	109.29	112.93
4	B	800	HW7	C05-C06-N01	-3.08	119.57	122.99
2	A	750	HEM	CHD-C4C-NC	3.06	127.39	124.73
3	A	760	H4B	C2-N1-C8A	3.04	121.93	117.61
4	A	800	HW7	C5'-N1'-C2'	2.99	113.46	105.92
3	A	760	H4B	C4A-N5-C6	-2.84	113.42	121.16
4	A	800	HW7	C26-C25-CL2	-2.84	115.63	119.14
4	A	800	HW7	C2'-C3'-C4'	2.67	105.57	103.48
2	A	750	HEM	CMA-C3A-C4A	-2.66	124.53	128.62
4	B	800	HW7	O09-C3'-C2'	-2.64	103.37	110.63
4	B	800	HW7	C4'-C5'-N1'	-2.51	100.69	105.75
3	A	760	H4B	C7-N8-C8A	-2.50	118.52	121.36
2	B	750	HEM	C4C-NC-C1C	2.50	108.14	105.53
4	B	800	HW7	C3'-C2'-N1'	-2.50	100.95	105.28
4	A	800	HW7	C05-C06-N01	-2.48	120.23	122.99
4	B	800	HW7	F23-C23-C24	2.45	122.00	118.22
4	B	800	HW7	C26-C25-CL2	-2.45	116.12	119.14
3	B	760	H4B	C7-C6-C9	-2.44	109.89	113.66
2	B	750	HEM	CBD-CAD-C3D	-2.42	109.10	114.37
4	A	800	HW7	F23-C23-C24	2.37	121.87	118.22
2	A	750	HEM	C4A-C3A-C2A	2.37	108.64	107.00
4	B	800	HW7	C5'-N1'-C2'	2.34	111.83	105.92
2	B	750	HEM	O2A-CGA-CBA	2.32	122.42	114.22
3	A	760	H4B	C4-C4A-N5	2.24	122.36	119.10
4	B	800	HW7	C26-C25-C24	-2.21	118.59	121.65
2	B	750	HEM	C1B-NB-C4B	2.20	107.41	105.16
3	A	760	H4B	N3-C2-N1	-2.19	118.71	121.78
3	B	760	H4B	N2-C2-N1	-2.19	117.35	120.31
2	B	750	HEM	CAD-C3D-C4D	2.17	128.44	124.53
3	A	760	H4B	C4A-C4-N3	2.16	119.61	114.06
2	B	750	HEM	CMA-C3A-C4A	-2.11	125.37	128.62
3	B	760	H4B	C4A-N5-C6	-2.10	115.44	121.16
4	B	800	HW7	C10-O09-C3'	2.10	118.99	113.93
3	A	760	H4B	C7-C6-C9	-2.05	110.50	113.66
2	B	750	HEM	C1A-CHA-C4D	-2.05	124.78	127.47
2	A	750	HEM	O1A-CGA-CBA	-2.03	116.03	123.03

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/422 (96%)	0.65	26 (6%) 19 21	33, 62, 99, 125	0
1	B	411/422 (97%)	0.53	11 (2%) 52 56	32, 52, 77, 103	0
All	All	818/844 (96%)	0.59	37 (4%) 32 34	32, 56, 93, 125	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	348	VAL	6.6
1	B	718	GLY	4.2
1	B	352	ASP	3.6
1	B	351	LYS	3.6
1	A	355	PHE	3.6
1	B	389	GLU	3.4
1	A	393	THR	3.3
1	A	554	PHE	3.3
1	B	350	THR	3.2
1	B	300	PHE	3.2
1	A	351	LYS	2.8
1	B	321	THR	2.7
1	A	366	TYR	2.6
1	A	466	THR	2.6
1	A	713	THR	2.5
1	A	370	LYS	2.4
1	B	375	LYS	2.4
1	A	606	ILE	2.4
1	A	552	ASP	2.4
1	A	350	THR	2.4
1	A	394	TYR	2.4
1	A	551	PHE	2.3
1	A	506	ILE	2.3
1	A	486	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	352	ASP	2.3
1	A	322	LEU	2.2
1	A	299	ARG	2.2
1	B	717	LYS	2.2
1	A	330	ILE	2.2
1	A	557	LEU	2.2
1	B	619	ARG	2.2
1	A	385	ASN	2.1
1	A	577	LEU	2.1
1	A	361	PHE	2.1
1	A	556	ASP	2.1
1	A	300	PHE	2.1
1	A	715	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	ACT	A	860	4/4	0.29	7.52	63,63,63,66	0
6	ZN	A	900	1/1	0.15	3.04	50,50,50,50	0
5	ACT	B	860	4/4	0.20	2.35	57,59,60,61	0
2	HEM	B	750	43/43	0.19	1.35	29,44,53,67	0
4	HW7	A	800	28/28	0.17	0.83	54,62,73,75	0
4	HW7	B	800	28/28	0.16	0.70	43,55,58,62	0
2	HEM	A	750	43/43	0.16	0.32	35,39,57,66	0
3	H4B	B	760	17/17	0.14	-0.16	35,45,49,50	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	H4B	A	760	17/17	0.15	-0.18	49,52,55,59	0

6.5 Other polymers ⓘ

There are no such residues in this entry.