



wwPDB X-ray Structure Validation Summary Report

Nov 2, 2014 – 10:45 PM EST

PDB ID : 3UGT
Title : Crystal structure of the yeast mitochondrial threonyl-tRNA synthetase - orthorhombic crystal form
Authors : Peterson, K.M.; Ling, J.; Simonovic, I.; Cho, C.; Soll, D.; Simonovic, M.
Deposited on : 2011-11-02
Resolution : 3.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.16 November 2013
Xtriage (Phenix)	:	dev-1439
EDS	:	stable24103
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.1.3
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable24103

1 Overall quality at a glance

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1020 (3.86-3.34)
Clashscore	79885	1155 (3.80-3.40)
Ramachandran outliers	78287	1109 (3.80-3.40)
Sidechain outliers	78261	1108 (3.80-3.40)
RSRZ outliers	66119	1000 (3.84-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	460	
1	B	460	
1	C	460	
1	D	460	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13459 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Threonyl-tRNA synthetase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	419	Total	C	N	O	S	0	3	0
			3397	2195	571	613	18			
1	B	419	Total	C	N	O	S	3	4	0
			3285	2115	553	601	16			
1	C	427	Total	C	N	O	S	1	2	0
			3432	2206	574	634	18			
1	D	418	Total	C	N	O	S	0	1	0
			3295	2121	552	605	17			

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	MET	-	EXPRESSION TAG	UNP P07236
A	4	GLY	-	EXPRESSION TAG	UNP P07236
A	5	SER	-	EXPRESSION TAG	UNP P07236
A	6	SER	-	EXPRESSION TAG	UNP P07236
A	7	HIS	-	EXPRESSION TAG	UNP P07236
A	8	HIS	-	EXPRESSION TAG	UNP P07236
A	9	HIS	-	EXPRESSION TAG	UNP P07236
A	10	HIS	-	EXPRESSION TAG	UNP P07236
A	11	HIS	-	EXPRESSION TAG	UNP P07236
A	12	HIS	-	EXPRESSION TAG	UNP P07236
A	13	SER	-	EXPRESSION TAG	UNP P07236
A	14	SER	-	EXPRESSION TAG	UNP P07236
A	15	GLY	-	EXPRESSION TAG	UNP P07236
A	16	LEU	-	EXPRESSION TAG	UNP P07236
A	17	VAL	-	EXPRESSION TAG	UNP P07236
A	18	PRO	-	EXPRESSION TAG	UNP P07236
A	19	ARG	-	EXPRESSION TAG	UNP P07236
A	20	GLY	-	EXPRESSION TAG	UNP P07236
A	21	SER	-	EXPRESSION TAG	UNP P07236
A	22	HIS	-	EXPRESSION TAG	UNP P07236
A	23	MET	-	EXPRESSION TAG	UNP P07236

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	ALA	-	EXPRESSION TAG	UNP P07236
A	25	SER	-	EXPRESSION TAG	UNP P07236
B	3	MET	-	EXPRESSION TAG	UNP P07236
B	4	GLY	-	EXPRESSION TAG	UNP P07236
B	5	SER	-	EXPRESSION TAG	UNP P07236
B	6	SER	-	EXPRESSION TAG	UNP P07236
B	7	HIS	-	EXPRESSION TAG	UNP P07236
B	8	HIS	-	EXPRESSION TAG	UNP P07236
B	9	HIS	-	EXPRESSION TAG	UNP P07236
B	10	HIS	-	EXPRESSION TAG	UNP P07236
B	11	HIS	-	EXPRESSION TAG	UNP P07236
B	12	HIS	-	EXPRESSION TAG	UNP P07236
B	13	SER	-	EXPRESSION TAG	UNP P07236
B	14	SER	-	EXPRESSION TAG	UNP P07236
B	15	GLY	-	EXPRESSION TAG	UNP P07236
B	16	LEU	-	EXPRESSION TAG	UNP P07236
B	17	VAL	-	EXPRESSION TAG	UNP P07236
B	18	PRO	-	EXPRESSION TAG	UNP P07236
B	19	ARG	-	EXPRESSION TAG	UNP P07236
B	20	GLY	-	EXPRESSION TAG	UNP P07236
B	21	SER	-	EXPRESSION TAG	UNP P07236
B	22	HIS	-	EXPRESSION TAG	UNP P07236
B	23	MET	-	EXPRESSION TAG	UNP P07236
B	24	ALA	-	EXPRESSION TAG	UNP P07236
B	25	SER	-	EXPRESSION TAG	UNP P07236
C	3	MET	-	EXPRESSION TAG	UNP P07236
C	4	GLY	-	EXPRESSION TAG	UNP P07236
C	5	SER	-	EXPRESSION TAG	UNP P07236
C	6	SER	-	EXPRESSION TAG	UNP P07236
C	7	HIS	-	EXPRESSION TAG	UNP P07236
C	8	HIS	-	EXPRESSION TAG	UNP P07236
C	9	HIS	-	EXPRESSION TAG	UNP P07236
C	10	HIS	-	EXPRESSION TAG	UNP P07236
C	11	HIS	-	EXPRESSION TAG	UNP P07236
C	12	HIS	-	EXPRESSION TAG	UNP P07236
C	13	SER	-	EXPRESSION TAG	UNP P07236
C	14	SER	-	EXPRESSION TAG	UNP P07236
C	15	GLY	-	EXPRESSION TAG	UNP P07236
C	16	LEU	-	EXPRESSION TAG	UNP P07236
C	17	VAL	-	EXPRESSION TAG	UNP P07236
C	18	PRO	-	EXPRESSION TAG	UNP P07236
C	19	ARG	-	EXPRESSION TAG	UNP P07236

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	20	GLY	-	EXPRESSION TAG	UNP P07236
C	21	SER	-	EXPRESSION TAG	UNP P07236
C	22	HIS	-	EXPRESSION TAG	UNP P07236
C	23	MET	-	EXPRESSION TAG	UNP P07236
C	24	ALA	-	EXPRESSION TAG	UNP P07236
C	25	SER	-	EXPRESSION TAG	UNP P07236
D	3	MET	-	EXPRESSION TAG	UNP P07236
D	4	GLY	-	EXPRESSION TAG	UNP P07236
D	5	SER	-	EXPRESSION TAG	UNP P07236
D	6	SER	-	EXPRESSION TAG	UNP P07236
D	7	HIS	-	EXPRESSION TAG	UNP P07236
D	8	HIS	-	EXPRESSION TAG	UNP P07236
D	9	HIS	-	EXPRESSION TAG	UNP P07236
D	10	HIS	-	EXPRESSION TAG	UNP P07236
D	11	HIS	-	EXPRESSION TAG	UNP P07236
D	12	HIS	-	EXPRESSION TAG	UNP P07236
D	13	SER	-	EXPRESSION TAG	UNP P07236
D	14	SER	-	EXPRESSION TAG	UNP P07236
D	15	GLY	-	EXPRESSION TAG	UNP P07236
D	16	LEU	-	EXPRESSION TAG	UNP P07236
D	17	VAL	-	EXPRESSION TAG	UNP P07236
D	18	PRO	-	EXPRESSION TAG	UNP P07236
D	19	ARG	-	EXPRESSION TAG	UNP P07236
D	20	GLY	-	EXPRESSION TAG	UNP P07236
D	21	SER	-	EXPRESSION TAG	UNP P07236
D	22	HIS	-	EXPRESSION TAG	UNP P07236
D	23	MET	-	EXPRESSION TAG	UNP P07236
D	24	ALA	-	EXPRESSION TAG	UNP P07236
D	25	SER	-	EXPRESSION TAG	UNP P07236

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	18	Total 18	O 18	0	0
3	B	7	Total 7	O 7	0	0
3	C	15	Total 15	O 15	0	0
3	D	6	Total 6	O 6	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Threonyl-tRNA synthetase, mitochondrial

Chain A:



- Molecule 1: Threonyl-tRNA synthetase, mitochondrial

Chain B:





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	154.08Å 157.94Å 237.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.11 – 3.60 35.11 – 3.50	Depositor EDS
% Data completeness (in resolution range)	97.5 (35.11-3.60) 95.3 (35.11-3.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 3.48Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.237 , 0.275 0.226 , 0.267	Depositor DCC
R_{free} test set	1629 reflections (4.96%)	DCC
Wilson B-factor (Å ²)	89.8	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 33.1	EDS
Estimated twinning fraction	0.016 for -k,-h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	2 of 35048 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	13459	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/3494	0.40	0/4729
1	B	0.22	0/3372	0.41	0/4574
1	C	0.23	0/3529	0.41	0/4786
1	D	0.22	0/3380	0.41	0/4590
All	All	0.22	0/13775	0.41	0/18679

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	3
All	All	0	5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	439	ARG	Peptide
1	B	62	LEU	Peptide
1	C	132	ASN	Peptide
1	C	296	GLN	Peptide
1	C	297	LEU	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3397	0	3262	130	0
1	B	3285	0	3069	156	0
1	C	3432	0	3241	136	0
1	D	3295	0	3062	167	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	18	0	0	0	0
3	B	7	0	0	0	0
3	C	15	0	0	0	0
3	D	6	0	0	0	0
All	All	13459	0	12634	556	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 21.

The worst 5 of 556 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:119:THR:H	1:C:120:ASP:HA	1.28	0.98
1:C:296:GLN:HA	1:C:298:PRO:HD2	1.46	0.97
1:B:269:PHE:HB2	1:B:270:TYR:HA	1.50	0.92
1:C:92:PRO:HB3	1:D:58:SER:HB3	1.53	0.90
1:D:40:SER:O	1:D:44:GLN:HB2	1.73	0.87

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/460 (90%)	377 (91%)	37 (9%)	1 (0%)	56	94
1	B	415/460 (90%)	368 (89%)	46 (11%)	1 (0%)	56	94
1	C	426/460 (93%)	390 (92%)	36 (8%)	0	100	100
1	D	412/460 (90%)	370 (90%)	42 (10%)	0	100	100
All	All	1668/1840 (91%)	1505 (90%)	161 (10%)	2 (0%)	59	96

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	394	VAL
1	A	394	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/416 (87%)	361 (100%)	1 (0%)	96	99
1	B	335/416 (80%)	333 (99%)	2 (1%)	92	98
1	C	364/416 (88%)	363 (100%)	1 (0%)	96	99
1	D	338/416 (81%)	338 (100%)	0	100	100
All	All	1399/1664 (84%)	1395 (100%)	4 (0%)	96	99

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	169	LEU
1	B	62	LEU
1	B	341	TRP
1	C	320	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	393	ASN
1	C	132	ASN
1	D	308	GLN
1	B	414	ASN
1	B	429[A]	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	419/460 (91%)	-0.19	0 100 100	64, 83, 110, 132	0
1	B	419/460 (91%)	-0.03	4 (0%) 79 53	64, 98, 138, 149	1 (0%)
1	C	427/460 (92%)	-0.15	0 100 100	57, 79, 105, 115	3 (0%)
1	D	418/460 (90%)	-0.09	0 100 100	64, 93, 126, 137	1 (0%)
All	All	1683/1840 (91%)	-0.12	4 (0%) 93 84	57, 88, 125, 149	5 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	239	LEU	2.5
1	B	237	GLY	2.3
1	B	234	HIS	2.1
1	B	238	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ZN	B	2	1/1	0.26	0.35	92,92,92,92	0
2	ZN	A	1	1/1	0.21	-0.64	70,70,70,70	0
2	ZN	C	463	1/1	0.23	-1.30	59,59,59,59	0
2	ZN	D	463	1/1	0.20	-2.10	103,103,103,103	0

6.5 Other polymers ⓘ

There are no such residues in this entry.