



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 08:26 PM GMT

PDB ID : 3UXH
Title : Design, Synthesis and Biological Evaluation of Potetent Quinoline and Pyrrolo-quinoline Ammosamide Analogues as Inhibitors of Quinone Reductase 2
Authors : Cushman, M.; Mesecar, A.D.; Fanwick, P.E.; Narasimha, R.; Jensen, K.C.
Deposited on : 2011-12-05
Resolution : 1.53 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1031 (1.56-1.52)
Clashscore	79885	1155 (1.56-1.52)
Ramachandran outliers	78287	1127 (1.56-1.52)
Sidechain outliers	78261	1125 (1.56-1.52)
RSRZ outliers	66119	1031 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	230	
1	B	230	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	UXH	A	303	X	-
4	UXH	B	302	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4190 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribosyldihyronicotinamidedehydrogenase [quinone].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	1	0
			1829	1178	304	338	9			
1	B	230	Total	C	N	O	S	0	2	0
			1834	1181	304	340	9			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	PHE	LEU	NATURAL VARIANT	UNP P16083
B	46	PHE	LEU	NATURAL VARIANT	UNP P16083

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is 6,8-DIAMINO-7-CHLORO-1-METHYL-2-OXO-1,2-DIHYDROPYRRR OLO[4,3,2-DE]QUINOLINE-4-CARBOXAMIDE (three-letter code: UXH) (formula: $C_{12}H_{10}ClN_5O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			20	12	1	5	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	Cl	N	O	0	0
			20	12	1	5	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	170	Total	O	0	0
			170	170		
5	B	209	Total	O	0	0
			209	209		


3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribosyldihydronicotinamidedehydrogenase [quinone]



- Molecule 1: Ribosyldihydronicotinamidedehydrogenase [quinone]



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.45Å 84.06Å 106.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.11 – 1.53 39.11 – 1.53	Depositor EDS
% Data completeness (in resolution range)	97.0 (39.11-1.53) 97.0 (39.11-1.53)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.58 (at 1.53Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.162 , 0.184 0.180 , 0.201	Depositor DCC
R_{free} test set	3780 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.087	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 31.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 74965 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4190	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FAD, UXH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.34	4/1882 (0.2%)	1.15	10/2552 (0.4%)
1	B	1.25	2/1890 (0.1%)	1.26	12/2563 (0.5%)
All	All	1.30	6/3772 (0.2%)	1.21	22/5115 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	193	GLU	CD-OE1	6.59	1.32	1.25
1	A	33	GLY	N-CA	5.62	1.54	1.46
1	B	182	LYS	CE-NZ	5.40	1.62	1.49
1	B	193	GLU	CD-OE1	5.29	1.31	1.25
1	A	13	GLU	CD-OE2	5.12	1.31	1.25
1	A	153	GLU	CG-CD	5.04	1.59	1.51

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	78	ARG	NE-CZ-NH1	15.18	127.89	120.30
1	B	78	ARG	NE-CZ-NH2	-12.47	114.06	120.30
1	B	49	ARG	NE-CZ-NH1	9.33	124.96	120.30
1	B	118	ARG	NE-CZ-NH2	-8.98	115.81	120.30
1	B	127[A]	ASP	CB-CG-OD1	8.15	125.64	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	127[B]	ASP	CB-CG-OD1	8.15	125.64	118.30
1	A	104	TYR	CB-CG-CD2	-7.54	116.48	121.00
1	B	52	ASP	CB-CG-OD1	7.34	124.91	118.30
1	B	127[A]	ASP	CB-CG-OD2	-7.33	111.71	118.30
1	B	127[B]	ASP	CB-CG-OD2	-7.33	111.71	118.30
1	A	143	LEU	CB-CG-CD1	-7.07	98.98	111.00
1	A	31	ARG	NE-CZ-NH1	6.80	123.70	120.30
1	A	27	ASP	CB-CG-OD2	-6.44	112.50	118.30
1	A	31	ARG	NE-CZ-NH2	-6.22	117.19	120.30
1	A	65	PHE	CB-CG-CD2	6.19	125.14	120.80
1	B	49	ARG	CG-CD-NE	6.17	124.75	111.80
1	A	184	LEU	CB-CG-CD1	-6.04	100.73	111.00
1	A	101	PHE	CB-CG-CD1	-5.98	116.62	120.80
1	B	86	ASP	CB-CG-OD1	5.20	122.98	118.30
1	A	52	ASP	CB-CG-OD1	5.14	122.93	118.30
1	B	117	ASP	CB-CG-OD1	5.09	122.88	118.30
1	A	1	ALA	C-N-CA	5.06	132.92	122.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	ALA	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1829	0	0	9	0
1	B	1834	0	0	3	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	53	0	31	3	0
3	B	53	0	31	1	0
4	A	20	0	10	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	20	0	10	0	0
5	A	170	0	0	5	0
5	B	209	0	0	2	0
All	All	4190	0	82	14	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (14) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:1:ALA:CB	1:A:34:CYS:SG	2.61	0.89
3:A:302:FAD:H2B	5:A:561:HOH:O	1.88	0.73
1:A:193:GLU:CD	5:A:566:HOH:O	2.31	0.69
3:A:302:FAD:O1A	5:A:566:HOH:O	2.09	0.69
1:A:197:GLU:OE2	1:A:200:ARG:NH1	2.33	0.61
1:B:200:ARG:CD	3:B:303:FAD:O4B	2.51	0.59
1:A:161:ASN:ND2	4:A:303:UXH:HN1C	2.04	0.55
1:A:196:SER:O	1:A:198:GLU:O	2.25	0.54
1:A:193:GLU:CG	5:A:566:HOH:O	2.57	0.53
1:B:45:ASN:OD1	5:B:596:HOH:O	2.19	0.50
1:B:24:VAL:CG1	5:B:493:HOH:O	2.62	0.46
1:A:197:GLU:C	1:A:198:GLU:O	2.55	0.44
1:A:153:GLU:CG	5:A:505:HOH:O	2.64	0.44
1:A:200:ARG:CB	3:A:302:FAD:C2A	2.98	0.42

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	229/230 (100%)	218 (95%)	10 (4%)	1 (0%)	43 16
1	B	230/230 (100%)	222 (96%)	8 (4%)	0	100 100
All	All	459/460 (100%)	440 (96%)	18 (4%)	1 (0%)	56 25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	198	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/194 (100%)	191 (98%)	4 (2%)	66	32
1	B	196/194 (101%)	192 (98%)	4 (2%)	68	34
All	All	391/388 (101%)	383 (98%)	8 (2%)	66	34

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LYS
1	A	157	LYS
1	A	201	LYS
1	A	230	GLN
1	B	53	LYS
1	B	157	LYS
1	B	197	GLU
1	B	230	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	FAD	A	302	-	58,58,58	2.06	15 (25%)	85,89,89	2.10	22 (25%)
4	UXH	A	303	-	22,22,22	2.72	11 (50%)	32,35,35	3.45	14 (43%)
4	UXH	B	302	-	22,22,22	2.88	11 (50%)	32,35,35	3.89	17 (53%)
3	FAD	B	303	-	58,58,58	1.88	13 (22%)	85,89,89	2.14	23 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	302	-	-	0/34/50/50	0/1/6/6
4	UXH	A	303	-	-	4/4/4/4	0/0/3/3
4	UXH	B	302	-	-	0/4/4/4	0/0/3/3
3	FAD	B	303	-	-	0/34/50/50	0/1/6/6

All (50) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302	FAD	C1'-C2'	8.48	1.59	1.51
4	A	303	UXH	C5-C6	5.73	1.43	1.36
3	B	303	FAD	C1'-C2'	5.69	1.57	1.51
3	A	302	FAD	C10-N1	5.57	1.45	1.35
4	A	303	UXH	C9-C8	5.54	1.48	1.40
3	B	303	FAD	C9A-N10	5.39	1.46	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	302	UXH	C5-C6	5.00	1.42	1.36
4	B	302	UXH	C3-C10	-4.97	1.36	1.42
4	B	302	UXH	C9-C10	4.83	1.49	1.42
4	B	302	UXH	C9-C8	4.21	1.46	1.40
4	A	303	UXH	O19-C17	4.20	1.32	1.24
4	A	303	UXH	C7-C8	4.10	1.46	1.40
4	B	302	UXH	O19-C17	4.09	1.32	1.24
4	B	302	UXH	C7-C8	4.08	1.46	1.40
3	A	302	FAD	P-O3P	3.88	1.66	1.59
3	B	303	FAD	C4-N3	3.65	1.43	1.37
3	A	302	FAD	PA-O3P	3.61	1.66	1.59
3	B	303	FAD	C9A-C5X	3.60	1.50	1.42
3	B	303	FAD	P-O3P	3.59	1.66	1.59
4	B	302	UXH	C7-C2	3.44	1.48	1.43
4	A	303	UXH	C9-C10	3.42	1.47	1.42
4	A	303	UXH	C7-C2	3.21	1.48	1.43
3	A	302	FAD	C4X-C10	3.20	1.46	1.40
3	A	302	FAD	C5A-C4A	3.20	1.47	1.40
3	B	303	FAD	C5A-C4A	3.12	1.47	1.40
4	B	302	UXH	C9-N13	3.11	1.42	1.36
3	B	303	FAD	O2-C2	2.99	1.29	1.23
3	A	302	FAD	C10-N10	-2.92	1.32	1.38
3	A	302	FAD	C4-C4X	2.85	1.45	1.41
4	A	303	UXH	C6-C14	-2.78	1.45	1.51
3	B	303	FAD	C4A-N9A	-2.71	1.33	1.37
3	B	303	FAD	C4X-C10	2.67	1.45	1.40
4	A	303	UXH	C2-N1	-2.62	1.31	1.37
4	A	303	UXH	C9-N13	2.60	1.41	1.36
3	A	302	FAD	C4A-N9A	-2.57	1.34	1.37
4	B	302	UXH	C8-CL1	2.46	1.78	1.72
3	A	302	FAD	O2-C2	2.45	1.28	1.23
3	B	303	FAD	O3'-C3'	2.44	1.48	1.43
3	A	302	FAD	C1'-N10	2.28	1.50	1.48
4	B	302	UXH	C10-N18	-2.24	1.36	1.39
4	A	303	UXH	C5-C4	-2.22	1.37	1.42
4	A	303	UXH	C4-C3	-2.20	1.36	1.45
3	A	302	FAD	C2-N1	-2.16	1.30	1.35
3	B	303	FAD	C10-N1	2.15	1.39	1.35
3	A	302	FAD	C9-C9A	-2.14	1.36	1.40
3	A	302	FAD	O4B-C1B	2.13	1.44	1.41
3	A	302	FAD	O4'-C4'	-2.12	1.38	1.43
4	B	302	UXH	C4-C3	-2.08	1.37	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	303	FAD	C8-C7	2.05	1.46	1.40
3	B	303	FAD	C8M-C8	-2.03	1.46	1.51

All (76) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	302	UXH	C4-C3-C2	-14.60	119.92	124.10
4	A	303	UXH	C4-C3-C2	-11.69	120.76	124.10
3	A	302	FAD	N3A-C2A-N1A	-6.87	122.97	128.71
4	A	303	UXH	C14-C6-N1	-6.83	109.33	117.04
4	A	303	UXH	C6-C5-C4	-6.75	115.25	120.50
4	B	302	UXH	C6-C14-N15	6.71	121.80	115.96
3	A	302	FAD	C4X-C10-N1	-6.20	116.53	122.73
3	B	303	FAD	N3A-C2A-N1A	-5.74	123.91	128.71
3	B	303	FAD	C2'-C1'-N10	-5.74	104.84	112.45
3	A	302	FAD	C2'-C1'-N10	-5.52	105.12	112.45
3	A	302	FAD	N3-C2-N1	5.43	132.74	121.19
3	B	303	FAD	N3A-C4A-N9A	5.27	134.95	125.43
4	B	302	UXH	C8-C7-N11	5.18	128.38	121.99
4	B	302	UXH	C10-C3-C2	4.83	130.45	121.04
4	B	302	UXH	C9-C8-C7	4.71	131.14	122.80
4	A	303	UXH	C8-C9-C10	-4.70	112.03	118.11
3	A	302	FAD	C4A-C5A-N7A	-4.70	105.50	109.52
4	B	302	UXH	C8-C7-C2	-4.52	112.27	118.11
3	B	303	FAD	C5X-C9A-N10	-4.47	112.40	116.80
3	B	303	FAD	C4-N3-C2	-4.44	116.28	125.39
3	B	303	FAD	O4B-C1B-N9A	4.36	112.50	108.44
3	A	302	FAD	C4-N3-C2	-4.34	116.47	125.39
3	A	302	FAD	N3A-C4A-N9A	4.34	133.27	125.43
3	A	302	FAD	C1B-N9A-C4A	-4.34	119.14	126.64
4	B	302	UXH	C14-C6-N1	-4.33	112.15	117.04
3	B	303	FAD	O2A-PA-O3P	4.33	125.69	105.14
4	B	302	UXH	C8-C9-C10	-4.24	112.62	118.11
4	A	303	UXH	C10-C3-C2	4.16	129.15	121.04
3	B	303	FAD	C4A-C5A-N7A	-4.11	106.00	109.52
3	B	303	FAD	C8A-N9A-C4A	3.92	109.89	106.90
4	A	303	UXH	O16-C14-C6	3.91	123.56	119.65
3	B	303	FAD	N3-C2-N1	3.81	129.28	121.19
3	B	303	FAD	C4X-C10-N10	3.78	122.39	120.51
4	B	302	UXH	C9-C10-C3	-3.77	117.25	122.71
4	A	303	UXH	C9-C8-C7	3.51	129.02	122.80
4	A	303	UXH	C10-C9-N13	3.49	125.86	121.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	302	UXH	C7-C8-CL1	-3.40	112.80	118.12
4	A	303	UXH	C17-C4-C3	-3.34	102.65	106.36
4	A	303	UXH	C4-C17-N18	3.22	111.88	108.55
4	B	302	UXH	C5-C4-C17	3.20	138.90	132.58
3	A	302	FAD	C2B-C1B-N9A	3.19	121.46	113.27
3	A	302	FAD	O2'-C2'-C1'	3.13	117.48	109.71
3	B	303	FAD	C4B-O4B-C1B	-3.01	106.48	109.75
3	A	302	FAD	N1-C10-N10	2.98	123.82	115.97
4	B	302	UXH	C17-N18-C10	-2.96	103.96	108.59
3	B	303	FAD	C4X-N5-C5X	2.96	120.02	116.69
3	B	303	FAD	C1'-N10-C9A	-2.93	116.02	118.87
3	A	302	FAD	C8A-N9A-C1B	2.92	132.14	126.38
4	B	302	UXH	C17-C4-C3	-2.89	103.15	106.36
3	B	303	FAD	C5A-C4A-N3A	-2.89	119.40	125.70
3	A	302	FAD	C4-C4X-C10	2.77	121.42	116.95
3	A	302	FAD	C5A-C4A-N3A	-2.75	119.71	125.70
3	B	303	FAD	C4X-C10-N1	-2.72	120.02	122.73
3	B	303	FAD	O4'-C4'-C5'	-2.71	104.56	110.12
3	B	303	FAD	C1B-N9A-C4A	-2.64	122.08	126.64
4	B	302	UXH	C10-C9-N13	2.59	124.65	121.14
3	A	302	FAD	O2A-PA-O3P	2.50	117.00	105.14
3	B	303	FAD	C5'-C4'-C3'	-2.50	107.35	112.06
4	A	303	UXH	C9-C10-C3	-2.49	119.09	122.71
3	A	302	FAD	C2A-N3A-C4A	2.46	121.01	114.01
3	A	302	FAD	C1'-N10-C9A	-2.46	116.48	118.87
4	A	303	UXH	C8-C7-C2	-2.44	114.96	118.11
3	A	302	FAD	O4B-C1B-N9A	2.43	110.70	108.44
3	A	302	FAD	C8A-N9A-C4A	2.38	108.72	106.90
3	A	302	FAD	O4'-C4'-C3'	-2.37	103.15	109.05
3	B	303	FAD	O2A-PA-O5B	-2.36	96.60	108.51
3	A	302	FAD	C9A-C5X-N5	2.35	125.97	122.37
4	A	303	UXH	C7-C2-N1	2.35	124.25	119.59
4	B	302	UXH	C4-C17-N18	2.32	110.94	108.55
3	B	303	FAD	C2A-N3A-C4A	2.28	120.49	114.01
4	A	303	UXH	C4-C3-C10	2.24	110.01	106.05
3	B	303	FAD	C9-C9A-N10	2.15	126.22	121.59
4	B	302	UXH	O16-C14-C6	-2.14	117.50	119.65
4	B	302	UXH	C5-C6-C14	2.05	127.47	118.33
3	B	303	FAD	C1'-N10-C10	2.03	122.05	119.17
3	A	302	FAD	C2A-N1A-C6A	2.02	122.41	118.77

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	303	UXH	O16-C14-C6-N1
4	A	303	UXH	O16-C14-C6-C5
4	A	303	UXH	C5-C6-C14-N15
4	A	303	UXH	N1-C6-C14-N15

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/230 (100%)	0.43	22 (9%) 8 7	10, 22, 46, 72	0
1	B	230/230 (100%)	0.20	7 (3%) 48 52	10, 18, 38, 59	0
All	All	460/460 (100%)	0.32	29 (6%) 19 21	10, 20, 42, 72	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	230	GLN	7.3
1	B	2	GLY	6.6
1	B	1	ALA	6.1
1	B	230	GLN	4.2
1	A	57	GLY	4.0
1	A	229	GLY	3.9
1	A	1	ALA	3.9
1	B	197	GLU	3.8
1	A	2	GLY	3.5
1	A	63	GLU	3.4
1	B	198	GLU	3.3
1	B	201	LYS	3.2
1	A	112	LEU	3.1
1	A	158	THR	3.1
1	A	197	GLU	2.9
1	A	58	THR	2.9
1	A	108	VAL	2.8
1	B	112	LEU	2.8
1	A	171	LEU	2.7
1	A	157	LYS	2.6
1	A	153	GLU	2.5
1	A	59	LEU	2.4
1	A	194	ILE	2.3
1	A	225	HIS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	198	GLU	2.2
1	A	196	SER	2.2
1	A	78	ARG	2.2
1	A	101	PHE	2.1
1	A	201	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	FAD	A	302	53/53	0.14	0.40	11,18,62,64	0
3	FAD	B	303	53/53	0.14	0.36	12,17,47,49	0
4	UXH	B	302	20/20	0.08	-0.13	15,19,26,26	0
4	UXH	A	303	20/20	0.07	-0.73	15,18,23,25	0
2	ZN	B	301	1/1	0.03	-2.89	21,21,21,21	0
2	ZN	A	301	1/1	0.04	-3.95	26,26,26,26	0

6.5 Other polymers ⓘ

There are no such residues in this entry.