



wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 03:51 AM GMT

PDB ID : 2UY1
Title : CRYSTAL STRUCTURE OF CSTF-77
Authors : Legrand, P.; Pinaud, N.; Minvielle-Sebastia, L.; Fribourg, S.
Deposited on : 2007-04-02
Resolution : 2.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4888 (2.00-2.00)
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	493	
2	B	493	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7223 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CLEAVAGE STIMULATION FACTOR 77.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	438	Total	C	N	O	S	0	2	0
			3668	2374	601	672	21			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	95	GLU	LEU	CONFLICT	UNP Q8SWC5

- Molecule 2 is a protein called CLEAVAGE STIMULATION FACTOR 77.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	384	Total	C	N	O	S	0	1	0
			3235	2112	530	574	19			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	42	TRP	LYS	CONFLICT	UNP Q8SWC5
B	68	PHE	TYR	CONFLICT	UNP Q8SWC5
B	88	PHE	TYR	CONFLICT	UNP Q8SWC5
B	152	LEU	ILE	CONFLICT	UNP Q8SWC5
B	210	PHE	TYR	CONFLICT	UNP Q8SWC5


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	221	Total	O	0	0
			221	221		
3	B	99	Total	O	0	0
			99	99		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CLEAVAGE STIMULATION FACTOR 77



• Molecule 2: CLEAVAGE STIMULATION FACTOR 77



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.44Å 148.28Å 91.44Å 90.00° 92.15° 90.00°	Depositor
Resolution (Å)	91.29 – 2.00 29.84 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.7 (91.29-2.00) 99.4 (29.84-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.225 , 0.280 0.227 , 0.221	Depositor DCC
R_{free} test set	3954 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	23.8	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 28.7	EDS
Estimated twinning fraction	0.075 for h,-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 79075 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7223	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.16	7/3757 (0.2%)	0.99	11/5041 (0.2%)
2	B	0.98	3/3319 (0.1%)	0.91	3/4455 (0.1%)
All	All	1.08	10/7076 (0.1%)	0.95	14/9496 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8
2	B	0	4
All	All	0	12

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	251	TYR	C-N	-13.68	1.02	1.34
1	A	331	TYR	CD2-CE2	6.55	1.49	1.39
1	A	95[A]	GLU	CA-C	6.08	1.68	1.52
1	A	95[B]	GLU	CA-C	6.08	1.68	1.52
2	B	372	ASN	CG-OD1	5.93	1.36	1.24

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	356	LEU	CA-CB-CG	9.93	138.13	115.30
1	A	404	ARG	NE-CZ-NH2	-8.03	116.28	120.30
1	A	95[A]	GLU	CA-C-O	-6.97	105.46	120.10
1	A	95[B]	GLU	CA-C-O	-6.97	105.46	120.10
1	A	404	ARG	NE-CZ-NH1	6.76	123.68	120.30

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	312	LEU	Peptide
1	A	95[A]	GLU	Mainchain,Peptide
1	A	95[B]	GLU	Mainchain,Peptide
1	A	96	GLY	Peptide
1	A	97	LYS	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3668	0	3621	83	0
2	B	3235	0	3180	98	0
3	A	221	0	0	12	0
3	B	99	0	0	11	0
All	All	7223	0	6801	173	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 173 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:109:ARG:HA	2:B:111:GLY:N	1.37	1.37
2:B:109:ARG:CA	2:B:111:GLY:H	1.37	1.35
2:B:45:TYR:HB2	2:B:207:ASP:OD2	1.42	1.19
1:A:192:ARG:HH11	1:A:192:ARG:HG2	1.09	1.12
2:B:171:SER:HB3	2:B:175:ALA:H	1.08	1.09

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	432/493 (88%)	412 (95%)	17 (4%)	3 (1%)	30	20
2	B	371/493 (75%)	358 (96%)	10 (3%)	3 (1%)	27	17
All	All	803/986 (81%)	770 (96%)	27 (3%)	6 (1%)	27	20

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	100	ASP
1	A	314	ASN
2	B	49	LEU
2	B	110	ASN
2	B	316	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	395/444 (89%)	379 (96%)	16 (4%)	41	35
2	B	346/444 (78%)	340 (98%)	6 (2%)	73	75
All	All	741/888 (83%)	719 (97%)	22 (3%)	53	50

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	315	GLU
1	A	371	GLU
2	B	244	ASP
1	A	365	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	366	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	409	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	438/493 (88%)	-0.13	4 (0%) 81 82	11, 22, 44, 58	0
2	B	384/493 (77%)	0.06	9 (2%) 57 57	12, 31, 52, 61	0
All	All	822/986 (83%)	-0.04	13 (1%) 68 69	11, 26, 50, 61	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	430	VAL	5.5
2	B	42	TRP	4.9
2	B	315	GLU	3.4
1	A	430	VAL	3.2
2	B	68	PHE	3.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.