



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 01:35 AM GMT

PDB ID : 1V08
Title : CRYSTAL STRUCTURE OF THE ZEA MAZE BETA-GLUCOSIDASE-1 IN
COMPLEX WITH GLUCO-TETRAZOLE
Authors : Moriniere, J.; Verdoucq, L.; Bevan, D.R.; Esen, A.; Henrissat, B.; Czjzek, M.
Deposited on : 2004-03-25
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	512	
1	B	512	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NTZ	A	1503	-	X
2	NTZ	B	1503	-	X

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 8829 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BETA-GLUCOSIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	491	Total	C	N	O	S	0	0	1
			3967	2543	657	749	18			
1	B	491	Total	C	N	O	S	0	0	1
			3967	2543	657	749	18			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	191	ASP	GLU	ENGINEERED MUTATION	UNP P49235

- Molecule 2 is NOJIRIMYCINE TETRAZOLE (three-letter code: NTZ) (formula: $C_6H_{10}N_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	6	4	4		
2	A	1	Total	C	N	O	0	0
			14	6	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			14	6	4	4		
2	B	1	Total	C	N	O	0	0
			14	6	4	4		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	450	Total	O	0	0
			450	450		
3	B	389	Total	O	0	0
			389	389		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: BETA-GLUCOSIDASE

Chain A:



• Molecule 1: BETA-GLUCOSIDASE

Chain B:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.18Å 104.80Å 119.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.75 – 1.90 29.69 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.75-1.90) 96.3 (29.69-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.173 , 0.205 0.184 , 0.211	Depositor DCC
R_{free} test set	4604 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	22.9	Xtriage
Anisotropy	0.638	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 40.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 91959 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8829	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NTZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/4088	0.77	14/5551 (0.3%)
1	B	0.43	0/4088	0.75	15/5551 (0.3%)
All	All	0.43	0/8176	0.76	29/11102 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	297	ASP	CB-CG-OD2	7.28	124.85	118.30
1	A	27	ASP	CB-CG-OD2	6.84	124.46	118.30
1	B	54	ASP	CB-CG-OD2	6.80	124.42	118.30
1	B	297	ASP	CB-CG-OD2	6.77	124.39	118.30
1	A	461	ASP	CB-CG-OD2	6.65	124.29	118.30
1	A	411	ASP	CB-CG-OD2	6.62	124.26	118.30
1	B	411	ASP	CB-CG-OD2	6.53	124.17	118.30
1	A	158	ASP	CB-CG-OD2	6.32	123.99	118.30
1	B	158	ASP	CB-CG-OD2	6.29	123.96	118.30
1	B	144	ASP	CB-CG-OD2	6.27	123.94	118.30
1	B	177	ASP	CB-CG-OD2	6.15	123.84	118.30
1	B	27	ASP	CB-CG-OD2	6.12	123.81	118.30
1	B	461	ASP	CB-CG-OD2	6.12	123.80	118.30
1	B	274	ASP	CB-CG-OD2	6.09	123.78	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	144	ASP	CB-CG-OD2	6.05	123.74	118.30
1	A	253	ASP	CB-CG-OD2	6.00	123.70	118.30
1	B	243	ASP	CB-CG-OD2	5.99	123.69	118.30
1	B	355	ASP	CB-CG-OD2	5.88	123.59	118.30
1	B	117	ASP	CB-CG-OD2	5.88	123.59	118.30
1	A	54	ASP	CB-CG-OD2	5.69	123.42	118.30
1	A	479	ASP	CB-CG-OD2	5.57	123.31	118.30
1	B	167	ASP	CB-CG-OD2	5.56	123.30	118.30
1	A	117	ASP	CB-CG-OD2	5.47	123.22	118.30
1	A	46	ASP	CB-CG-OD2	5.40	123.16	118.30
1	A	177	ASP	CB-CG-OD2	5.35	123.11	118.30
1	B	479	ASP	CB-CG-OD2	5.26	123.04	118.30
1	B	252	ASP	CB-CG-OD2	5.24	123.02	118.30
1	A	366	ASP	CB-CG-OD2	5.14	122.93	118.30
1	A	446	ASP	CB-CG-OD2	5.10	122.89	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	458	SER	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3967	0	3757	26	0
1	B	3967	0	3757	23	1
2	A	28	0	20	1	0
2	B	28	0	20	0	0
3	A	450	0	0	14	1
3	B	389	0	0	9	0
All	All	8829	0	7554	48	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including

hydrogens) of the entry. The overall clashscore for this entry is 3.

All (48) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:458:SER:HA	3:B:2355:HOH:O	1.46	1.15
1:B:75:ASN:ND2	3:B:2063:HOH:O	2.25	0.69
1:A:184:LYS:NZ	3:A:2193:HOH:O	2.32	0.63
1:B:73:GLY:O	3:B:2061:HOH:O	2.16	0.62
1:A:75:ASN:ND2	3:A:2082:HOH:O	2.34	0.60
1:A:304:SER:O	1:A:307:ARG:HD3	2.04	0.56
1:A:52:ASN:HD21	1:A:144:ASP:HA	1.71	0.55
1:A:279:GLU:HG3	3:A:2265:HOH:O	2.06	0.55
1:B:90:MET:HE1	1:B:475:ILE:HD12	1.88	0.55
1:B:52:ASN:HD22	1:B:52:ASN:N	2.05	0.54
1:B:79:MET:O	1:B:82:THR:HG22	2.07	0.54
1:B:414:THR:HG23	1:B:417:THR:H	1.72	0.53
1:A:442:LYS:NZ	3:A:2410:HOH:O	2.41	0.53
1:A:123:ARG:HD3	3:A:2185:HOH:O	2.09	0.53
1:A:452:GLN:HG3	3:A:2418:HOH:O	2.09	0.52
1:A:452:GLN:CD	3:A:2418:HOH:O	2.49	0.51
1:B:52:ASN:HD21	1:B:144:ASP:HA	1.76	0.51
1:A:307:ARG:HG3	3:A:2285:HOH:O	2.11	0.50
1:B:457:TRP:O	1:B:458:SER:HB2	2.11	0.50
1:A:90:MET:HE1	1:A:492:ALA:HA	1.93	0.50
1:A:52:ASN:N	1:A:52:ASN:HD22	2.10	0.49
1:A:247:LYS:NZ	3:A:2230:HOH:O	2.45	0.49
1:A:419:LEU:HD13	1:A:424:ALA:HB2	1.93	0.49
1:B:279:GLU:HG3	3:B:2088:HOH:O	2.12	0.49
1:A:295:ARG:HD2	1:B:273:LEU:HD11	1.97	0.46
1:B:480:ARG:HD3	3:B:2061:HOH:O	2.15	0.46
1:A:414:THR:HG23	1:A:417:THR:H	1.81	0.46
1:A:111:GLU:HG2	1:A:112:GLY:N	2.30	0.46
1:A:101:TRP:HB3	1:A:102:PRO:HD3	1.98	0.45
1:A:502:LYS:N	3:A:2447:HOH:O	2.49	0.45
1:A:224:LEU:HD13	1:A:350:VAL:O	2.17	0.45
1:A:82:THR:HG22	3:A:2030:HOH:O	2.18	0.44
1:B:85:ARG:CZ	3:B:2075:HOH:O	2.65	0.44
1:B:320:LYS:HE2	3:B:2177:HOH:O	2.18	0.44
3:A:2282:HOH:O	1:B:307:ARG:CZ	2.64	0.44
1:A:90:MET:HE1	1:A:475:ILE:HD12	2.00	0.43
3:A:2286:HOH:O	1:B:307:ARG:HG3	2.17	0.43
1:A:376:ASN:HB2	1:A:377:PRO:CD	2.49	0.43
1:B:75:ASN:HD21	1:B:78:HIS:HD2	1.67	0.43
1:A:452:GLN:CG	3:A:2418:HOH:O	2.65	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:194:THR:HG21	2:A:1503:NTZ:H2	2.02	0.42
1:B:376:ASN:HB2	1:B:377:PRO:CD	2.49	0.41
1:B:101:TRP:N	1:B:102:PRO:HD2	2.36	0.41
1:A:232:HIS:CE1	1:A:313:PHE:CE2	3.08	0.41
1:B:75:ASN:ND2	3:B:2062:HOH:O	2.54	0.40
1:B:143:TRP:CZ3	1:B:205:PHE:CE2	3.09	0.40
1:B:429:LYS:HD3	3:B:2340:HOH:O	2.22	0.40
1:B:141:PHE:CZ	1:B:143:TRP:HA	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:79:MET:SD	3:A:2413:HOH:O[2_464]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	489/512 (96%)	470 (96%)	18 (4%)	1 (0%)	56	44
1	B	489/512 (96%)	471 (96%)	15 (3%)	3 (1%)	33	19
All	All	978/1024 (96%)	941 (96%)	33 (3%)	4 (0%)	43	29

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	458	SER
1	A	458	SER
1	B	158	ASP
1	B	482	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	423/441 (96%)	402 (95%)	21 (5%)	34	20
1	B	423/441 (96%)	408 (96%)	15 (4%)	48	34
All	All	846/882 (96%)	810 (96%)	36 (4%)	40	26

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	MET
1	A	52	ASN
1	A	70	SER
1	A	75	ASN
1	A	82	THR
1	A	85	ARG
1	A	86	LEU
1	A	111	GLU
1	A	224	LEU
1	A	285	ASN
1	A	330	LEU
1	A	344	SER
1	A	346	ASN
1	A	348	SER
1	A	386	LEU
1	A	419	LEU
1	A	442	LYS
1	A	458	SER
1	A	459	LEU
1	A	462	ASN
1	A	482	ASN
1	B	52	ASN
1	B	86	LEU
1	B	111	GLU
1	B	224	LEU
1	B	252	ASP
1	B	285	ASN
1	B	316	GLU
1	B	328	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	330	LEU
1	B	348	SER
1	B	386	LEU
1	B	419	LEU
1	B	459	LEU
1	B	462	ASN
1	B	482	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	ASN
1	A	132	ASN
1	A	178	ASN
1	A	246	ASN
1	A	276	GLN
1	A	452	GLN
1	A	462	ASN
1	A	481	ASN
1	B	52	ASN
1	B	78	HIS
1	B	132	ASN
1	B	246	ASN
1	B	452	GLN
1	B	462	ASN
1	B	481	ASN
1	B	482	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NTZ	A	1502	-	15,15,15	2.76	5 (33%)	18,22,22	7.99	6 (33%)
2	NTZ	A	1503	-	15,15,15	2.60	5 (33%)	18,22,22	7.97	9 (50%)
2	NTZ	B	1502	-	15,15,15	2.60	5 (33%)	18,22,22	8.44	6 (33%)
2	NTZ	B	1503	-	15,15,15	2.77	5 (33%)	18,22,22	6.08	6 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NTZ	A	1502	-	-	0/2/22/22	0/0/2/2
2	NTZ	A	1503	-	-	0/2/22/22	0/0/2/2
2	NTZ	B	1502	-	-	0/2/22/22	0/0/2/2
2	NTZ	B	1503	-	-	0/2/22/22	0/0/2/2

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1502	NTZ	N21-N18	-6.42	1.25	1.34
2	B	1502	NTZ	N21-N18	-6.10	1.26	1.34
2	B	1503	NTZ	N21-N18	-5.96	1.26	1.34
2	A	1503	NTZ	N21-N18	-5.77	1.26	1.34
2	A	1503	NTZ	N17-N18	5.41	1.41	1.30
2	B	1503	NTZ	N17-N18	5.23	1.41	1.30
2	A	1502	NTZ	C1-C2	-5.01	1.44	1.51
2	A	1502	NTZ	C5-N1	-4.55	1.40	1.49
2	B	1503	NTZ	C1-C2	-4.34	1.45	1.51
2	B	1502	NTZ	N17-N18	4.33	1.39	1.30
2	B	1502	NTZ	C5-N1	-4.26	1.41	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1502	NTZ	C1-C2	-4.18	1.45	1.51
2	A	1502	NTZ	N17-N18	4.13	1.39	1.30
2	B	1503	NTZ	C5-N1	-4.07	1.41	1.49
2	B	1503	NTZ	N17-N1	3.84	1.42	1.34
2	A	1503	NTZ	C1-C2	-3.66	1.46	1.51
2	A	1503	NTZ	N17-N1	3.61	1.41	1.34
2	A	1503	NTZ	C5-N1	-2.97	1.43	1.49
2	B	1502	NTZ	N17-N1	2.86	1.40	1.34
2	A	1502	NTZ	N17-N1	2.68	1.40	1.34

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1502	NTZ	C2-C1-N21	34.30	134.31	129.12
2	A	1502	NTZ	C2-C1-N21	32.41	134.02	129.12
2	A	1503	NTZ	C2-C1-N21	31.30	133.85	129.12
2	B	1503	NTZ	C2-C1-N21	22.97	132.59	129.12
2	B	1503	NTZ	C1-N21-N18	9.90	113.99	105.36
2	B	1502	NTZ	C1-N21-N18	8.23	112.53	105.36
2	A	1503	NTZ	C1-N21-N18	7.98	112.31	105.36
2	A	1502	NTZ	C1-N21-N18	7.32	111.74	105.36
2	A	1503	NTZ	C3-C4-C5	-5.43	102.98	111.62
2	A	1503	NTZ	C1-N1-N17	-4.63	102.33	108.75
2	B	1503	NTZ	C1-N1-N17	-3.74	103.56	108.75
2	A	1502	NTZ	C3-C4-C5	-3.62	105.86	111.62
2	B	1502	NTZ	C3-C4-C5	-3.60	105.90	111.62
2	A	1502	NTZ	C1-N1-N17	-3.45	103.96	108.75
2	B	1502	NTZ	C1-N1-N17	-3.05	104.52	108.75
2	A	1503	NTZ	N21-N18-N17	-3.00	106.64	110.11
2	A	1503	NTZ	O2-C2-C3	2.90	114.80	108.59
2	A	1503	NTZ	C4-C3-C2	-2.84	107.12	110.64
2	B	1503	NTZ	O3-C3-C2	-2.73	103.24	108.61
2	A	1503	NTZ	O4-C4-C5	2.59	114.52	109.80
2	B	1503	NTZ	N21-N18-N17	-2.48	107.24	110.11
2	A	1502	NTZ	O3-C3-C2	-2.31	104.05	108.61
2	B	1502	NTZ	C4-C5-N1	-2.21	103.55	107.81
2	B	1503	NTZ	C4-C3-C2	2.18	113.33	110.64
2	A	1502	NTZ	N21-C1-N1	-2.15	106.66	109.34
2	A	1503	NTZ	O2-C2-C1	2.10	118.14	110.89
2	B	1502	NTZ	N21-N18-N17	-2.04	107.75	110.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	491/512 (95%)	-0.06	14 (2%)	49	50	16, 22, 34, 44	1 (0%)
1	B	491/512 (95%)	0.10	15 (3%)	47	48	16, 26, 37, 44	1 (0%)
All	All	982/1024 (95%)	0.02	29 (2%)	49	49	16, 24, 36, 44	2 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	502	LYS	4.7
1	B	346	ASN	4.2
1	B	501	ALA	4.2
1	A	501	ALA	3.4
1	B	415	LYS	3.2
1	B	416	GLU	3.1
1	A	458	SER	3.0
1	A	346	ASN	3.0
1	B	457	TRP	2.9
1	B	252	ASP	2.8
1	A	502	LYS	2.8
1	A	457	TRP	2.6
1	A	26	SER	2.6
1	A	252	ASP	2.6
1	A	415	LYS	2.5
1	A	416	GLU	2.5
1	B	345	PRO	2.4
1	A	347	TYR	2.4
1	B	111	GLU	2.3
1	B	428	TYR	2.3
1	B	458	SER	2.3
1	A	422	GLU	2.2
1	A	366	ASP	2.1
1	B	12	MET	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	152	LYS	2.1
1	A	12	MET	2.1
1	B	33	ALA	2.1
1	B	253	ASP	2.0
1	A	15	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NTZ	A	1503	14/14	0.28	7.58	39,44,45,46	0
2	NTZ	B	1503	14/14	0.29	4.91	37,42,44,44	0
2	NTZ	A	1502	14/14	0.14	0.36	18,20,21,21	0
2	NTZ	B	1502	14/14	0.14	-0.32	18,20,23,24	0

6.5 Other polymers ⓘ

There are no such residues in this entry.