



# Full wwPDB X-ray Structure Validation Report

Oct 3, 2014 – 12:15 AM EDT

PDB ID : 3V0J  
Title : Crystal structure of Ciona intestinalis voltage sensor-containing phosphatase (Ci-VSP), residues 241-576(C363S), Deletion of 401-405  
Authors : Liu, L.; Kohout, S.C.; Xu, Q.; Muller, S.; Kimberlin, C.; Isacoff, E.Y.; Minor, D.L.  
Deposited on : 2011-12-08  
Resolution : 1.72 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1439  
EDS : stable23828  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable23828

# 1 Overall quality at a glance



The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2979 (1.74-1.70)
Clashscore	79885	3506 (1.74-1.70)
Ramachandran outliers	78287	3449 (1.74-1.70)
Sidechain outliers	78261	3449 (1.74-1.70)
RSRZ outliers	66119	2979 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	334	
1	B	334	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5804 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Voltage-sensor containing phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	10	0
			2634	1675	456	490	13			
1	B	320	Total	C	N	O	S	0	15	0
			2654	1690	454	497	13			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	238	GLY	-	EXPRESSION TAG	UNP Q4W8A1
A	239	HIS	-	EXPRESSION TAG	UNP Q4W8A1
A	240	MET	-	EXPRESSION TAG	UNP Q4W8A1
A	363	SER	CYS	ENGINEERED MUTATION	UNP Q4W8A1
A	?	-	ASP	DELETION	UNP Q4W8A1
A	?	-	PHE	DELETION	UNP Q4W8A1
A	?	-	GLU	DELETION	UNP Q4W8A1
A	?	-	VAL	DELETION	UNP Q4W8A1
A	?	-	GLY	DELETION	UNP Q4W8A1
B	238	GLY	-	EXPRESSION TAG	UNP Q4W8A1
B	239	HIS	-	EXPRESSION TAG	UNP Q4W8A1
B	240	MET	-	EXPRESSION TAG	UNP Q4W8A1
B	363	SER	CYS	ENGINEERED MUTATION	UNP Q4W8A1
B	?	-	ASP	DELETION	UNP Q4W8A1
B	?	-	PHE	DELETION	UNP Q4W8A1
B	?	-	GLU	DELETION	UNP Q4W8A1
B	?	-	VAL	DELETION	UNP Q4W8A1
B	?	-	GLY	DELETION	UNP Q4W8A1

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	275	Total	O	0	0
			275	275		
3	B	231	Total	O	0	0
			231	231		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Voltage-sensor containing phosphatase

Chain A: 



- Molecule 1: Voltage-sensor containing phosphatase

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.56Å 76.80Å 84.01Å 90.00° 92.73° 90.00°	Depositor
Resolution (Å)	83.91 – 1.72 42.39 – 1.72	Depositor EDS
% Data completeness (in resolution range)	99.8 (83.91-1.72) 99.8 (42.39-1.72)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.66 (at 1.72Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.185 , 0.222 0.185 , 0.223	Depositor DCC
$R_{free}$ test set	3441 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.2	Xtriage
Anisotropy	0.257	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 32.4	EDS
Estimated twinning fraction	0.037 for h,-k,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 68067 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5804	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/2723	0.57	0/3668
1	B	0.40	0/2758	0.55	0/3715
All	All	0.41	0/5481	0.56	0/7383

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2634	0	2623	9	0
1	B	2654	0	2656	12	0
2	A	5	0	0	1	0
2	B	5	0	0	1	0
3	A	275	0	0	0	0
3	B	231	0	0	2	0
All	All	5804	0	5279	21	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (21) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:468[A]:ARG:HG2	1:B:468[A]:ARG:HH21	0.92	1.06
1:B:468[A]:ARG:NH2	1:B:468[A]:ARG:HG2	1.62	0.99
1:B:468[A]:ARG:CG	1:B:468[A]:ARG:HH21	1.80	0.94
1:A:342[B]:LYS:NZ	1:A:342[B]:LYS:HB3	2.06	0.70
1:B:283:SER:HB3	1:B:286:ARG:HG2	1.76	0.67
1:A:342[B]:LYS:HZ3	1:A:342[B]:LYS:HB3	1.67	0.59
1:B:468[A]:ARG:NH2	1:B:468[A]:ARG:CG	2.46	0.58
1:B:502:VAL:HG22	1:B:573:ALA:HB1	1.92	0.51
1:A:363:SER:OG	2:A:2:PO4:O1	2.26	0.48
1:B:545:ARG:HD2	1:B:557:TRP:CD1	2.49	0.48
1:A:546:GLU:HG3	3:B:95:HOH:O	2.14	0.47
1:B:281:ARG:H	1:B:281:ARG:NE	2.14	0.46
1:B:254:ARG:HG2	1:B:256:ARG:HG3	1.99	0.45
1:B:391:LEU:HD21	1:B:419:VAL:HG21	1.99	0.45
1:A:434:PRO:HG2	1:A:534[B]:SER:OG	2.17	0.44
1:B:364[B]:LYS:HE3	3:B:655:HOH:O	2.17	0.44
1:A:342[B]:LYS:NZ	1:A:342[B]:LYS:CB	2.78	0.44
1:A:342[B]:LYS:HZ2	1:A:342[B]:LYS:HB3	1.82	0.44
1:B:363:SER:OG	2:B:1:PO4:O1	2.29	0.43
1:A:391:LEU:HD21	1:A:419:VAL:HG21	2.03	0.41
1:A:518:LEU:HD12	1:A:519:PRO:HD2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/334 (98%)	308 (94%)	17 (5%)	2 (1%)	33	13
1	B	333/334 (100%)	321 (96%)	11 (3%)	1 (0%)	50	28
All	All	660/668 (99%)	629 (95%)	28 (4%)	3 (0%)	38	17



All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	550	ASN
1	B	550	ASN
1	A	284	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/297 (99%)	288 (98%)	6 (2%)	68	47
1	B	300/297 (101%)	294 (98%)	6 (2%)	68	47
All	All	594/594 (100%)	582 (98%)	12 (2%)	71	47

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	281	ARG
1	A	286	ARG
1	A	398	ARG
1	A	407	PHE
1	A	558[A]	LYS
1	A	558[B]	LYS
1	B	281	ARG
1	B	282	GLN
1	B	398	ARG
1	B	436	MET
1	B	468[A]	ARG
1	B	468[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	408	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	A	2	-	4,4,4	0.52	0	6,6,6	0.31	0
2	PO4	B	1	-	4,4,4	0.45	0	6,6,6	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	2	-	-	0/0/0/0	0/0/0/0
2	PO4	B	1	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	319/334 (95%)	0.03	14 (4%)	33 37	14, 23, 39, 53	0
1	B	320/334 (95%)	-0.01	12 (3%)	38 43	15, 24, 42, 55	0
All	All	639/668 (95%)	0.01	26 (4%)	35 40	14, 24, 41, 55	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	407	PHE	5.2
1	A	575	ASP	5.1
1	A	281	ARG	5.1
1	A	282	GLN	5.0
1	A	253	ARG	5.0
1	A	284	LEU	4.9
1	A	285	PHE	4.4
1	A	286	ARG	4.2
1	B	281	ARG	4.2
1	B	284	LEU	3.9
1	B	575	ASP	3.7
1	A	283	SER	3.6
1	B	282	GLN	3.4
1	A	400	ASP	3.4
1	A	280	GLY	3.4
1	A	574[A]	GLU	3.1
1	B	574	GLU	3.1
1	B	407	PHE	2.9
1	B	285	PHE	2.7
1	B	400	ASP	2.4
1	A	279	SER	2.3
1	A	572	ASP	2.3
1	B	537	GLU	2.2
1	B	280	GLY	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	573	ALA	2.1
1	B	408	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	PO4	A	2	5/5	0.10	0.30	23,24,26,28	0
2	PO4	B	1	5/5	0.07	-0.39	23,24,26,27	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.