



# wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 02:33 PM GMT

PDB ID : 2V2T  
Title : X-RAY STRUCTURE OF A NF-KB P50-RELB-DNA COMPLEX  
Authors : Moorthy, A.K.; Huang, D.B.; Wang, V.Y.; Vu, D.; Ghosh, G.  
Deposited on : 2007-06-07  
Resolution : 3.05 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2629 (3.12-3.00)
Ramachandran outliers	78287	2536 (3.12-3.00)
Sidechain outliers	78261	2539 (3.12-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	288	
2	B	326	
3	C	11	
3	D	11	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5134 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTION FACTOR RELB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	279	Total	C	N	O	S	0	0	0
			2200	1373	395	417	15			

- Molecule 2 is a protein called NUCLEAR FACTOR NF-KAPPA-B P105 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	313	Total	C	N	O	S	0	0	0
			2458	1557	429	460	12			

- Molecule 3 is a DNA chain called 5'-D(\*CP\*GP\*GP\*GP\*AP\*AP\*TP\*TP\*CP\*CP\*CP)-3'.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	11	Total	C	N	O	P	0	0	0
			221	106	41	64	10			
3	D	11	Total	C	N	O	P	0	0	0
			221	106	41	64	10			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	14	Total	O	0	0
			14	14		
4	B	17	Total	O	0	0
			17	17		
4	C	1	Total	O	0	0
			1	1		
4	D	2	Total	O	0	0
			2	2		




Chain C: 



- Molecule 3: 5'-D(\*CP\*GP\*GP\*GP\*AP\*AP\*TP\*TP\*CP\*CP\*CP)-3'

Chain D: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.44Å 91.44Å 419.97Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.90 – 3.05	Depositor
% Data completeness (in resolution range)	91.6 (19.90-3.05)	Depositor
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.238 , 0.276	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5134	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/2248	0.67	2/3047 (0.1%)
2	B	0.41	0/2510	0.65	0/3391
3	C	0.42	0/247	0.73	0/379
3	D	0.39	0/247	0.70	0/379
All	All	0.40	0/5252	0.67	2/7196 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	285	ARG	CB-CA-C	5.03	120.47	110.40
1	A	330	ASP	CB-CA-C	5.03	120.46	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2200	0	2154	267	0
2	B	2458	0	2451	248	0
3	C	221	0	125	14	0
3	D	221	0	125	9	0
4	A	14	0	0	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	17	0	0	1	0
4	C	1	0	0	0	0
4	D	2	0	0	0	0
All	All	5134	0	4855	518	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 52.

The worst 5 of 518 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:285:ARG:HG2	2:B:267:TYR:OH	1.09	1.24
1:A:110:GLN:HG2	1:A:268:GLU:HB2	1.27	1.16
2:B:207:LEU:HD12	2:B:207:LEU:H	1.05	1.14
1:A:194:VAL:HG13	1:A:195:SER:H	0.96	1.07
1:A:314:VAL:HG12	1:A:323:ARG:HB3	1.33	1.07

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/288 (96%)	192 (69%)	43 (16%)	42 (15%)	0	1
2	B	311/326 (95%)	233 (75%)	46 (15%)	32 (10%)	1	4
All	All	588/614 (96%)	425 (72%)	89 (15%)	74 (13%)	0	3

5 of 74 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	134	SER
1	A	137	GLU
1	A	138	ALA
1	A	164	TRP
1	A	167	TRP



### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/257 (96%)	209 (85%)	37 (15%)	4	19
2	B	268/282 (95%)	245 (91%)	23 (9%)	15	50
All	All	514/539 (95%)	454 (88%)	60 (12%)	8	30

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	309	GLU
1	A	351	GLU
2	B	254	ASP
1	A	340	LYS
1	A	359	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	307	GLN
2	B	43	GLN
2	B	177	GLN
1	A	287	ASN
2	B	185	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.