



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 04:30 PM GMT

PDB ID : 2V3C
Title : CRYSTAL STRUCTURE OF THE SRP54-SRP19-7S.S SRP RNA COM-
PLEX OF M. JANNASCHII
Authors : Hainzl, T.; Huang, S.; Sauer-Eriksson, A.E.
Deposited on : 2007-06-15
Resolution : 2.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)
RNA backbone	1838	1107 (3.10-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	87	
1	B	87	
2	C	432	
2	D	432	
3	M	96	
3	N	96	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12336 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SIGNAL RECOGNITION PARTICLE 19 KDA PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	87	Total	C	N	O	S	0	0	0
			727	468	130	125	4			
1	B	87	Total	C	N	O	S	0	0	0
			727	468	130	125	4			

- Molecule 2 is a protein called SIGNAL RECOGNITION 54 KDA PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	404	Total	C	N	O	S	0	0	1
			3149	1998	541	600	10			
2	D	402	Total	C	N	O	S	0	0	1
			3133	1989	538	597	9			

- Molecule 3 is a RNA chain called 7S RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	96	Total	C	N	O	P	0	0	0
			2063	919	385	664	95			
3	N	96	Total	C	N	O	P	0	0	0
			2063	919	385	664	95			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	23	Total	O	0	0
			23	23		
4	B	26	Total	O	0	0
			26	26		
4	C	107	Total	O	0	0
			107	107		
4	D	107	Total	O	0	0
			107	107		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	M	92	Total 92	O 92	0	0
4	N	119	Total 119	O 119	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SIGNAL RECOGNITION PARTICLE 19 KDA PROTEIN

Chain A: 




• Molecule 1: SIGNAL RECOGNITION PARTICLE 19 KDA PROTEIN


Chain B: 



• Molecule 2: SIGNAL RECOGNITION 54 KDA PROTEIN


Chain C: 






• Molecule 2: SIGNAL RECOGNITION 54 KDA PROTEIN

Chain D:




• Molecule 3: 7S RNA

Chain M:



• Molecule 3: 7S RNA

Chain N:



C202	A203	G204	G205	C206	C207	C208	G209	G210	A211	A212	G213	G214	G215	A216	G217	C218	A219	A220	C221	G222	G223	U224	A225	G226	G227	C228	A229	G230	G231	A232	C233	G234	U235	C236	G237
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.29Å 129.40Å 163.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 32.35 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.6 (20.00-2.50) 93.6 (32.35-2.40)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.244 , 0.294 0.263 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	57.8	Xtriage
Anisotropy	0.510	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 52.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	2 of 55298 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12336	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.19	1/740 (0.1%)	1.26	4/984 (0.4%)
1	B	1.15	1/740 (0.1%)	1.17	6/984 (0.6%)
2	C	0.83	1/3176 (0.0%)	1.05	16/4248 (0.4%)
2	D	0.63	0/3160	0.99	16/4227 (0.4%)
3	M	2.58	140/2309 (6.1%)	3.88	600/3603 (16.7%)
3	N	2.26	91/2309 (3.9%)	3.38	476/3603 (13.2%)
All	All	1.62	234/12434 (1.9%)	2.46	1118/17649 (6.3%)

The worst 5 of 234 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	M	218	C	N1-C6	-14.80	1.28	1.37
3	M	217	G	C8-N7	12.07	1.38	1.30
3	M	217	G	N7-C5	11.67	1.46	1.39
3	M	168	G	N3-C4	-11.48	1.27	1.35
3	N	172	A	N9-C4	11.11	1.44	1.37

The worst 5 of 1118 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	M	209	G	C5-C6-O6	-24.70	113.78	128.60
3	M	194	A	O4'-C1'-N9	-20.43	91.86	108.20
3	M	217	G	C5-C6-O6	-20.20	116.48	128.60
3	N	217	G	C5-C6-O6	-19.78	116.73	128.60
3	M	142	G	O4'-C1'-N9	-18.30	93.56	108.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	727	0	786	80	0
1	B	727	0	786	90	0
2	C	3149	0	3364	579	0
2	D	3133	0	3349	620	0
3	M	2063	0	1043	162	0
3	N	2063	0	1041	136	0
4	A	23	0	0	2	0
4	B	26	0	0	2	0
4	C	107	0	0	17	0
4	D	107	0	0	21	0
4	M	92	0	0	11	0
4	N	119	0	0	4	0
All	All	12336	0	10369	1633	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 75.

The worst 5 of 1633 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:18:ALA:CB	2:D:71:LYS:HD2	1.54	1.34
2:C:325:LYS:HA	2:C:325:LYS:NZ	1.51	1.21
2:D:18:ALA:HB1	2:D:71:LYS:CD	1.71	1.19
2:D:290:ILE:HG23	2:D:294:LEU:HD21	1.22	1.18
2:C:64:THR:HG22	2:C:66:LYS:HB2	1.25	1.15

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	85/87 (98%)	70 (82%)	12 (14%)	3 (4%)	6	7
1	B	85/87 (98%)	66 (78%)	14 (16%)	5 (6%)	2	2
2	C	400/432 (93%)	258 (64%)	97 (24%)	45 (11%)	1	0
2	D	398/432 (92%)	227 (57%)	116 (29%)	55 (14%)	0	0
All	All	968/1038 (93%)	621 (64%)	239 (25%)	108 (11%)	1	0

5 of 108 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	18	ALA
2	C	19	ALA
2	C	24	LYS
2	C	64	THR
2	C	68	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/80 (100%)	69 (86%)	11 (14%)	5	9
1	B	80/80 (100%)	63 (79%)	17 (21%)	1	2
2	C	340/364 (93%)	264 (78%)	76 (22%)	1	2
2	D	338/364 (93%)	253 (75%)	85 (25%)	1	1
All	All	838/888 (94%)	649 (77%)	189 (23%)	1	2

5 of 189 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	303	LEU
2	D	4	LYS
2	D	327	THR
2	C	313	LYS
2	C	334	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	334	GLN
2	C	340	ASN
2	D	101	ASN
2	C	329	ASN
2	D	226	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	M	95/96 (98%)	52 (54%)	16 (16%)
3	N	95/96 (98%)	41 (43%)	7 (7%)
All	All	190/192 (98%)	93 (48%)	23 (12%)

5 of 93 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	M	145	G
3	M	146	G
3	M	147	U
3	M	149	G
3	M	153	A

5 of 23 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
3	M	181	C
3	M	193	G
3	N	193	G
3	M	185	U
3	M	195	A

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	87/87 (100%)	0.35	2 (2%) 57 60	27, 42, 50, 53	0
1	B	87/87 (100%)	0.42	2 (2%) 57 60	27, 42, 53, 62	0
2	C	404/432 (93%)	1.04	66 (16%) 2 2	29, 57, 77, 88	0
2	D	402/432 (93%)	1.73	128 (31%) 1 1	31, 67, 80, 89	0
3	M	96/96 (100%)	-0.18	0 100 100	18, 37, 50, 61	0
3	N	96/96 (100%)	-0.11	0 100 100	20, 37, 52, 70	0
All	All	1172/1230 (95%)	0.98	198 (16%) 2 2	18, 56, 77, 89	0

The worst 5 of 198 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	141	PRO	13.4
2	D	301	SER	12.8
2	D	302	LEU	11.7
2	C	314	THR	11.3
2	D	251	GLY	10.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.