



wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 03:59 AM GMT

PDB ID : 3V4I
Title : Crystal structure of HIV-1 reverse transcriptase (RT) with DNA and AZTTP
Authors : Das, K.; Martinez, S.E.; Arnold, E.
Deposited on : 2011-12-15
Resolution : 2.80 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1799 (2.80-2.80)
Clashscore	79885	2295 (2.80-2.80)
Ramachandran outliers	78287	2252 (2.80-2.80)
Sidechain outliers	78261	2254 (2.80-2.80)
RSRZ outliers	66119	1802 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	556	
1	C	556	
2	B	428	
2	D	428	
3	E	27	
3	T	27	
4	F	21	
4	P	21	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 17696 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 Reverse Transcriptase P66 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	555	Total	C	N	O	S	0	0	0
			4511	2920	751	832	8			
1	C	555	Total	C	N	O	S	0	0	0
			4511	2920	751	832	8			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP P03366
A	0	VAL	-	EXPRESSION TAG	UNP P03366
A	258	CYS	GLN	ENGINEERED MUTATION	UNP P03366
A	280	SER	CYS	ENGINEERED MUTATION	UNP P03366
A	498	ASN	ASP	ENGINEERED MUTATION	UNP P03366
C	-1	MET	-	EXPRESSION TAG	UNP P03366
C	0	VAL	-	EXPRESSION TAG	UNP P03366
C	258	CYS	GLN	ENGINEERED MUTATION	UNP P03366
C	280	SER	CYS	ENGINEERED MUTATION	UNP P03366
C	498	ASN	ASP	ENGINEERED MUTATION	UNP P03366

- Molecule 2 is a protein called HIV-1 Reverse Transcriptase P51 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	412	Total	C	N	O	S	0	0	0
			3400	2212	563	619	6			
2	D	412	Total	C	N	O	S	0	0	0
			3400	2212	563	619	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	280	SER	CYS	ENGINEERED MUTATION	UNP P03366
D	280	SER	CYS	ENGINEERED MUTATION	UNP P03366

- Molecule 3 is a DNA chain called DNA (5'-D(*AP*TP*GP*GP*AP*AP*GP*GP*CP*GP*CP*CP*GP*AP*AP*CP*AP*GP*GP*GP*AP*CP*TP*GP*TP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	T	24	Total	C	N	O	P	0	0	0
			497	234	102	138	23			
3	E	24	Total	C	N	O	P	0	0	0
			497	234	102	138	23			


- Molecule 4 is a DNA chain called DNA (5'-D(*AP*CP*AP*GP*TP*CP*CP*CP*TP*GP*TP*TP*CP*GP*GP*(MRG)P*CP*GP*CP*CP*(ATM))-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	P	20	Total	C	N	O	P	S	0	0	0
			408	195	72	121	19	1			
4	F	20	Total	C	N	O	P	S	0	0	0
			408	195	72	121	19	1			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	C	1	Total	Mg	0	0
			1	1		

- Molecule 6 is 3'-AZIDO-3'-DEOXYTHYMIDINE-5'-TRIPHOSPHATE (three-letter code: AZT) (formula: C₁₀H₁₆N₅O₁₃P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
6	C	1	Total 31	C 10	N 5	O 13	P 3	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HIV-1 Reverse Transcriptase P66 subunit


Chain A: 



- Molecule 1: HIV-1 Reverse Transcriptase P66 subunit


Chain C: 






• Molecule 2: HIV-1 Reverse Transcriptase P51 subunit


Chain B:



• Molecule 2: HIV-1 Reverse Transcriptase P51 subunit

Chain D:





- Molecule 3: DNA (5'-D(*AP*TP*GP*GP*AP*AP*GP*GP*CP*GP*CP*CP*CP*GP*AP*AP*CP*AP*GP*GP*GP*AP*CP*TP*GP*TP*G)-3')

Chain T:




- Molecule 3: DNA (5'-D(*AP*TP*GP*GP*AP*AP*GP*GP*CP*GP*CP*CP*CP*GP*AP*AP*CP*AP*GP*GP*GP*AP*CP*TP*GP*TP*G)-3')

Chain E:




- Molecule 4: DNA (5'-D(*AP*CP*AP*GP*TP*CP*CP*CP*TP*GP*TP*TP*CP*GP*GP*(M RG)P*CP*GP*CP*CP*(ATM))-3')

Chain P:



- Molecule 4: DNA (5'-D(*AP*CP*AP*GP*TP*CP*CP*CP*TP*GP*TP*TP*CP*GP*GP*(M RG)P*CP*GP*CP*CP*(ATM))-3')

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.05Å 132.65Å 138.01Å 90.00° 98.12° 90.00°	Depositor
Resolution (Å)	45.54 – 2.80 45.54 – 2.80	Depositor EDS
% Data completeness (in resolution range)	92.3 (45.54-2.80) 92.3 (45.54-2.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.229 , 0.262 0.222 , 0.258	Depositor DCC
R_{free} test set	2208 reflections (3.02%)	DCC
Wilson B-factor (Å ²)	65.4	Xtriage
Anisotropy	0.370	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 24.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 73180 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	17696	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ATM, MG, MRG, AZT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/4629	0.63	0/6290
1	C	0.45	0/4629	0.60	0/6290
2	B	0.50	0/3497	0.62	0/4751
2	D	0.49	0/3497	0.62	1/4751 (0.0%)
3	E	0.81	0/560	1.58	10/864 (1.2%)
3	T	0.79	0/560	1.55	13/864 (1.5%)
4	F	0.80	0/400	1.55	8/612 (1.3%)
4	P	0.79	0/400	1.70	9/612 (1.5%)
All	All	0.52	0/18172	0.80	41/25034 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

There are no bond length outliers.

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	P	807	DC	O4'-C1'-N1	12.48	116.73	108.00
4	P	820	DC	O4'-C4'-C3'	-10.09	99.95	106.00
4	P	816	DG	O4'-C1'-N9	9.64	114.75	108.00
3	E	718	DA	O4'-C1'-N9	9.52	114.67	108.00
4	F	807	DC	O4'-C1'-N1	9.05	114.34	108.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	420	PRO	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4511	0	4570	217	0
1	C	4511	0	4570	180	0
2	B	3400	0	3433	145	0
2	D	3400	0	3433	142	0
3	E	497	0	268	13	0
3	T	497	0	268	13	0
4	F	408	0	231	19	0
4	P	408	0	231	12	0
5	A	1	0	0	0	0
5	C	1	0	0	0	0
6	A	31	0	12	2	0
6	C	31	0	12	2	0
All	All	17696	0	17028	709	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 20.

The worst 5 of 709 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:115:TYR:HD2	2:D:156:SER:HB3	1.15	1.07
2:D:115:TYR:CD2	2:D:156:SER:HB3	1.91	1.05
4:F:807:DC:H2"	4:F:808:DC:H5"	1.39	1.04
2:B:266:TRP:CD1	2:B:425:LEU:HD22	1.91	1.04
1:A:5:ILE:HG22	1:A:212:TRP:HD1	1.22	1.04

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	553/556 (100%)	524 (95%)	27 (5%)	2 (0%)	43	80
1	C	553/556 (100%)	521 (94%)	30 (5%)	2 (0%)	43	80
2	B	408/428 (95%)	387 (95%)	21 (5%)	0	100	100
2	D	408/428 (95%)	384 (94%)	24 (6%)	0	100	100
All	All	1922/1968 (98%)	1816 (94%)	102 (5%)	4 (0%)	56	88

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	GLU
1	C	217	PRO
1	A	217	PRO
1	C	59	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	495/497 (100%)	472 (95%)	23 (5%)	37	73
1	C	495/497 (100%)	473 (96%)	22 (4%)	39	75
2	B	374/390 (96%)	356 (95%)	18 (5%)	35	72
2	D	374/390 (96%)	360 (96%)	14 (4%)	45	81
All	All	1738/1774 (98%)	1661 (96%)	77 (4%)	39	75

5 of 77 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	269	GLN
1	C	5	ILE
2	D	326	ILE
2	B	280	SER
2	B	424	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	407	GLN
1	C	509	GLN
2	D	336	GLN
1	C	222	GLN
2	D	348	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MRG	F	817	1,3,4	26,28,29	2.62	10 (38%)	33,39,42	2.83	6 (18%)
4	ATM	F	822	3,4	21,23,24	1.17	2 (9%)	24,32,35	2.52	7 (29%)
4	MRG	P	817	1,3,4	26,28,29	2.69	10 (38%)	33,39,42	3.29	6 (18%)
4	ATM	P	822	3,4	21,23,24	1.24	4 (19%)	24,32,35	2.28	6 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MRG	F	817	1,3,4	-	0/11/26/27	0/1/3/3
4	ATM	F	822	3,4	-	0/8/24/25	0/2/2/2
4	MRG	P	817	1,3,4	-	0/11/26/27	0/1/3/3
4	ATM	P	822	3,4	-	0/8/24/25	0/2/2/2

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	P	817	MRG	C4-N3	6.10	1.46	1.35
4	P	817	MRG	C2-N2	5.83	1.49	1.32
4	F	817	MRG	C2-N2	5.63	1.48	1.32
4	F	817	MRG	C4-N3	5.62	1.45	1.35
4	F	817	MRG	P-OP1	5.00	1.52	1.46

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	P	817	MRG	C6-C5-N7	-16.70	131.89	134.14
4	F	817	MRG	C6-C5-N7	-13.75	132.29	134.14
4	F	822	ATM	C6-N1-C2	-8.48	120.00	122.41
4	P	822	ATM	C6-N1-C2	-7.09	120.39	122.41
4	F	822	ATM	N3-C2-N1	5.99	120.97	115.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	AZT	A	823	5	32,32,32	2.50	7 (21%)	44,49,49	2.01	12 (27%)
6	AZT	C	823	5	32,32,32	2.52	7 (21%)	44,49,49	2.14	12 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	AZT	A	823	5	-	0/22/37/37	0/2/2/2
6	AZT	C	823	5	-	0/22/37/37	0/2/2/2

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	823	AZT	O4-C4	8.93	1.42	1.24
6	A	823	AZT	O4-C4	8.75	1.41	1.24
6	A	823	AZT	O2-C2	7.13	1.38	1.23
6	C	823	AZT	O2-C2	7.09	1.38	1.23
6	A	823	AZT	C4-N3	4.30	1.44	1.37

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	823	AZT	C6-N1-C2	-8.19	120.08	122.41
6	A	823	AZT	C6-N1-C2	-6.28	120.62	122.41
6	C	823	AZT	O3B-PB-O3A	5.29	112.42	101.66
6	A	823	AZT	N3-C2-N1	5.14	120.26	115.97
6	C	823	AZT	N3-C2-N1	5.09	120.22	115.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	555/556 (99%)	0.78	74 (13%) 4 3	20, 62, 118, 137	0
1	C	555/556 (99%)	0.78	67 (12%) 5 4	21, 62, 117, 137	0
2	B	412/428 (96%)	0.60	31 (7%) 14 12	21, 51, 109, 122	0
2	D	412/428 (96%)	0.62	42 (10%) 7 6	25, 54, 113, 127	0
3	E	24/27 (88%)	0.20	2 (8%) 11 10	42, 86, 136, 149	0
3	T	24/27 (88%)	0.44	3 (12%) 5 4	40, 80, 137, 146	0
4	F	20/21 (95%)	-0.31	0 100 100	42, 72, 129, 129	0
4	P	20/21 (95%)	0.01	1 (5%) 28 28	42, 69, 130, 131	0
All	All	2022/2064 (97%)	0.68	220 (10%) 6 5	20, 57, 117, 149	0

The worst 5 of 220 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	214	LEU	16.4
2	D	4	PRO	10.7
1	A	133	PRO	10.2
3	T	702	DT	8.4
2	B	4	PRO	8.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ATM	P	822	22/23	0.19	0.64	56,63,78,82	0
4	ATM	F	822	22/23	0.20	0.08	55,64,81,83	0
4	MRG	P	817	26/27	0.17	-0.19	58,74,93,106	0
4	MRG	F	817	26/27	0.13	-0.80	66,81,99,115	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	MG	C	600	1/1	0.24	0.68	82,82,82,82	0
5	MG	A	600	1/1	0.17	-0.23	87,87,87,87	0
6	AZT	A	823	31/31	0.17	-0.54	72,90,106,109	0
6	AZT	C	823	31/31	0.18	-0.75	69,85,104,111	0

6.5 Other polymers

There are no such residues in this entry.