



# wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 04:22 PM GMT

PDB ID : 2V5H  
Title : CONTROLLING THE STORAGE OF NITROGEN AS ARGININE: THE  
COMPLEX OF PII AND ACETYLGLUTAMATE KINASE FROM SYNE-  
CHOCOCCUS ELONGATUS PCC 7942  
Authors : Llacer, J.L.; Marco-Marin, C.; Gil-Ortiz, F.; Fita, I.; Rubio, V.  
Deposited on : 2007-07-04  
Resolution : 2.75 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2995 (2.80-2.72)
Ramachandran outliers	78287	2941 (2.80-2.72)
Sidechain outliers	78261	2944 (2.80-2.72)
RSRZ outliers	66119	2409 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	321	
1	B	321	
1	C	321	
1	D	321	
1	E	321	
1	F	321	
2	G	112	
2	H	112	
2	I	112	
2	J	112	
2	K	112	
2	L	112	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	NA	A	1293	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 18185 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ACETYLGLUTAMATE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	284	Total	C	N	O	S	0	1	0
			2126	1333	386	396	11			
1	B	289	Total	C	N	O	S	0	2	0
			2165	1357	390	407	11			
1	C	284	Total	C	N	O	S	0	1	0
			2128	1334	383	400	11			
1	D	286	Total	C	N	O	S	0	1	0
			2137	1339	385	402	11			
1	E	286	Total	C	N	O	S	0	2	0
			2142	1341	385	405	11			
1	F	287	Total	C	N	O	S	0	1	0
			2142	1342	386	403	11			

- Molecule 2 is a protein called NITROGEN REGULATORY PROTEIN P-II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	111	Total	C	N	O	S	0	0	0
			853	540	149	162	2			
2	H	108	Total	C	N	O	S	0	0	0
			840	531	148	159	2			
2	I	109	Total	C	N	O	S	0	0	0
			838	531	147	158	2			
2	J	110	Total	C	N	O	S	0	0	0
			850	537	150	161	2			
2	K	108	Total	C	N	O	S	0	0	0
			840	531	148	159	2			
2	L	109	Total	C	N	O	S	0	0	0
			841	532	148	159	2			

- Molecule 3 is N-ACETYL-L-GLUTAMATE (three-letter code: NLG) (formula: C<sub>7</sub>H<sub>11</sub>NO<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			13	7	1	5		
3	B	1	Total	C	N	O	0	0
			13	7	1	5		
3	C	1	Total	C	N	O	0	0
			13	7	1	5		
3	D	1	Total	C	N	O	0	0
			13	7	1	5		
3	E	1	Total	C	N	O	0	0
			13	7	1	5		
3	F	1	Total	C	N	O	0	0
			13	7	1	5		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Na	0	0
			1	1		
4	E	1	Total	Na	0	0
			1	1		
4	B	1	Total	Na	0	0
			1	1		
4	C	1	Total	Na	0	0
			1	1		
4	A	1	Total	Na	0	0
			1	1		
4	F	1	Total	Na	0	0
			1	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula:  $Cl$ ).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	2	Total	Cl	0	0
			2	2		
6	G	1	Total	Cl	0	0
			1	1		
6	J	2	Total	Cl	0	0
			2	2		
6	K	1	Total	Cl	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	29	Total	O	0	0
			29	29		
7	B	22	Total	O	0	0
			22	22		
7	C	16	Total	O	0	0
			16	16		
7	D	16	Total	O	0	0
			16	16		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	E	22	Total 22	O 22	0	0
7	F	23	Total 23	O 23	0	0
7	G	9	Total 9	O 9	0	0
7	H	11	Total 11	O 11	0	0
7	I	8	Total 8	O 8	0	0
7	J	10	Total 10	O 10	0	0
7	K	7	Total 7	O 7	0	0
7	L	14	Total 14	O 14	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: ACETYLGLUTAMATE KINASE

Chain A:




#### • Molecule 1: ACETYLGLUTAMATE KINASE

Chain B:



#### • Molecule 1: ACETYLGLUTAMATE KINASE


Chain C:



#### • Molecule 1: ACETYLGLUTAMATE KINASE




Chain D:




- Molecule 1: ACETYLGLUTAMATE KINASE

Chain E:




- Molecule 1: ACETYLGLUTAMATE KINASE

Chain F:



- Molecule 2: NITROGEN REGULATORY PROTEIN P-II

Chain G:



- Molecule 2: NITROGEN REGULATORY PROTEIN P-II

Chain H:




- Molecule 2: NITROGEN REGULATORY PROTEIN P-II

Chain I: 




- Molecule 2: NITROGEN REGULATORY PROTEIN P-II

Chain J: 



- Molecule 2: NITROGEN REGULATORY PROTEIN P-II

Chain K: 



- Molecule 2: NITROGEN REGULATORY PROTEIN P-II

Chain L: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.38Å 161.03Å 91.56Å 90.00° 106.53° 90.00°	Depositor
Resolution (Å)	50.00 – 2.75 45.79 – 2.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-2.75) 100.0 (45.79-2.75)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.16 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.200 , 0.236 0.203 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	45.7	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 15.1	EDS
Estimated twinning fraction	0.022 for l,-k,h	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 65125 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	18185	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, NLG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/2155	0.71	2/2920 (0.1%)
1	B	0.56	0/2196	0.70	2/2979 (0.1%)
1	C	0.52	0/2157	0.69	2/2923 (0.1%)
1	D	0.53	1/2166 (0.0%)	0.65	0/2935
1	E	0.55	0/2171	0.65	0/2943
1	F	0.50	1/2171 (0.0%)	0.61	0/2943
2	G	0.50	0/860	0.66	0/1155
2	H	0.52	0/847	0.71	0/1137
2	I	0.51	0/844	0.67	0/1133
2	J	0.52	0/857	0.70	0/1151
2	K	0.55	0/847	0.72	0/1137
2	L	0.57	0/848	0.71	1/1139 (0.1%)
All	All	0.54	2/18119 (0.0%)	0.68	7/24495 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	2
1	E	0	1
1	F	0	2
All	All	0	5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	218	LYS	CD-CE	6.18	1.66	1.51
1	F	251	CYS	CB-SG	-5.38	1.73	1.81

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	HIS	CB-CA-C	9.76	129.91	110.40
1	C	11	ALA	CB-CA-C	-8.28	97.68	110.10
1	C	10	ALA	N-CA-C	6.89	129.59	111.00
1	B	10	ALA	CB-CA-C	-6.07	100.99	110.10
1	B	10	ALA	N-CA-C	5.95	127.06	111.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	12	ASP	Peptide
1	D	9	GLY	Peptide
1	E	9	GLY	Peptide
1	F	11	ALA	Peptide
1	F	9	GLY	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2126	0	2176	54	0
1	B	2165	0	2190	53	0
1	C	2128	0	2173	50	0
1	D	2137	0	2178	58	0
1	E	2142	0	2172	54	0
1	F	2142	0	2181	48	1
2	G	853	0	892	12	0
2	H	840	0	880	16	0
2	I	838	0	874	16	0
2	J	850	0	887	14	0
2	K	840	0	880	15	1
2	L	841	0	879	8	0
3	A	13	0	9	1	0
3	B	13	0	9	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	13	0	9	1	0
3	D	13	0	9	2	0
3	E	13	0	9	0	0
3	F	13	0	9	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
5	E	6	0	8	1	0
6	G	1	0	0	0	0
6	H	2	0	0	1	0
6	J	2	0	0	1	0
6	K	1	0	0	4	0
7	A	29	0	0	3	0
7	B	22	0	0	0	0
7	C	16	0	0	2	0
7	D	16	0	0	1	0
7	E	22	0	0	5	0
7	F	23	0	0	1	0
7	G	9	0	0	2	0
7	H	11	0	0	2	0
7	I	8	0	0	0	0
7	J	10	0	0	3	0
7	K	7	0	0	1	0
7	L	14	0	0	1	0
All	All	18185	0	18424	357	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

The worst 5 of 357 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:J:38:ARG:HD2	2:J:52:THR:O	1.69	0.93
1:C:12:ASP:N	1:C:13:ARG:HB2	1.85	0.91
1:A:211:ARG:HG2	7:A:2020:HOH:O	1.71	0.90
1:D:211:ARG:HH11	1:D:211:ARG:HG3	1.35	0.88
1:B:11:ALA:HA	1:B:13:ARG:HG3	1.65	0.79

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:F:80:ARG:NH2	2:K:22:ASN:O[1.455]	2.02	0.18

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/321 (88%)	274 (97%)	5 (2%)	4 (1%)	16	45
1	B	289/321 (90%)	277 (96%)	8 (3%)	4 (1%)	16	45
1	C	283/321 (88%)	274 (97%)	7 (2%)	2 (1%)	30	67
1	D	285/321 (89%)	280 (98%)	2 (1%)	3 (1%)	21	53
1	E	286/321 (89%)	280 (98%)	4 (1%)	2 (1%)	30	67
1	F	286/321 (89%)	273 (96%)	9 (3%)	4 (1%)	16	45
2	G	109/112 (97%)	103 (94%)	5 (5%)	1 (1%)	25	60
2	H	106/112 (95%)	102 (96%)	4 (4%)	0	100	100
2	I	105/112 (94%)	101 (96%)	4 (4%)	0	100	100
2	J	108/112 (96%)	101 (94%)	6 (6%)	1 (1%)	25	60
2	K	106/112 (95%)	101 (95%)	4 (4%)	1 (1%)	25	60
2	L	107/112 (96%)	104 (97%)	3 (3%)	0	100	100
All	All	2353/2598 (91%)	2270 (96%)	61 (3%)	22 (1%)	25	60

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	178	ASP
1	F	178	ASP
1	A	10	ALA
1	A	70	GLY
1	A	89	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/256 (88%)	203 (91%)	21 (9%)	13	32
1	B	226/256 (88%)	205 (91%)	21 (9%)	13	33
1	C	225/256 (88%)	203 (90%)	22 (10%)	12	30
1	D	225/256 (88%)	201 (89%)	24 (11%)	10	25
1	E	225/256 (88%)	204 (91%)	21 (9%)	13	33
1	F	225/256 (88%)	200 (89%)	25 (11%)	9	23
2	G	90/93 (97%)	81 (90%)	9 (10%)	11	28
2	H	90/93 (97%)	82 (91%)	8 (9%)	14	36
2	I	88/93 (95%)	81 (92%)	7 (8%)	17	41
2	J	90/93 (97%)	81 (90%)	9 (10%)	11	28
2	K	90/93 (97%)	83 (92%)	7 (8%)	18	42
2	L	89/93 (96%)	82 (92%)	7 (8%)	18	42
All	All	1887/2094 (90%)	1706 (90%)	181 (10%)	12	31

5 of 181 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	221	GLU
1	E	211	ARG
2	K	13	LEU
1	D	261	ARG
1	E	45	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	200	ASN
1	D	26	GLN
1	F	156	ASN
1	B	156	ASN
1	C	156	ASN



### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 12 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	NLG	A	1292	-	12,12,12	1.78	1 (8%)	15,15,15	1.14	1 (6%)
3	NLG	B	1297	-	12,12,12	1.76	1 (8%)	15,15,15	1.03	1 (6%)
3	NLG	C	1292	-	12,12,12	1.76	1 (8%)	15,15,15	1.30	2 (13%)
3	NLG	D	1294	-	12,12,12	1.82	1 (8%)	15,15,15	0.94	0
3	NLG	E	1294	-	12,12,12	1.89	2 (16%)	15,15,15	1.16	2 (13%)
5	GOL	E	1295	-	5,5,5	0.37	0	5,5,5	0.40	0
3	NLG	F	1293	-	12,12,12	1.83	1 (8%)	15,15,15	1.07	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NLG	A	1292	-	-	0/13/13/13	0/0/0/0
3	NLG	B	1297	-	-	0/13/13/13	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NLG	C	1292	-	-	0/13/13/13	0/0/0/0
3	NLG	D	1294	-	-	0/13/13/13	0/0/0/0
3	NLG	E	1294	-	-	0/13/13/13	0/0/0/0
5	GOL	E	1295	-	-	0/4/4/4	0/0/0/0
3	NLG	F	1293	-	-	0/13/13/13	0/0/0/0

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1294	NLG	C8-C7	-5.52	1.39	1.50
3	D	1294	NLG	C8-C7	-5.51	1.39	1.50
3	F	1293	NLG	C8-C7	-5.49	1.39	1.50
3	C	1292	NLG	C8-C7	-5.46	1.39	1.50
3	A	1292	NLG	C8-C7	-5.22	1.39	1.50

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1292	NLG	O7-C7-C8	-2.61	116.94	122.04
3	A	1292	NLG	O7-C7-C8	-2.31	117.53	122.04
3	E	1294	NLG	CB-CA-N2	-2.20	106.64	110.83
3	E	1294	NLG	O7-C7-C8	-2.16	117.83	122.04
3	C	1292	NLG	OE1-CD-CG	-2.06	115.93	123.03

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	284/321 (88%)	0.40	7 (2%) 54 58	31, 37, 46, 50	0
1	B	289/321 (90%)	0.32	6 (2%) 60 64	31, 37, 46, 50	1 (0%)
1	C	284/321 (88%)	0.28	6 (2%) 60 64	31, 37, 45, 50	0
1	D	286/321 (89%)	0.32	9 (3%) 47 50	31, 37, 46, 52	0
1	E	286/321 (89%)	0.34	8 (2%) 50 54	31, 37, 46, 54	1 (0%)
1	F	287/321 (89%)	0.39	11 (3%) 38 41	31, 37, 45, 50	0
2	G	111/112 (99%)	0.56	7 (6%) 19 20	31, 38, 49, 53	0
2	H	108/112 (96%)	0.50	7 (6%) 18 19	31, 37, 46, 56	0
2	I	109/112 (97%)	0.82	13 (11%) 5 5	31, 37, 48, 53	0
2	J	110/112 (98%)	0.38	1 (0%) 81 83	31, 38, 46, 52	0
2	K	108/112 (96%)	0.30	2 (1%) 64 67	31, 38, 46, 55	0
2	L	109/112 (97%)	0.37	2 (1%) 65 68	31, 38, 45, 48	0
All	All	2371/2598 (91%)	0.38	79 (3%) 44 47	31, 37, 46, 56	2 (0%)

The worst 5 of 79 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	293	TYR	7.7
2	G	111	ALA	6.7
2	G	41	GLY	6.2
2	I	109	ALA	6.0
1	F	6	ILE	5.9

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	NA	A	1293	1/1	0.24	2.19	57,57,57,57	0
5	GOL	E	1295	6/6	0.26	1.62	56,56,57,58	0
3	NLG	F	1293	13/13	0.23	1.28	42,46,47,49	0
3	NLG	B	1297	13/13	0.20	0.51	42,44,47,48	0
3	NLG	C	1292	13/13	0.20	0.45	38,39,40,41	0
3	NLG	A	1292	13/13	0.24	0.43	50,53,54,54	0
3	NLG	D	1294	13/13	0.20	0.31	41,45,49,49	0
3	NLG	E	1294	13/13	0.20	0.28	41,43,44,46	0
6	CL	G	1112	1/1	0.21	0.03	43,43,43,43	0
4	NA	D	1295	1/1	0.19	-0.01	50,50,50,50	0
6	CL	J	1112	1/1	0.16	-0.35	62,62,62,62	0
4	NA	B	1298	1/1	0.17	-0.83	37,37,37,37	0
4	NA	F	1294	1/1	0.12	-1.04	47,47,47,47	0
4	NA	E	1296	1/1	0.13	-1.44	46,46,46,46	0
6	CL	H	1110	1/1	0.12	-1.74	63,63,63,63	0
4	NA	C	1293	1/1	0.13	-2.00	36,36,36,36	0
6	CL	H	1109	1/1	0.12	-2.26	35,35,35,35	0
6	CL	J	1111	1/1	0.12	-2.34	30,30,30,30	0
6	CL	K	1109	1/1	0.08	-2.96	39,39,39,39	0

### 6.5 Other polymers ⓘ

There are no such residues in this entry.