



wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 04:23 PM GMT

PDB ID : 2V5M
Title : STRUCTURAL BASIS FOR DSCAM ISOFORM SPECIFICITY
Authors : Meijers, R.; Puettmann-Holgado, R.; Skiniotis, G.; Liu, J.-H.; Walz, T.;
Schmucker, D.; Wang, J.-H.
Deposited on : 2007-07-06
Resolution : 1.95 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1321 (1.96-1.96)
Clashscore	79885	1488 (1.96-1.96)
Ramachandran outliers	78287	1475 (1.96-1.96)
Sidechain outliers	78261	1475 (1.96-1.96)
RSRZ outliers	66119	1321 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	388	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3915 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DSCAM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3144	1980	545	606	13	0	25	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	207	ILE	VAL	VARIANT SEE REMARK 99	UNP Q9NBA1
A	211	ALA	PRO	VARIANT SEE REMARK 99	UNP Q9NBA1
A	213	ARG	LYS	VARIANT SEE REMARK 99	UNP Q9NBA1
A	214	THR	ILE	VARIANT SEE REMARK 99	UNP Q9NBA1
A	215	PRO	ASN	VARIANT SEE REMARK 99	UNP Q9NBA1
A	216	ALA	THR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	218	VAL	THR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	219	GLN	TYR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	222	LEU	ASN	VARIANT SEE REMARK 99	UNP Q9NBA1
A	223	GLU	ILE	VARIANT SEE REMARK 99	UNP Q9NBA1
A	224	LEU	VAL	VARIANT SEE REMARK 99	UNP Q9NBA1
A	225	MET	GLU	VARIANT SEE REMARK 99	UNP Q9NBA1
A	226	VAL	SER	VARIANT SEE REMARK 99	UNP Q9NBA1
A	227	ALA	MET	VARIANT SEE REMARK 99	UNP Q9NBA1
A	228	HIS	ALA	VARIANT SEE REMARK 99	UNP Q9NBA1
A	229	THR	SER	VARIANT SEE REMARK 99	UNP Q9NBA1
A	230	ILE	THR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	231	SER	ALA	VARIANT SEE REMARK 99	UNP Q9NBA1
A	232	LEU	ILE	VARIANT SEE REMARK 99	UNP Q9NBA1
A	239	PHE	TYR	VARIANT SEE REMARK 99	UNP Q9NBA1

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	2	28	16	2	10	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			28	16	2	10		

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	207	ILE	VAL	VARIANT SEE REMARK 99	UNP Q9NBA1
A	211	ALA	PRO	VARIANT SEE REMARK 99	UNP Q9NBA1
A	213	ARG	LYS	VARIANT SEE REMARK 99	UNP Q9NBA1
A	214	THR	ILE	VARIANT SEE REMARK 99	UNP Q9NBA1
A	215	PRO	ASN	VARIANT SEE REMARK 99	UNP Q9NBA1
A	216	ALA	THR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	218	VAL	THR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	219	GLN	TYR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	222	LEU	ASN	VARIANT SEE REMARK 99	UNP Q9NBA1
A	223	GLU	ILE	VARIANT SEE REMARK 99	UNP Q9NBA1
A	224	LEU	VAL	VARIANT SEE REMARK 99	UNP Q9NBA1
A	225	MET	GLU	VARIANT SEE REMARK 99	UNP Q9NBA1
A	226	VAL	SER	VARIANT SEE REMARK 99	UNP Q9NBA1
A	227	ALA	MET	VARIANT SEE REMARK 99	UNP Q9NBA1
A	228	HIS	ALA	VARIANT SEE REMARK 99	UNP Q9NBA1
A	229	THR	SER	VARIANT SEE REMARK 99	UNP Q9NBA1
A	230	ILE	THR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	231	SER	ALA	VARIANT SEE REMARK 99	UNP Q9NBA1
A	232	LEU	ILE	VARIANT SEE REMARK 99	UNP Q9NBA1
A	239	PHE	TYR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	207	ILE	VAL	VARIANT SEE REMARK 99	UNP Q9NBA1
A	211	ALA	PRO	VARIANT SEE REMARK 99	UNP Q9NBA1
A	213	ARG	LYS	VARIANT SEE REMARK 99	UNP Q9NBA1
A	214	THR	ILE	VARIANT SEE REMARK 99	UNP Q9NBA1
A	215	PRO	ASN	VARIANT SEE REMARK 99	UNP Q9NBA1
A	216	ALA	THR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	218	VAL	THR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	219	GLN	TYR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	222	LEU	ASN	VARIANT SEE REMARK 99	UNP Q9NBA1
A	223	GLU	ILE	VARIANT SEE REMARK 99	UNP Q9NBA1
A	224	LEU	VAL	VARIANT SEE REMARK 99	UNP Q9NBA1
A	225	MET	GLU	VARIANT SEE REMARK 99	UNP Q9NBA1
A	226	VAL	SER	VARIANT SEE REMARK 99	UNP Q9NBA1
A	227	ALA	MET	VARIANT SEE REMARK 99	UNP Q9NBA1
A	228	HIS	ALA	VARIANT SEE REMARK 99	UNP Q9NBA1
A	229	THR	SER	VARIANT SEE REMARK 99	UNP Q9NBA1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	230	ILE	THR	VARIANT SEE REMARK 99	UNP Q9NBA1
A	231	SER	ALA	VARIANT SEE REMARK 99	UNP Q9NBA1
A	232	LEU	ILE	VARIANT SEE REMARK 99	UNP Q9NBA1
A	239	PHE	TYR	VARIANT SEE REMARK 99	UNP Q9NBA1

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	709	Total	O	0	0
			709	709		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DSCAM



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 2 2	Depositor
Cell constants a, b, c, α , β , γ	99.17Å 99.17Å 163.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95 29.71 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.6 (20.00-1.95) 98.6 (29.71-1.95)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.47 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.169 , 0.205 0.179 , 0.214	Depositor DCC
R_{free} test set	3006 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	35.0	Xtriage
Anisotropy	0.377	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 61.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 59598 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3915	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	2/3291 (0.1%)	1.35	24/4458 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	386	LYS	C-O	5.77	1.34	1.23
1	A	387	LEU	C-O	5.43	1.33	1.23

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	193	ARG	NE-CZ-NH2	-22.43	109.08	120.30
1	A	193	ARG	NE-CZ-NH1	18.34	129.47	120.30
1	A	198	LYS	CD-CE-NZ	8.24	130.65	111.70
1	A	72	ARG	NE-CZ-NH1	-7.75	116.42	120.30
1	A	85	ARG	NE-CZ-NH2	-7.68	116.46	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	58[A]	ASP	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3144	0	3111	120	0
2	A	56	0	50	3	0
3	A	6	0	6	1	0
4	A	709	0	0	66	6
All	All	3915	0	3167	122	6

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 19.

The worst 5 of 122 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:115:VAL:HB	4:A:2284:HOH:O	1.26	1.26
1:A:14:ILE:HG23	4:A:2037:HOH:O	1.50	1.10
1:A:235:PRO:CD	4:A:2482:HOH:O	2.00	1.07
1:A:57[B]:SER:O	1:A:58[B]:ASP:HB2	1.52	1.07
1:A:235:PRO:CG	4:A:2482:HOH:O	2.04	1.05

The worst 5 of 6 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:A:2185:HOH:O	4:A:2211:HOH:O[8_665]	1.50	0.70
4:A:2400:HOH:O	4:A:2400:HOH:O[2_765]	1.74	0.46
4:A:2029:HOH:O	4:A:2450:HOH:O[5_755]	1.82	0.38
4:A:2429:HOH:O	4:A:2429:HOH:O[6_565]	2.00	0.20
4:A:2279:HOH:O	4:A:2316:HOH:O[5_755]	2.02	0.18

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	411/388 (106%)	398 (97%)	11 (3%)	2 (0%)	38 23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	58[A]	ASP
1	A	58[B]	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	356/332 (107%)	348 (98%)	8 (2%)	64 55

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	193	ARG
1	A	355	LEU
1	A	319	VAL
1	A	124	LYS
1	A	251	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	HIS
1	A	147	ASN
1	A	111	ASN
1	A	31	ASN
1	A	141	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	A	401	1,2	12,14,15	0.82	0	15,19,21	3.14	9 (60%)
2	NAG	A	402	2	12,14,15	0.58	0	15,19,21	2.59	7 (46%)
2	NAG	A	501	1,2	12,14,15	0.57	0	15,19,21	2.07	4 (26%)
2	NAG	A	502	2	12,14,15	0.69	0	15,19,21	1.17	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	401	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	402	2	-	0/6/23/26	0/1/1/1
2	NAG	A	501	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	502	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NAG	O6-C6-C5	-6.84	87.82	111.36
2	A	402	NAG	O5-C5-C6	6.13	113.41	106.98
2	A	401	NAG	C3-C4-C5	5.56	120.12	110.20
2	A	501	NAG	C6-C5-C4	-5.08	100.72	113.00
2	A	401	NAG	C6-C5-C4	-4.34	102.51	113.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	601	-	5,5,5	0.42	0	5,5,5	5.05	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	601	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	GOL	C3-C2-C1	-10.06	66.86	111.26
3	A	601	GOL	O2-C2-C1	4.30	127.83	108.22
3	A	601	GOL	O2-C2-C3	2.27	118.58	108.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/388 (100%)	0.86	49 (12%) 4 5	49, 58, 68, 81	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	388	GLY	6.7
1	A	321	PHE	5.9
1	A	318	THR	4.7
1	A	387	LEU	4.4
1	A	360	VAL	4.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	A	402	14/15	0.21	4.06	59,66,74,80	0
2	NAG	A	502	14/15	0.22	2.14	64,68,83,85	0
2	NAG	A	501	14/15	0.21	1.27	62,69,77,84	0
2	NAG	A	401	14/15	0.12	-0.37	43,53,62,71	0

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	A	601	6/6	0.11	-1.58	40,46,53,53	0

6.5 Other polymers

There are no such residues in this entry.