



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 02:39 PM GMT

PDB ID : 3V6Z
Title : Crystal Structure of Hepatitis B Virus e-antigen
Authors : Dimattia, M.A.; Watts, N.R.; Stahl, S.J.; Grimes, J.M.; Steven, A.C.; Stuart, D.I.; Wingfield, P.T.
Deposited on : 2011-12-20
Resolution : 3.34 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1421 (3.48-3.20)
Clashscore	79885	1077 (3.46-3.22)
Ramachandran outliers	78287	1053 (3.46-3.22)
Sidechain outliers	78261	1052 (3.46-3.22)
RSRZ outliers	66119	1422 (3.48-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	224	
1	C	224	
2	B	219	
2	D	219	
3	E	159	
3	F	159	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9082 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab e6 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S	0	0	0
			1652	1039	265	339	9			
1	C	221	Total	C	N	O	S	0	0	0
			1652	1039	265	339	9			

- Molecule 2 is a protein called Fab e6 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	219	Total	C	N	O	S	0	0	0
			1707	1066	284	348	9			
2	D	219	Total	C	N	O	S	0	0	0
			1707	1066	284	348	9			

- Molecule 3 is a protein called e-antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	148	Total	C	N	O	S	0	0	0
			1182	773	194	211	4			
3	F	148	Total	C	N	O	S	0	0	0
			1182	773	194	211	4			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	58	ALA	CYS	engineered mutation	UNP Q9QMH8
E	74	ASP	GLU	natural variant	UNP Q9QMH8
E	103	VAL	MET	natural variant	UNP Q9QMH8
E	117	ALA	CYS	engineered mutation	UNP Q9QMH8
E	133	ALA	GLY	engineered mutation	UNP Q9QMH8
F	58	ALA	CYS	engineered mutation	UNP Q9QMH8
F	74	ASP	GLU	natural variant	UNP Q9QMH8
F	103	VAL	MET	natural variant	UNP Q9QMH8

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	117	ALA	CYS	engineered mutation	UNP Q9QMH8
F	133	ALA	GLY	engineered mutation	UNP Q9QMH8

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fab e6 Heavy Chain

Chain A: 



- Molecule 1: Fab e6 Heavy Chain

Chain C: 



- Molecule 2: Fab e6 Light Chain

Chain B: 




- Molecule 2: Fab e6 Light Chain

Chain D: 




- Molecule 3: e-antigen

Chain E: 



- Molecule 3: e-antigen

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	66.66Å 75.76Å 88.70Å 96.77° 103.81° 116.04°	Depositor
Resolution (Å)	46.20 – 3.34 46.22 – 3.34	Depositor EDS
% Data completeness (in resolution range)	92.6 (46.20-3.34) 92.9 (46.22-3.34)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 3.32Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.234 , 0.233 0.270 , 0.279	Depositor DCC
R_{free} test set	1003 reflections (5.11%)	DCC
Wilson B-factor (Å ²)	90.7	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 71.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 19651 reflections	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	9082	wwPDB-VP
Average B, all atoms (Å ²)	110.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.43	0/1694	0.79	2/2312 (0.1%)
1	C	0.43	0/1694	0.79	2/2312 (0.1%)
2	B	0.39	0/1746	0.74	0/2369
2	D	0.39	0/1746	0.74	0/2369
3	E	0.43	0/1221	0.63	1/1676 (0.1%)
3	F	0.43	0/1221	0.63	1/1676 (0.1%)
All	All	0.41	0/9322	0.73	6/12714 (0.0%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	11	LEU	CD1-CG-CD2	-6.67	90.49	110.50
1	A	11	LEU	CD1-CG-CD2	-6.34	91.48	110.50
1	C	171	SER	N-CA-C	5.24	125.14	111.00
3	E	93	ASP	CB-CG-OD2	5.23	123.01	118.30
1	A	171	SER	N-CA-C	5.20	125.04	111.00
3	F	93	ASP	CB-CG-OD2	5.19	122.97	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1652	0	0	3	0
1	C	1652	0	0	5	0
2	B	1707	0	2	6	0
2	D	1707	0	2	7	0
3	E	1182	0	18	5	0
3	F	1182	0	18	5	0
All	All	9082	0	40	27	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (27) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:E:87:GLU:O	3:E:92:ARG:HG3	1.43	1.19
3:F:87:GLU:O	3:F:92:ARG:HG3	1.43	1.18
3:E:87:GLU:O	3:E:92:ARG:CG	2.28	0.80
3:F:87:GLU:O	3:F:92:ARG:CG	2.29	0.79
1:C:165:ASN:CB	1:C:168:SER:OG	2.44	0.66
1:A:165:ASN:CB	1:A:168:SER:OG	2.44	0.66
2:B:166:VAL:CG1	2:B:186:LEU:CD1	2.75	0.64
2:D:166:VAL:CG1	2:D:186:LEU:CD1	2.75	0.64
2:B:115:ARG:NE	2:B:116:ALA:O	2.31	0.63
2:D:115:ARG:NE	2:D:116:ALA:O	2.32	0.62
2:B:218:ARG:O	2:B:219:ASN:OD1	2.18	0.61
2:D:218:ARG:O	2:D:219:ASN:OD1	2.19	0.60
1:C:11:LEU:CD2	1:C:157:PRO:CG	2.84	0.56
1:A:11:LEU:CD2	1:A:157:PRO:CG	2.86	0.54
1:A:40:THR:OG1	1:A:42:ASP:OD1	2.29	0.50
1:C:40:THR:OG1	1:C:42:ASP:OD1	2.29	0.50
3:F:93:ASP:O	3:F:93:ASP:OD1	2.34	0.45
3:E:93:ASP:O	3:E:93:ASP:OD1	2.34	0.45
2:B:73:SER:OG	2:D:73:SER:CB	2.66	0.44
2:B:73:SER:CB	2:D:73:SER:OG	2.66	0.44
3:F:4:CYS:SG	3:F:6:GLY:O	2.76	0.42
3:E:4:CYS:SG	3:E:6:GLY:O	2.77	0.42
1:C:11:LEU:CD2	1:C:157:PRO:CB	2.98	0.41
3:E:87:GLU:O	3:E:92:ARG:CD	2.68	0.41
3:F:87:GLU:O	3:F:92:ARG:CD	2.68	0.41
2:B:73:SER:OG	2:D:73:SER:N	2.54	0.41
1:C:114:GLY:O	2:D:49:SER:CB	2.69	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/224 (97%)	215 (99%)	1 (0%)	1 (0%)	38	85
1	C	217/224 (97%)	215 (99%)	1 (0%)	1 (0%)	38	85
2	B	217/219 (99%)	208 (96%)	9 (4%)	0	100	100
2	D	217/219 (99%)	208 (96%)	9 (4%)	0	100	100
3	E	146/159 (92%)	130 (89%)	16 (11%)	0	100	100
3	F	146/159 (92%)	130 (89%)	16 (11%)	0	100	100
All	All	1160/1204 (96%)	1106 (95%)	52 (4%)	2 (0%)	56	94

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	104	GLY
1	C	104	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/192 (100%)	184 (96%)	7 (4%)	45	83
1	C	191/192 (100%)	184 (96%)	7 (4%)	45	83
2	B	196/196 (100%)	181 (92%)	15 (8%)	18	61
2	D	196/196 (100%)	181 (92%)	15 (8%)	18	61
3	E	127/138 (92%)	107 (84%)	20 (16%)	4	20
3	F	127/138 (92%)	107 (84%)	20 (16%)	4	20
All	All	1028/1052 (98%)	944 (92%)	84 (8%)	17	58

All (84) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	LEU
1	A	13	LYS
1	A	64	VAL
1	A	86	LEU
1	A	117	THR
1	A	125	LYS
1	A	160	VAL
2	B	3	MET
2	B	21	MET
2	B	24	LYS
2	B	34	ASN
2	B	35	GLN
2	B	39	LEU
2	B	56	TRP
2	B	59	THR
2	B	112	GLU
2	B	141	CYS
2	B	143	LEU
2	B	154	LYS
2	B	163	GLN
2	B	201	CYS
2	B	210	SER
1	C	11	LEU
1	C	13	LYS
1	C	64	VAL
1	C	86	LEU
1	C	117	THR
1	C	125	LYS
1	C	160	VAL
2	D	3	MET
2	D	21	MET
2	D	24	LYS
2	D	34	ASN
2	D	35	GLN
2	D	39	LEU
2	D	56	TRP
2	D	59	THR
2	D	112	GLU
2	D	141	CYS
2	D	143	LEU
2	D	154	LYS
2	D	163	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	201	CYS
2	D	210	SER
3	E	4	CYS
3	E	8	LEU
3	E	9	TRP
3	E	11	MET
3	E	13	ILE
3	E	26	LEU
3	E	42	ASP
3	E	52	LEU
3	E	56	GLU
3	E	70	LEU
3	E	78	LEU
3	E	86	LEU
3	E	87	GLU
3	E	94	LEU
3	E	110	LEU
3	E	124	THR
3	E	136	ILE
3	E	137	ARG
3	E	138	THR
3	E	146	ASN
3	F	4	CYS
3	F	8	LEU
3	F	9	TRP
3	F	11	MET
3	F	13	ILE
3	F	26	LEU
3	F	42	ASP
3	F	52	LEU
3	F	56	GLU
3	F	70	LEU
3	F	78	LEU
3	F	86	LEU
3	F	87	GLU
3	F	94	LEU
3	F	110	LEU
3	F	124	THR
3	F	136	ILE
3	F	137	ARG
3	F	138	THR
3	F	146	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	221/224 (98%)	0.10	2 (0%) 81 38	59, 89, 134, 157	0
1	C	221/224 (98%)	0.13	2 (0%) 81 38	56, 93, 145, 166	0
2	B	219/219 (100%)	0.21	0 100 100	67, 98, 136, 146	0
2	D	219/219 (100%)	0.19	1 (0%) 88 52	69, 101, 141, 157	0
3	E	148/159 (93%)	0.40	5 (3%) 43 11	64, 149, 201, 226	0
3	F	148/159 (93%)	0.38	5 (3%) 43 11	63, 144, 193, 218	0
All	All	1176/1204 (97%)	0.22	15 (1%) 74 28	56, 103, 169, 226	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	89	PRO	6.7
3	E	89	PRO	4.9
3	E	88	ASP	3.6
1	C	169	LEU	3.5
1	A	169	LEU	3.1
3	F	88	ASP	3.0
3	F	142	TYR	3.0
3	E	142	TYR	2.9
1	C	167	GLY	2.6
3	F	66	ARG	2.3
3	F	92	ARG	2.3
3	E	91	SER	2.2
2	D	163	GLN	2.2
1	A	170	SER	2.2
3	E	90	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.