



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 03:34 AM GMT

PDB ID : 1V7L  
Title : Structure of 3-isopropylmalate isomerase small subunit from *Pyrococcus horikoshii*  
Authors : Yao, M.; Kirita, T.; Sakai, N.; Tanaka, I.  
Deposited on : 2003-12-18  
Resolution : 1.98 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	6577 (2.00-1.96)
Clashscore	79885	8091 (2.00-1.96)
Ramachandran outliers	78287	7989 (2.00-1.96)
Sidechain outliers	78261	7987 (2.00-1.96)
RSRZ outliers	66119	6578 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	163	
1	B	163	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2822 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-isopropylmalate dehydratase small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1258	805	216	234	3			
1	B	163	Total	C	N	O	S	0	0	0
			1270	811	220	236	3			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	157	Total	O	0	0
			157	157		
2	B	137	Total	O	0	0
			137	137		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-isopropylmalate dehydratase small subunit



- Molecule 1: 3-isopropylmalate dehydratase small subunit



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.21 Å 95.21 Å 152.26 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 1.98 24.25 – 1.98	Depositor EDS
% Data completeness (in resolution range)	99.8 (10.00-1.98) 99.7 (24.25-1.98)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.93 (at 1.98 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.206 , 0.247 0.207 , 0.247	Depositor DCC
$R_{free}$ test set	2914 reflections (10.13%)	DCC
Wilson B-factor (Å <sup>2</sup> )	25.3	Xtriage
Anisotropy	0.334	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 43.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 29033 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2822	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/1278	0.72	0/1723
1	B	0.76	2/1290 (0.2%)	1.13	9/1737 (0.5%)
All	All	0.63	2/2568 (0.1%)	0.95	9/3460 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	26	LEU	C-N	-19.05	0.90	1.34
1	B	23	ARG	C-N	7.56	1.51	1.34

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	26	LEU	O-C-N	-19.34	91.75	122.70
1	B	23	ARG	O-C-N	18.25	151.91	122.70
1	B	23	ARG	CA-C-N	-13.39	87.73	117.20
1	B	26	LEU	CA-C-N	12.92	145.63	117.20
1	B	24	TYR	C-N-CA	-6.33	105.86	121.70
1	B	25	ASN	CA-CB-CG	-6.19	99.79	113.40
1	B	25	ASN	C-N-CA	5.81	136.22	121.70
1	B	24	TYR	N-CA-C	5.39	125.55	111.00
1	B	24	TYR	CB-CA-C	-5.32	99.77	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	26	LEU	Mainchain

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1258	0	1291	27	0
1	B	1270	0	1302	74	0
2	A	157	0	0	6	0
2	B	137	0	0	3	0
All	All	2822	0	2593	100	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 20.

All (100) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:25:ASN:C	1:B:26:LEU:HD23	1.72	1.07
1:B:23:ARG:HH21	1:B:32:GLU:HB3	0.98	1.07
1:B:23:ARG:HH21	1:B:32:GLU:CB	1.68	1.04
1:B:23:ARG:NH2	1:B:32:GLU:HB3	1.71	1.03
1:B:23:ARG:HG2	1:B:36:ILE:CD1	1.90	1.02
1:B:24:TYR:CD2	1:B:69:SER:OG	2.14	0.99
1:B:24:TYR:O	1:B:26:LEU:N	1.97	0.98
1:B:26:LEU:N	1:B:26:LEU:CD2	2.25	0.98
1:A:34:ALA:HB1	1:A:77:LEU:HD13	1.49	0.94
1:B:26:LEU:HD23	1:B:26:LEU:N	1.81	0.91
1:B:23:ARG:HG2	1:B:36:ILE:HD11	1.53	0.90
1:B:36:ILE:H	1:B:36:ILE:HD13	1.37	0.87
1:B:24:TYR:CG	1:B:24:TYR:O	2.30	0.83
1:B:89:ARG:H	1:B:89:ARG:NE	1.77	0.82
1:B:24:TYR:HD2	1:B:69:SER:HG	0.87	0.82
1:B:24:TYR:HD2	1:B:69:SER:OG	1.56	0.82
1:B:23:ARG:HD2	1:B:33:LEU:HD23	1.61	0.81
1:B:33:LEU:HA	1:B:36:ILE:HD11	1.64	0.77

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:66:SER:HB2	1:B:90:ILE:HD13	1.70	0.74
1:B:24:TYR:OH	1:B:67:ARG:HB3	1.88	0.73
1:B:23:ARG:HG3	1:B:24:TYR:N	2.03	0.72
1:B:33:LEU:O	1:B:36:ILE:HD13	1.91	0.70
1:B:26:LEU:N	1:B:26:LEU:HD22	2.06	0.69
1:B:24:TYR:O	1:B:25:ASN:C	2.27	0.69
1:B:89:ARG:H	1:B:89:ARG:HE	1.41	0.69
1:B:23:ARG:C	1:B:23:ARG:HD3	2.14	0.68
1:B:23:ARG:HD2	1:B:33:LEU:CD2	2.24	0.66
1:B:20:THR:O	1:B:67:ARG:NH1	2.29	0.65
1:B:23:ARG:CG	1:B:24:TYR:N	2.57	0.65
1:B:36:ILE:HD13	1:B:36:ILE:N	2.08	0.64
1:B:23:ARG:HB2	1:B:23:ARG:NH1	2.13	0.63
1:B:127:LYS:O	1:B:130:GLU:HG2	2.00	0.62
1:A:90:ILE:HD12	1:A:93:ARG:NH2	2.15	0.61
1:A:28:LYS:HB3	1:A:28:LYS:NZ	2.15	0.61
1:B:23:ARG:HH11	1:B:23:ARG:CA	2.14	0.60
1:B:45:PHE:CZ	1:B:49:VAL:HG21	2.37	0.60
1:B:33:LEU:HA	1:B:36:ILE:CD1	2.30	0.59
1:A:71:ALA:O	1:A:100:ILE:HD11	2.03	0.59
1:B:24:TYR:HD2	1:B:69:SER:CB	2.16	0.59
1:B:31:LYS:O	1:B:35:LYS:HD3	2.03	0.58
1:A:47:ARG:HH11	1:A:47:ARG:HG3	1.68	0.58
1:B:23:ARG:C	1:B:23:ARG:HH11	2.07	0.58
1:B:24:TYR:CE1	1:B:67:ARG:CZ	2.87	0.57
1:B:89:ARG:H	1:B:89:ARG:CD	2.18	0.57
1:B:24:TYR:CZ	1:B:67:ARG:HB3	2.40	0.57
1:B:23:ARG:HG2	1:B:36:ILE:CG1	2.34	0.56
1:B:25:ASN:C	1:B:26:LEU:CD2	2.55	0.56
1:B:151:LEU:HD13	1:B:151:LEU:O	2.05	0.56
1:B:45:PHE:CE2	1:B:49:VAL:HG21	2.41	0.55
1:B:24:TYR:CD2	1:B:24:TYR:O	2.58	0.55
1:B:162:ILE:O	1:B:163:ARG:CB	2.55	0.55
1:B:45:PHE:O	1:B:49:VAL:HG23	2.08	0.53
1:A:132:LEU:N	1:A:132:LEU:HD12	2.24	0.53
1:B:66:SER:HA	1:B:90:ILE:HG21	1.90	0.53
1:B:24:TYR:CE1	1:B:25:ASN:ND2	2.77	0.52
1:B:22:GLY:O	1:B:23:ARG:HB3	2.10	0.52
1:B:29:ASP:OD1	1:B:32:GLU:HG2	2.09	0.52
1:B:162:ILE:O	1:B:163:ARG:HB3	2.11	0.51
1:A:94:ASN:O	1:A:98:ILE:HG12	2.11	0.51
1:B:23:ARG:CG	1:B:36:ILE:HG13	2.40	0.50

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:23:ARG:CB	1:B:23:ARG:HH11	2.23	0.50
1:B:24:TYR:CD1	1:B:24:TYR:C	2.79	0.50
1:A:26:LEU:HD22	1:A:26:LEU:N	2.26	0.50
1:A:24:TYR:N	2:A:189:HOH:O	2.45	0.49
1:B:23:ARG:HH21	1:B:32:GLU:CG	2.22	0.49
1:B:23:ARG:CB	1:B:23:ARG:NH1	2.76	0.49
1:A:40:GLU:HG2	2:A:303:HOH:O	2.12	0.49
1:B:36:ILE:CD1	1:B:36:ILE:N	2.75	0.49
1:B:89:ARG:NH2	2:B:222:HOH:O	2.45	0.48
1:B:23:ARG:HG2	1:B:36:ILE:HD12	1.87	0.48
1:A:153:TYR:CE1	1:A:157:ARG:HD2	2.48	0.48
1:A:157:ARG:HG2	1:A:157:ARG:NH1	2.29	0.47
1:B:89:ARG:CD	1:B:89:ARG:N	2.78	0.47
1:A:67:ARG:NH2	2:A:300:HOH:O	2.48	0.46
1:B:23:ARG:HG2	1:B:36:ILE:HG13	1.98	0.46
1:B:93:ARG:HD3	2:B:188:HOH:O	2.15	0.46
1:B:151:LEU:HD13	1:B:151:LEU:C	2.36	0.46
1:B:24:TYR:CE2	1:B:67:ARG:HB3	2.52	0.45
1:A:47:ARG:HH11	1:A:47:ARG:CG	2.28	0.45
1:B:24:TYR:CD1	1:B:67:ARG:NH1	2.84	0.45
1:B:23:ARG:HD3	1:B:27:THR:HG23	1.98	0.45
1:B:24:TYR:CE2	1:B:67:ARG:CB	3.00	0.44
1:A:127:LYS:NZ	2:A:262:HOH:O	2.49	0.44
1:A:157:ARG:HH11	1:A:157:ARG:HG2	1.82	0.44
1:B:60:ASN:HB3	1:B:63:ILE:CG2	2.48	0.43
1:B:23:ARG:HB3	1:B:36:ILE:HG13	1.99	0.43
1:A:23:ARG:C	2:A:189:HOH:O	2.55	0.43
1:A:121:GLU:HA	1:A:142:LEU:CD1	2.48	0.43
1:B:163:ARG:NH1	2:B:190:HOH:O	2.51	0.43
1:A:143:GLU:OE2	1:A:147:GLU:OE2	2.37	0.42
1:A:158:GLY:O	1:A:160:LEU:HG	2.20	0.42
1:A:25:ASN:O	1:A:27:THR:N	2.52	0.42
1:A:67:ARG:NE	2:A:300:HOH:O	2.47	0.41
1:A:115:LEU:HD13	1:B:4:THR:CG2	2.50	0.41
1:B:150:ILE:O	1:B:153:TYR:HB3	2.20	0.41
1:A:39:ILE:HG23	1:A:40:GLU:HG3	2.03	0.41
1:A:28:LYS:HB3	1:A:28:LYS:HZ3	1.85	0.40
1:A:36:ILE:HD12	1:A:36:ILE:C	2.41	0.40
1:A:86:SER:HA	1:A:104:LEU:HD11	2.03	0.40
1:B:24:TYR:CD1	1:B:67:ARG:CZ	3.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/163 (98%)	152 (95%)	6 (4%)	2 (1%)	18	7
1	B	161/163 (99%)	150 (93%)	5 (3%)	6 (4%)	5	1
All	All	321/326 (98%)	302 (94%)	11 (3%)	8 (2%)	9	2

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	23	ARG
1	B	26	LEU
1	B	27	THR
1	B	21	PRO
1	B	28	LYS
1	A	26	LEU
1	B	25	ASN
1	A	158	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/133 (99%)	129 (98%)	3 (2%)	63	61
1	B	133/133 (100%)	123 (92%)	10 (8%)	19	11
All	All	265/266 (100%)	252 (95%)	13 (5%)	35	25

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	77	LEU
1	A	115	LEU
1	B	23	ARG
1	B	25	ASN
1	B	26	LEU
1	B	27	THR
1	B	36	ILE
1	B	67	ARG
1	B	75	LYS
1	B	89	ARG
1	B	115	LEU
1	B	127	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	ASN
1	B	25	ASN
1	B	94	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	162/163 (99%)	0.36	8 (4%) 28 29	14, 23, 46, 74	0
1	B	163/163 (100%)	0.81	15 (9%) 9 9	16, 27, 63, 91	0
All	All	325/326 (99%)	0.58	23 (7%) 16 16	14, 25, 59, 91	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	24	TYR	18.3
1	B	23	ARG	15.3
1	B	26	LEU	9.0
1	B	22	GLY	8.0
1	B	28	LYS	6.1
1	A	158	GLY	5.6
1	B	25	ASN	5.6
1	A	162	ILE	4.9
1	A	159	ASP	3.7
1	B	66	SER	3.6
1	A	131	ILE	3.3
1	B	27	THR	3.1
1	B	65	SER	2.8
1	A	157	ARG	2.7
1	B	162	ILE	2.7
1	B	1	MET	2.6
1	A	129	ASP	2.5
1	B	67	ARG	2.5
1	A	156	ARG	2.4
1	B	55	VAL	2.3
1	B	130	GLU	2.2
1	B	30	PRO	2.1
1	A	22	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.