



Full wwPDB X-ray Structure Validation Report

(i)

Feb 27, 2014 – 01:55 PM GMT

PDB ID : 2V7A
Title : CRYSTAL STRUCTURE OF THE T315I ABL MUTANT IN COMPLEX WITH THE INHIBITOR PHA-739358
Authors : Modugno, M.; Casale, E.; Soncini, C.; Rosettani, P.; Colombo, R.; Lupi, R.; Rusconi, L.; Fancelli, D.; Carpinelli, P.; Cameron, A.D.; Isacchi, A.; Moll, J.
Deposited on : 2007-07-27
Resolution : 2.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance (i)


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.



The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	MG	A	1505	-	X
3	MG	B	1505	-	X

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4638 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROTO-ONCOGENE TYROSINE-PROTEIN KINASE ABL1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	271	Total 2204	C 1417	N 359	O 410	P 1	S 17	0	0	1
1	B	270	Total 2196	C 1413	N 358	O 407	P 1	S 17	0	0	1

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	315	ILE	THR	ENGINEERED MUTATION	UNP P00519
B	315	ILE	THR	ENGINEERED MUTATION	UNP P00519

- Molecule 2 is N-[(3E)-5-[(2R)-2-METHOXY-2-PHENYLACETYL]PYRROLO[3,4-C]PYRAZOL-3(5H)-YLIDENE]-4-(4-METHYLPIPERAZIN-1-YL)BENZAMIDE (three-letter code: 627) (formula: C₂₆H₂₆N₆O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C N O 35 26 6 3	0	0
2	B	1	Total C N O 35 26 6 3	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	104	Total O 104 104	0	0
4	B	62	Total O 62 62	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTO-ONCOGENE TYROSINE-PROTEIN KINASE ABL1

Chain A:



- Molecule 1: PROTO-ONCOGENE TYROSINE-PROTEIN KINASE ABL1

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, α , β , γ	159.76Å 159.76Å 56.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.50 29.56 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-2.50) 99.9 (29.56-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.18 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R , R_{free}	0.207 , 0.246 0.201 , 0.241	Depositor DCC
R_{free} test set	1477 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	34.8	Xtriage
Anisotropy	0.198	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 17.4	EDS
Estimated twinning fraction	0.029 for h,-h-k,-l	Xtriage
L-test for twinning	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Outliers	1 of 29017 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4638	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 627, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/2242	0.73	1/3032 (0.0%)
1	B	0.60	0/2234	0.65	1/3021 (0.0%)
All	All	0.65	0/4476	0.69	2/6053 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	341	LEU	CA-CB-CG	5.50	127.96	115.30
1	A	341	LEU	CA-CB-CG	5.01	126.82	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2204	0	2156	15	0
1	B	2196	0	2152	26	0
2	A	35	0	26	1	0
2	B	35	0	26	1	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
4	A	104	0	0	2	0
4	B	62	0	0	1	0
All	All	4638	0	4360	43	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

All (43) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:318:MET:HE3	1:B:378:LYS:HD2	1.71	0.73
1:A:404:LYS:HE3	1:A:445:LEU:HD23	1.70	0.72
1:A:235:TRP:CD1	1:A:291:LYS:HG2	2.32	0.65
1:B:386:ARG:HE	1:B:393:PTR:P	2.22	0.63
1:B:318:MET:HE1	1:B:378:LYS:HE3	1.82	0.60
1:A:307:ARG:O	1:A:308:GLU:HB3	2.01	0.59
1:A:239:ARG:HB2	1:A:239:ARG:NH1	2.19	0.58
1:A:236:GLU:HG2	4:A:2003:HOH:O	2.07	0.53
1:B:360:ILE:HD13	1:B:388:MET:HE3	1.89	0.53
1:B:406:THR:HG22	1:B:410:SER:HB2	1.90	0.52
1:B:370:LEU:HD22	1:B:380:ALA:CB	2.40	0.51
1:A:363:ASP:OD1	4:A:2050:HOH:O	2.19	0.51
1:B:386:ARG:NE	1:B:393:PTR:O3P	2.44	0.50
1:B:273:LEU:HD23	1:B:311:PHE:HD1	1.78	0.49
1:A:308:GLU:HB2	1:A:309:PRO:HD2	1.94	0.48
1:B:277:THR:HB	1:B:279:GLU:H	1.79	0.47
1:B:388:MET:HE2	1:B:393:PTR:HB3	1.97	0.47
1:B:370:LEU:HD22	1:B:380:ALA:HB2	1.96	0.47
1:B:305:CYS:HB2	1:B:312:TYR:HB2	1.97	0.47
1:A:273:LEU:HD21	1:A:280:VAL:HG22	1.98	0.46
1:A:290:MET:HB3	1:A:301:LEU:HB2	1.97	0.46
1:B:290:MET:HB3	1:B:301:LEU:HB2	1.98	0.46
1:B:268:VAL:HG12	1:B:302:LEU:CD1	2.46	0.46
1:A:293:ILE:HD12	1:A:298:LEU:HD13	1.99	0.45
1:B:283:PHE:HD2	1:B:284:LEU:CD1	2.28	0.45
1:B:361:HIS:O	1:B:362:ARG:HB2	2.17	0.44
1:B:384:LEU:O	1:B:386:ARG:HG2	2.18	0.43
1:B:318:MET:CE	1:B:378:LYS:HE3	2.48	0.43
1:B:316:GLU:HA	4:B:2008:HOH:O	2.18	0.43
1:B:343:MET:HG2	1:B:377:VAL:HG21	2.00	0.43
1:A:474:ALA:HB1	1:A:484:PRO:HD3	2.00	0.43
1:B:268:VAL:HG12	1:B:302:LEU:HD12	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:286:GLU:O	1:B:289:VAL:HG12	2.19	0.42
1:A:237:MET:SD	1:A:314:ILE:HD13	2.59	0.42
1:B:318:MET:HG3	1:B:370:LEU:HB3	2.01	0.42
1:A:279:GLU:HG2	1:A:282:GLU:OE2	2.20	0.41
1:B:297:ASN:O	1:B:378:LYS:HA	2.20	0.41
2:A:1504:627:H10	2:A:1504:627:H182	1.80	0.41
1:A:285:LYS:HA	1:A:285:LYS:HD3	1.96	0.41
1:B:494:GLU:O	1:B:498:GLN:HG2	2.20	0.41
1:B:302:LEU:HB2	1:B:314:ILE:HG22	2.03	0.41
1:A:235:TRP:NE1	1:A:291:LYS:HG2	2.35	0.40
2:B:1504:627:C15	2:B:1504:627:O8	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	268/286 (94%)	255 (95%)	13 (5%)	0	100 100
1	B	267/286 (93%)	248 (93%)	19 (7%)	0	100 100
All	All	535/572 (94%)	503 (94%)	32 (6%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	235/249 (94%)	223 (95%)	12 (5%)	33 57

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	234/249 (94%)	223 (95%)	11 (5%)	36 61
All	All	469/498 (94%)	446 (95%)	23 (5%)	35 59

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	233	ASP
1	A	238	GLU
1	A	265	SER
1	A	276	ASP
1	A	284	LEU
1	A	285	LYS
1	A	308	GLU
1	A	341	LEU
1	A	354	LEU
1	A	370	LEU
1	A	376	LEU
1	A	500	SER
1	B	277	THR
1	B	278	MET
1	B	299	VAL
1	B	308	GLU
1	B	319	THR
1	B	341	LEU
1	B	354	LEU
1	B	370	LEU
1	B	373	GLU
1	B	376	LEU
1	B	438	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	252	GLN
1	B	252	GLN
1	B	358	ASN

5.3.3 RNA

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	PTR	A	393	1	16,16,17	4.46	3 (18%)	20,22,24	1.96	1 (5%)
1	PTR	B	393	1	16,16,17	4.68	3 (18%)	20,22,24	1.52	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	393	1	-	0/9/11/13	0/1/1/1
1	PTR	B	393	1	-	0/9/11/13	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	393	PTR	O-C	17.45	1.23	1.11
1	A	393	PTR	O-C	16.24	1.22	1.11
1	A	393	PTR	OH-CZ	-6.02	1.25	1.40
1	B	393	PTR	OH-CZ	-5.33	1.27	1.40
1	A	393	PTR	P-OH	3.35	1.65	1.60
1	B	393	PTR	P-OH	2.70	1.64	1.60

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	393	PTR	C-CA-N	-8.12	105.72	113.83
1	B	393	PTR	C-CA-N	-5.48	108.36	113.83
1	B	393	PTR	P-OH-CZ	2.26	129.92	123.55
1	B	393	PTR	O3P-P-OH	2.22	114.37	105.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	627	A	1504	-	39,39,39	3.44	13 (33%)	52,55,55	3.42	16 (30%)
2	627	B	1504	-	39,39,39	3.37	12 (30%)	52,55,55	3.13	13 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	627	A	1504	-	-	0/21/45/45	0/3/5/5
2	627	B	1504	-	-	0/21/45/45	0/3/5/5

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1504	627	C18-N17	-12.66	1.27	1.46
2	B	1504	627	C18-N17	-12.03	1.28	1.46
2	B	1504	627	C15-C14	8.68	1.50	1.38
2	A	1504	627	C15-C14	8.45	1.50	1.38
2	B	1504	627	C16-C13	7.16	1.49	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1504	627	C15-N1	6.95	1.50	1.39
2	A	1504	627	C16-C13	6.91	1.49	1.38
2	B	1504	627	C15-N1	6.59	1.49	1.39
2	B	1504	627	C16-N1	5.77	1.48	1.39
2	A	1504	627	C16-N1	5.60	1.48	1.39
2	A	1504	627	C25-N1	-4.02	1.34	1.44
2	B	1504	627	C25-N1	-4.00	1.34	1.44
2	A	1504	627	C19-C18	-3.88	1.34	1.51
2	B	1504	627	C19-C18	-3.85	1.34	1.51
2	A	1504	627	C14-C3	-3.77	1.38	1.47
2	B	1504	627	C14-C3	-3.57	1.39	1.47
2	A	1504	627	N4-N2	3.20	1.42	1.27
2	B	1504	627	N4-N2	3.19	1.42	1.27
2	A	1504	627	C20-C25	-3.01	1.50	1.54
2	B	1504	627	C20-C25	-2.77	1.50	1.54
2	A	1504	627	C13-N2	-2.71	1.35	1.38
2	B	1504	627	C13-N2	-2.57	1.35	1.38
2	A	1504	627	C3-N5	2.35	1.38	1.30
2	B	1504	627	C6-N5	-2.35	1.32	1.37
2	A	1504	627	C6-N5	-2.30	1.32	1.37

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1504	627	C16-N1-C15	17.84	115.35	108.13
2	B	1504	627	C16-N1-C15	16.74	114.90	108.13
2	A	1504	627	C18-C19-N20	9.59	121.40	110.82
2	B	1504	627	C18-C19-N20	7.36	118.94	110.82
2	A	1504	627	N5-C3-N4	-6.14	119.42	129.58
2	B	1504	627	C19-C18-N17	5.49	120.96	110.62
2	B	1504	627	N5-C3-N4	-5.39	120.67	129.58
2	A	1504	627	C21-C22-N17	-4.55	102.04	110.62
2	B	1504	627	C22-N17-C18	4.44	120.80	111.67
2	A	1504	627	C19-C18-N17	3.66	117.51	110.62
2	B	1504	627	C14-C15-N1	-3.64	100.24	111.00
2	A	1504	627	C13-N2-N4	-3.64	107.08	109.23
2	A	1504	627	C22-N17-C18	3.63	119.14	111.67
2	A	1504	627	C14-C15-N1	-3.62	100.30	111.00
2	B	1504	627	C14-C13-N2	-3.54	109.63	112.68
2	B	1504	627	C13-N2-N4	-3.54	107.14	109.23
2	A	1504	627	C14-C13-N2	-3.45	109.71	112.68
2	A	1504	627	C14-C3-N5	3.07	129.50	122.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1504	627	C22-C21-N20	-3.01	107.50	110.82
2	B	1504	627	C7-C6-N5	2.86	119.01	115.10
2	A	1504	627	C21-N20-C19	2.82	113.48	109.54
2	A	1504	627	C11-C12-C7	-2.75	117.47	120.76
2	B	1504	627	C14-C3-N5	2.57	128.44	122.95
2	B	1504	627	C16-N1-C25	-2.50	119.69	125.90
2	A	1504	627	C16-N1-C25	-2.49	119.71	125.90
2	A	1504	627	O26-C25-N1	2.10	122.20	118.55
2	A	1504	627	C6-N5-C3	2.10	126.11	123.12
2	A	1504	627	C12-C7-C9	2.07	121.58	118.63
2	B	1504	627	O26-C25-N1	2.04	122.09	118.55

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	271/286 (94%)	-0.18	6 (2%) 59 61	12, 23, 51, 70	0
1	B	270/286 (94%)	0.16	13 (4%) 29 30	18, 40, 69, 76	0
All	All	541/572 (94%)	-0.01	19 (3%) 42 43	12, 30, 65, 76	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	276	ASP	4.9
1	B	502	ILE	4.8
1	A	278	MET	4.0
1	A	275	GLU	3.4
1	B	396	HIS	3.1
1	B	397	ALA	3.0
1	B	307	ARG	2.9
1	B	399	ALA	2.6
1	B	400	LYS	2.4
1	B	259	GLY	2.3
1	B	398	GLY	2.3
1	A	502	ILE	2.3
1	B	263	LYS	2.2
1	B	261	TRP	2.1
1	B	466	GLU	2.1
1	A	274	LYS	2.1
1	B	429	LEU	2.0
1	A	309	PRO	2.0
1	B	275	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	PTR	B	393	16/17	0.18	-0.19	57,59,63,65	0
1	PTR	A	393	16/17	0.11	-0.46	25,28,40,40	0

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MG	A	1505	1/1	0.25	8.74	32,32,32,32	0
3	MG	B	1505	1/1	0.25	2.45	49,49,49,49	0
2	627	A	1504	35/35	0.13	0.02	13,17,34,35	0
2	627	B	1504	35/35	0.12	-0.53	25,34,49,51	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.