



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 05:29 AM GMT

PDB ID : 3V7D
Title : Crystal Structure of ScSkp1-ScCdc4-pSic1 peptide complex
Authors : Tang, X.; Orlicky, S.; Mittag, T.; Csizmok, V.; Pawson, T.; Forman-Kay, J.;
Sicheri, F.; Tyers, M.
Deposited on : 2011-12-20
Resolution : 2.31 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	169	
1	C	169	
2	B	464	
2	D	464	
3	E	19	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9899 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Suppressor of kinetochore protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	139	Total	C	N	O	S	0	0	0
			1124	708	197	215	4			
1	C	139	Total	C	N	O	S	0	0	0
			1127	711	198	214	4			

There are 62 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP P52286
A	-1	ALA	-	EXPRESSION TAG	UNP P52286
A	0	HIS	-	EXPRESSION TAG	UNP P52286
A	?	-	HIS	DELETION	UNP P52286
A	?	-	ASP	DELETION	UNP P52286
A	?	-	SER	DELETION	UNP P52286
A	?	-	ASN	DELETION	UNP P52286
A	?	-	LEU	DELETION	UNP P52286
A	?	-	GLN	DELETION	UNP P52286
A	?	-	ASN	DELETION	UNP P52286
A	?	-	ASN	DELETION	UNP P52286
A	?	-	SER	DELETION	UNP P52286
A	?	-	ASP	DELETION	UNP P52286
A	?	-	SER	DELETION	UNP P52286
A	?	-	GLU	DELETION	UNP P52286
A	?	-	SER	DELETION	UNP P52286
A	?	-	ASP	DELETION	UNP P52286
A	?	-	SER	DELETION	UNP P52286
A	?	-	ASP	DELETION	UNP P52286
A	?	-	SER	DELETION	UNP P52286
A	?	-	GLU	DELETION	UNP P52286
A	?	-	THR	DELETION	UNP P52286
A	?	-	ASN	DELETION	UNP P52286
A	?	-	HIS	DELETION	UNP P52286
A	?	-	LYS	DELETION	UNP P52286

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	SER	DELETION	UNP P52286
A	?	-	LYS	DELETION	UNP P52286
A	?	-	ASP	DELETION	UNP P52286
A	?	-	ASN	DELETION	UNP P52286
A	?	-	ASN	DELETION	UNP P52286
A	?	-	ASN	DELETION	UNP P52286
C	-2	GLY	-	EXPRESSION TAG	UNP P52286
C	-1	ALA	-	EXPRESSION TAG	UNP P52286
C	0	HIS	-	EXPRESSION TAG	UNP P52286
C	?	-	HIS	DELETION	UNP P52286
C	?	-	ASP	DELETION	UNP P52286
C	?	-	SER	DELETION	UNP P52286
C	?	-	ASN	DELETION	UNP P52286
C	?	-	LEU	DELETION	UNP P52286
C	?	-	GLN	DELETION	UNP P52286
C	?	-	ASN	DELETION	UNP P52286
C	?	-	ASN	DELETION	UNP P52286
C	?	-	SER	DELETION	UNP P52286
C	?	-	ASP	DELETION	UNP P52286
C	?	-	SER	DELETION	UNP P52286
C	?	-	GLU	DELETION	UNP P52286
C	?	-	SER	DELETION	UNP P52286
C	?	-	ASP	DELETION	UNP P52286
C	?	-	SER	DELETION	UNP P52286
C	?	-	ASP	DELETION	UNP P52286
C	?	-	SER	DELETION	UNP P52286
C	?	-	GLU	DELETION	UNP P52286
C	?	-	THR	DELETION	UNP P52286
C	?	-	ASN	DELETION	UNP P52286
C	?	-	HIS	DELETION	UNP P52286
C	?	-	LYS	DELETION	UNP P52286
C	?	-	SER	DELETION	UNP P52286
C	?	-	LYS	DELETION	UNP P52286
C	?	-	ASP	DELETION	UNP P52286
C	?	-	ASN	DELETION	UNP P52286
C	?	-	ASN	DELETION	UNP P52286
C	?	-	ASN	DELETION	UNP P52286

- Molecule 2 is a protein called Cell division control protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	450	Total	C	N	O	S	0	1	0
			3633	2325	628	668	12			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	447	Total	C	N	O	S	0	0	0
			3610	2312	623	663	12			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	261	GLY	-	EXPRESSION TAG	UNP P07834
B	262	ALA	-	EXPRESSION TAG	UNP P07834
B	460	GLU	LYS	VARIANT	UNP P07834
B	?	-	ASN	DELETION	UNP P07834
B	?	-	ILE	DELETION	UNP P07834
B	?	-	TRP	DELETION	UNP P07834
B	?	-	ASN	DELETION	UNP P07834
B	608	LEU	CYS	ENGINEERED MUTATION	UNP P07834
B	?	-	SER	DELETION	UNP P07834
B	?	-	TYR	DELETION	UNP P07834
B	?	-	ALA	DELETION	UNP P07834
B	?	-	THR	DELETION	UNP P07834
B	?	-	ASN	DELETION	UNP P07834
B	?	-	SER	DELETION	UNP P07834
B	?	-	ALA	DELETION	UNP P07834
B	?	-	SER	DELETION	UNP P07834
B	?	-	PRO	DELETION	UNP P07834
B	?	-	CYS	DELETION	UNP P07834
B	?	-	ALA	DELETION	UNP P07834
B	?	-	LYS	DELETION	UNP P07834
B	?	-	ILE	DELETION	UNP P07834
B	?	-	GLY	DELETION	UNP P07834
B	?	-	ALA	DELETION	UNP P07834
D	261	GLY	-	EXPRESSION TAG	UNP P07834
D	262	ALA	-	EXPRESSION TAG	UNP P07834
D	460	GLU	LYS	VARIANT	UNP P07834
D	?	-	ASN	DELETION	UNP P07834
D	?	-	ILE	DELETION	UNP P07834
D	?	-	TRP	DELETION	UNP P07834
D	?	-	ASN	DELETION	UNP P07834
D	608	LEU	CYS	ENGINEERED MUTATION	UNP P07834
D	?	-	SER	DELETION	UNP P07834
D	?	-	TYR	DELETION	UNP P07834
D	?	-	ALA	DELETION	UNP P07834
D	?	-	THR	DELETION	UNP P07834
D	?	-	ASN	DELETION	UNP P07834

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	?	-	SER	DELETION	UNP P07834
D	?	-	ALA	DELETION	UNP P07834
D	?	-	SER	DELETION	UNP P07834
D	?	-	PRO	DELETION	UNP P07834
D	?	-	CYS	DELETION	UNP P07834
D	?	-	ALA	DELETION	UNP P07834
D	?	-	LYS	DELETION	UNP P07834
D	?	-	ILE	DELETION	UNP P07834
D	?	-	GLY	DELETION	UNP P07834
D	?	-	ALA	DELETION	UNP P07834

- Molecule 3 is a protein called Protein SIC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	10	Total	C	N	O	P	0	0	0
			85	47	15	21	2			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	25	Total	O	0	0
			25	25		
4	B	125	Total	O	0	0
			125	125		
4	C	23	Total	O	0	0
			23	23		
4	D	147	Total	O	0	0
			147	147		

4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	107.29Å 107.29Å 166.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.02 – 2.31 47.69 – 2.31	Depositor EDS
% Data completeness (in resolution range)	99.6 (38.02-2.31) 99.6 (47.69-2.31)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.32Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.196 , 0.221 0.187 , 0.211	Depositor DCC
R_{free} test set	4732 reflections (5.02%)	DCC
Wilson B-factor (Å ²)	41.7	Xtriage
Anisotropy	0.233	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 34.4	EDS
Estimated twinning fraction	0.013 for -h,-k,l 0.025 for h,-h-k,-l 0.013 for -k,-h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 94284 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9899	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/1142	0.43	0/1544
1	C	0.27	0/1145	0.42	0/1546
2	B	0.25	0/3718	0.45	0/5032
2	D	0.27	0/3692	0.48	0/4996
3	E	0.28	0/65	0.43	0/84
All	All	0.26	0/9762	0.46	0/13202

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1124	0	0	1	0
1	C	1127	0	0	2	0
2	B	3633	0	0	7	0
2	D	3610	0	0	3	0
3	E	85	0	4	1	0
4	A	25	0	0	1	0
4	B	125	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	23	0	0	0	0
4	D	147	0	0	1	0
All	All	9899	0	4	13	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

The worst 5 of 13 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:485:ARG:NH2	4:D:928:HOH:O	2.28	0.67
2:B:508:ASP:N	2:B:508:ASP:OD1	2.29	0.66
2:B:464:SER:OG	3:E:80:SEP:O3P	2.17	0.62
2:B:433:GLY:O	2:B:451:LYS:NZ	2.37	0.58
1:A:83:ARG:NH2	4:A:208:HOH:O	2.37	0.57

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/169 (79%)	130 (98%)	2 (2%)	1 (1%)	27	30
1	C	133/169 (79%)	129 (97%)	2 (2%)	2 (2%)	15	13
2	B	447/464 (96%)	435 (97%)	12 (3%)	0	100	100
2	D	443/464 (96%)	430 (97%)	12 (3%)	1 (0%)	56	68
3	E	7/19 (37%)	7 (100%)	0	0	100	100
All	All	1163/1285 (90%)	1131 (97%)	28 (2%)	4 (0%)	50	60

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	187	GLU
2	D	432	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	114	ALA
1	C	115	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/152 (83%)	125 (99%)	1 (1%)	89	96
1	C	126/152 (83%)	124 (98%)	2 (2%)	75	88
2	B	407/416 (98%)	400 (98%)	7 (2%)	73	87
2	D	403/416 (97%)	400 (99%)	3 (1%)	91	97
3	E	7/15 (47%)	7 (100%)	0	100	100
All	All	1069/1151 (93%)	1056 (99%)	13 (1%)	82	92

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	508	ASP
2	B	600	GLU
2	D	335	ASN
2	B	500	VAL
1	C	112	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SEP	E	76	3	9,9,10	6.13	3 (33%)	10,12,14	1.62	2 (20%)
3	SEP	E	80	3	9,9,10	6.18	3 (33%)	10,12,14	1.64	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SEP	E	76	3	-	0/6/8/10	0/0/0/0
3	SEP	E	80	3	-	0/6/8/10	0/0/0/0

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	80	SEP	O-C	17.89	1.23	1.11
3	E	76	SEP	O-C	17.73	1.23	1.11
3	E	76	SEP	P-O1P	3.24	1.62	1.51
3	E	80	SEP	P-O1P	3.04	1.61	1.51
3	E	80	SEP	CA-C	2.60	1.53	1.48

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	80	SEP	C-CA-N	-3.47	110.36	113.83
3	E	76	SEP	P-OG-CB	-3.40	108.35	118.19
3	E	76	SEP	C-CA-N	-3.07	110.77	113.83
3	E	80	SEP	P-OG-CB	-2.78	110.15	118.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	139/169 (82%)	-0.09	5 (3%) 41 51	32, 47, 99, 124	0
1	C	139/169 (82%)	-0.09	8 (5%) 22 31	32, 44, 98, 118	0
2	B	450/464 (96%)	-0.17	8 (1%) 65 74	28, 45, 75, 124	0
2	D	447/464 (96%)	-0.32	7 (1%) 68 77	25, 37, 58, 128	0
3	E	10/19 (52%)	1.05	2 (20%) 2 3	68, 76, 116, 140	0
All	All	1185/1285 (92%)	-0.20	30 (2%) 54 65	25, 42, 76, 140	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	103	PRO	6.9
2	B	500	VAL	5.9
2	B	498	SER	4.3
1	A	114	ALA	4.3
1	A	113	SER	4.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	SEP	E	80	10/11	0.28	0.73	65,86,115,180	0
3	SEP	E	76	10/11	0.12	-0.63	62,70,74,76	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.